

# Abdominal Aortic Aneurysm

Wall degeneration • morphology • complications

Anatomically grounded clinicopathologic interpretation

**Dr Abu Suraih Sakhri**

MBBS, MD (Pathology)


[www.abusuraihsakhri.com](http://www.abusuraihsakhri.com)

## Learning objectives

- Distinguish a true abdominal aortic aneurysm from pseudoaneurysm and dissection.
- Connect atherosclerosis, inflammation, extracellular matrix loss and medial injury.
- Recognize the morphology and inflammatory, IgG4-related and mycotic variants.
- Explain rupture, ischemia, embolism and compression as clinical outcomes.

# AAA is a true aneurysm

- An aneurysm is an abnormal, localized vessel dilation.
- In a true aneurysm, attenuated but continuous wall layers bound the dilation.
- Atherosclerotic aneurysms arise particularly in the abdominal aorta.



**Attenuated wall**


Simplified transverse schematic

# Aneurysm, pseudoaneurysm and dissection

| Feature            | True aneurysm                     | Pseudoaneurysm                                   | Dissection                              |
|--------------------|-----------------------------------|--------------------------------------------------|-----------------------------------------|
| Fundamental defect | Dilation of intact, weakened wall | Wall defect with extravascular hematoma          | Blood separates planes of arterial wall |
| Blood location     | Within dilated lumen              | Contained outside wall, communicating with lumen | Within a wall channel                   |
| Common danger      | Rupture; thromboembolism          | Rupture                                          | Rupture; branch compromise              |


# Why the abdominal aorta?

- Atherosclerosis-associated aneurysms develop most often in the abdominal aorta and common iliac arteries.
- The typical AAA lies distal to the renal arteries and proximal to the aortic bifurcation.
- Distribution is a key gross anatomic clue.



# Saccular versus fusiform geometry

- Saccular: Focal, asymmetric, roughly spherical outpouching.
- Fusiform: Circumferential dilation of an elongated arterial segment.
- Either shape may occur in the abdominal aorta; Shape alone does not establish etiology.




**Saccular**

**Fusiform**

# AAA distribution and neighboring arteries

- The renal arteries mark the usual upper boundary.
- Inferior mesenteric and iliac arterial origins may become involved.
- Smaller accompanying iliac aneurysms are not uncommon.



## Atherosclerosis is central—but not sufficient

### **Intimal disease**

### **Medial vulnerability**


- Atherosclerosis is a major cause of AAA.
- Extensive abdominal aortic atherosclerosis can exist without aneurysmal dilation.
- Aneurysm formation requires additional loss of structural wall integrity.

## Age, sex and smoking associations

- AAAs are reported more frequently in males and smokers.
- They rarely develop before age 50 years account.
- The risk profile contextualizes an incidental aortic dilation but does not replace morphologic assessment.


# Aortic wall architecture

- Intima: Luminal layer susceptible to atheromatous thickening.
- Media: Elastic lamellae and smooth muscle cells provide support.
- Adventitia: Supporting outer wall with vessels supplying outer media.



Layer relationships are schematic

# Vessel-wall strength is actively maintained



Aneurysmal dilation develops when the balance fails.


## Extracellular matrix is load-bearing

### Elastic recoil

### Structural strength

- Elastin contributes recoil during pulsatile pressure cycles.
- Collagens and other matrix molecules contribute wall integrity.
- Destruction or deficient repair leaves the aortic media unable to resist dilation.

# Inflammation accelerates matrix destruction



The pathology is remodeling failure—not merely luminal obstruction.

# Macrophages and metalloproteinases

- Macrophages can generate matrix metalloproteinases (MMPs).
- MMPs degrade elastin, collagen and additional matrix constituents.
- Proteolysis weakens the load-bearing aortic wall and facilitates expansion.

# Matrix degradation is normally restrained

## **MMPs: Degradation**

## **TIMPs: Restraint**

- MMP activity favors proteolytic remodeling.
- Tissue inhibitors of metalloproteinases (TIMPs) oppose excessive matrix breakdown.
- Increased MMPs or decreased TIMP expression can tip the balance toward matrix loss.

# Why loss of elastin matters


Loss and fragmentation of  
elastic components



Less effective arterial  
recoil and support



**Progressive dilation under  
pulsatile loading**




## Local cytokines modify proteolysis

- A shift toward T-helper 2 cytokine signaling in AAA-associated inflammation.
- Interleukin-4 and interleukin-13 can promote macrophage elastolytic MMP production.
- The inflammatory environment therefore influences structural degradation.


# Plaque thickness impairs medial nutrition

- Atherosclerotic intimal thickening increases the diffusion path from the lumen.
- Reduced nutrient and oxygen delivery can produce ischemic injury in the inner media.
- Smooth muscle cell loss impairs matrix maintenance and repair.



Layer relationships are schematic

## Smooth muscle loss reinforces wall failure



Medial thinning follows from impaired repair plus matrix degradation.

## Medial degeneration: A shared endpoint

- Elastic tissue injury and smooth muscle cell loss create a structurally weakened media.
- The term medial degeneration describes a pattern, not a specific etiology.
- Interpret this change with the lesion distribution, plaque burden and inflammatory context.

# AAA pathogenesis: An integrated model

Atherosclerotic plaque and local inflammation

MMP-driven extracellular matrix injury


Linked features of one disease process

Medial ischemia and smooth muscle cell loss

**Progressive weakening → enlargement**


# Typical AAA: Infrarenal distribution

- Usually between the renal arteries and the aortic bifurcation.
- Anatomic involvement can extend toward branch-vessel origins.
- The iliac arteries may contain additional aneurysms.



## AAA size and longitudinal extent


- AAA diameters greater than 3 cm.
- Examples may exceed 5.5 cm and extend up to 25 cm in length.
- These are descriptive textbook morphology, not stand-alone treatment instructions.



**Aneurysmal diameter**

# Gross wall appearance

- Severe complicated atherosclerosis accompanies many conventional AAAs.
- The underlying aortic media is destroyed and attenuated.
- Dilation produces a wall prone to rupture.




**Attenuated wall**

Simplified transverse schematic

# Mural thrombus: A frequent feature

- AAAs frequently contain bland, poorly organized mural thrombus.
- Thrombus may be laminated and occupy much of the aneurysmal sac.
- The thrombus is not a replacement for a structurally competent arterial wall.




**Mural thrombus**

Simplified transverse schematic

# How a mural thrombus is positioned

- Residual flowing lumen lies centrally or eccentrically.
- Layered thrombotic material abuts the internal wall.
- Atherosclerotic and thinned medial tissue remains external to the thrombus.




**Mural thrombus**

Simplified transverse schematic


# Branch-vessel ostia can be compromised

- Aneurysmal extension may directly involve nearby branch origins.
- Mural thrombus can obstruct a vessel ostium.
- Downstream ischemic injury then reflects the supplied vascular territory.



## Common iliac aneurysms may coexist

- Atherosclerosis-related aneurysmal disease can extend beyond one short aortic segment.
- Smaller iliac aneurysms may accompany the main AAA.
- The complete vascular distribution matters when interpreting gross findings.



# Reading the gross specimen systematically

- 1 Localize dilation relative to renal arteries and bifurcation.
- 2 Identify saccular or fusiform configuration.
- 3 Inspect plaque burden, media attenuation, mural thrombus and rupture site.
- 4 Assess branch ostia and iliac extension.

# Microscopic correlation: Common AAA

- Atheromatous intimal lesions overlie the affected region.
- The media shows depletion of smooth muscle and elastic support.
- Inflammatory and proteolytic injury can contribute to structural failure.

**Plaque thickens intima**

**Media: Elastic lamellae + SMC**

**Adventitia**

Layer relationships are schematic

## Inflammatory AAA is a distinct variant

**Inflamed aortic wall**

**Periaortic scarring**

- Inflammatory AAAs account for approximately 5%–10% of all AAAs.
- They exhibit conspicuous inflammation and periaortic fibrosis.
- Their recognition changes the expected presentation and differential.

## Clinical pattern of inflammatory AAA

- Patients tend to be younger than those with conventional AAA description.
- Back pain is a frequent presentation.
- Inflammatory markers such as erythrocyte sedimentation rate may be elevated.

## Inflammatory infiltrate in the wall


### **Lymphoplasmacytic cells**

### **Macrophages / giant cells**

- Numerous lymphocytes and plasma cells are present.
- Macrophages and sometimes giant cells accompany the inflammation.
- This infiltrate is linked to surrounding fibrosing tissue reaction.

# Periaortic fibrosis and regional extension

- Dense periaortic scarring is characteristic.
- Fibrosis can extend into the anterior retroperitoneum.
- Anatomic extension helps explain compressive consequences.




**Periaortic fibrosis**

## Inflammatory AAA may remain localized

- The inflammatory response is presumed to be directed against the abdominal aortic wall.
- Most such cases are not accompanied by inflammation of other arteries.
- Localized periaortitis should not automatically be labeled systemic vasculitis.

## A subset is associated with IgG4-related disease



Association is not synonymous with every inflammatory AAA being IgG4-related.

## IgG4-related disease: Tissue clues

- Affected tissue may show storiform fibrosis.
- IgG4-positive plasma cells can be present within the inflammatory infiltrate.
- Circulating IgG4 is elevated in many, but not all, affected patients.

# IgG4-related disease may involve other organs

Aorta and adjacent retroperitoneum

Pancreas


Linked features of one disease process

Thyroid

**Other vascular sites, including ascending  
aorta**

# Retroperitoneal fibrosis can affect ureters

- Dense fibrosis may involve tissues around both ureters.
- Obstructed urinary drainage can lead to bilateral hydronephrosis.
- This link makes an inflammatory aortic lesion clinically more than a focal aneurysm.



# Why identifying IgG4-related disease matters

## **Treatable inflammation**


## **Mechanical aneurysm risk**

- Response of IgG4-related disease to glucocorticoid and anti-B-cell therapies.
- Recognition may affect treatment of the systemic inflammatory/fibrosing component.
- Aneurysm integrity and rupture risk remain separate clinical concerns.

## “Mycotic” aneurysm means infection

- A mycotic AAA is infected; The term does not imply a fungal cause.
- Circulating microorganisms can seed the aortic wall.
- The infected lesion is pathobiologically different from uncomplicated atherosclerotic dilation.

## Infection further destroys the media



Destructive suppuration is the mechanistic warning sign.

# Conventional, inflammatory and mycotic AAA

| Feature                | Conventional AAA                   | Inflammatory AAA                            | Mycotic AAA                        |
|------------------------|------------------------------------|---------------------------------------------|------------------------------------|
| Dominant context       | Atherosclerosis with medial injury | Inflammatory infiltrates and fibrosis       | Infection of aortic wall           |
| Histologic clue        | Medial loss; atheroma; thrombus    | Lymphoplasmacytic cells and periaortic scar | Suppuration and medial destruction |
| Characteristic concern | Rupture, thrombus, branch effects  | Pain, inflammation, fibrosis/ureters        | Rapid dilation and rupture         |

## Most AAAs cause no symptoms

- Many are discovered incidentally rather than because of a specific symptom.
- A pulsating abdominal mass may be detected on physical examination.
- A mass-like presentation can simulate an abdominal tumor.


## An aneurysm can simulate a mass lesion

**Mass effect**

**Arterial pulsation**

- Palpable pulsation suggests a vascular origin of an abdominal mass.
- The lesion is a dilated arterial lumen and wall rather than a solid neoplasm.
- Gross location and thrombus help account for the mass-like enlargement.

# Rupture is the catastrophic complication



Wall integrity—not mere luminal diameter—is the immediate pathologic determinant.

## Rupture can enter two compartments


### **Intraperitoneal**

### **Retroperitoneal**


- Hemorrhage may extend into the peritoneal cavity.
- Alternatively, bleeding may enter retroperitoneal tissues.
- Both routes can produce massive blood loss.

# Branch obstruction causes territorial ischemia

- Direct extension or occlusion of branch ostia compromises arterial flow.
- The end-organ effects depend on which vessel is affected.
- Mural thrombus can be part of the obstructing material.




## Iliac involvement threatens leg perfusion




Aortic aneurysm can produce symptoms remote from the abdomen.

# Renal involvement threatens kidney perfusion



Differentiate arterial ischemia from ureteric obstruction due to fibrosis.

# Mesenteric compromise threatens bowel



The precise territory depends on the branch affected.

## Mural thrombus can embolize



Aneurysm-associated embolism is distinct from fixed ostial obstruction.

## Adjacent structures can be damaged

### **Ureteral compression**

### **Vertebral erosion**


- An enlarged AAA can compress a ureter.
- Long-standing impingement may erode vertebral structures.
- These manifestations arise from mass effect rather than embolic disease.

## Mechanism determines the complication

| Mechanism                | Pathologic event                          | Potential consequence   |
|--------------------------|-------------------------------------------|-------------------------|
| Rupture                  | Wall defect and major bleeding            | Fatal hemorrhage        |
| Fixed branch obstruction | Ostial coverage or aneurysmal involvement | Organ ischemia          |
| Embolism                 | Thrombus or atheroma migrates             | Distal ischemia         |
| Compression              | Expansion impinges on adjacent structures | Ureter/vertebral injury |

# Rupture risk increases with diameter

## Estimated annual rupture risk



Risk estimates are historical textbook descriptors, not modern individual-patient risk predictions.

## AAA enlargement is heterogeneous

- Most AAAs are described as expanding about 0.2–0.3 cm per year.
- Approximately one-fifth expand more rapidly account.
- Average enlargement therefore does not imply uniform individual behavior.

### Common expansion estimate




**0.2 cm/year**



**0.3 cm/year**

# Why increasing size changes management



Management requires specialist assessment; Textbook thresholds are not presented as current guidelines.

## Aggressive management at $\geq 5$ cm

- More active management for aneurysms at or above approximately 5 cm.
- Choice includes open graft reconstruction or an endovascular stent-graft approach.
- This is a source-specific teaching statement, not an updated eligibility algorithm.

## Open repair replaces a weakened segment


Aneurysmal segment is approached surgically




A prosthetic graft reconstructs the arterial channel



**Blood is carried through the graft instead of the untreated sac**



# Endovascular repair excludes the sac



Schematic principle of endovascular aneurysm repair (EVAR).

# The timing of repair matters

## Unruptured intervention

## After rupture

- Approximately 5% operative mortality for unruptured aneurysms.
- After rupture, fewer than half survive even with emergency surgery account.
- These historic textbook figures motivate prevention; They are not individualized outcome estimates.

# AAA is a marker of systemic atherosclerosis

Abdominal aorta: Aneurysmal disease

Coronary arteries: Ischemic heart disease  
risk

Linked features of one disease process

Cerebral circulation: Stroke risk

**Other vascular beds: Coexistent plaque  
disease**


## Clinical case: Incidental pulsatile mass

A man older than 60 with a smoking history is found to have a pulsatile abdominal mass.

Aneurysmal dilation is seen below the renal arteries.

**What mechanism best accounts for the weakening of the aortic wall?**

## Case interpretation: Wall injury, not just plaque



The correct mechanism combines inflammatory proteolysis and medial injury.

## Clinical case: Back pain and a fibrosing aorta

A patient has an infrarenal aneurysm, back pain and elevated inflammatory markers.

The periaortic tissue is densely fibrotic and lymphoplasmacytic.

**Which AAA variant is supported, and what associated disease warrants consideration?**

## Case interpretation: Inflammatory AAA

### Inflammatory phenotype

### Consider IgG4-RD subset

- The pattern supports inflammatory AAA.
- An IgG4-related component is possible in a subset, especially with characteristic fibrosis and other organ involvement.
- Bilateral hydronephrosis can reflect retroperitoneal fibrosing disease.

## Clinical case: Suppurative aortic lesion

An aortic aneurysm contains an infected wall with suppurative medial destruction.

The lesion is enlarging rapidly.

**What form of AAA is indicated and why is the lesion dangerous?**

## Case interpretation: Mycotic AAA



The defining feature is infection—not fungal etiology.

## Distinguishing related aortic disorders

| Entity                      | Typical textbook association               | Structural pattern                           |
|-----------------------------|--------------------------------------------|----------------------------------------------|
| Abdominal aortic aneurysm   | Atherosclerosis and smoking association    | Infrarenal dilation with medial loss         |
| Thoracic ascending aneurysm | Hypertension; selected inherited disorders | Aortic wall degeneration in thoracic segment |
| Aortic dissection           | Blood enters and separates wall planes     | Intramural blood-filled channel              |

# AAA pathology: Integrated synthesis

Risk and atherosclerotic context

Matrix degradation + smooth muscle cell  
loss

Linked features of one disease process

Medial weakening and aneurysmal  
dilation

**Rupture, embolism, obstruction or  
compression**

## Core diagnostic and clinical distinctions

- The infrarenal location, attenuated media, complicated plaque and mural thrombus typify common AAA.
- Inflammatory AAA adds lymphoplasmacytic inflammation and dense periaortic fibrosis.
- Mycotic AAA adds microbial infection and destructive suppuration.
- Diameter relates to rupture risk; Associated atherosclerosis signals broader vascular risk.

## Primary source and scope

- Robbins, Cotran & Kumar: Pathologic Basis of Disease, 11th edition.
- Chapter 11, “Blood Vessels”: “Aneurysms and Dissection” and “Abdominal Aortic Aneurysm,” printed pp. 458–462.
- All morphology, mechanisms and quantitative estimates in this lecture are from the supplied textbook extraction.
- Original schematic diagrams are educational models—not clinical photographs or histologic specimens.