

Acute Pancreatitis

Etiology · Pathogenesis · Morphology · Clinical consequences

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning objectives

- Explain how normal pancreatic defenses prevent enzyme-mediated injury.
- Classify obstructive, toxic, metabolic, vascular, infectious and hereditary triggers.
- Connect aberrant enzyme activation to edema, fat necrosis and hemorrhage.
- Interpret the clinical, laboratory and systemic consequences of acute pancreatitis.

Acute pancreatitis is a potentially reversible injury


- Acute pancreatic parenchymal injury is accompanied by an inflammatory response.
- Limited attacks may resolve without lasting structural or functional damage.
- Necrosis and systemic mediator release distinguish life-threatening disease.
- Recurrent injury can eventually produce chronic damage.

Exocrine anatomy explains the site of injury




Premature intrapancreatic activation converts a normal digestive system into a tissue-injury mechanism.

Zymogen storage separates synthesis from activity



Most pancreatic proteases should not be active within the gland.


Trypsin activation normally occurs outside the pancreas



The anatomical separation of zymogen synthesis and protease activation protects pancreatic tissue.

Trypsin inhibitors provide a second defense

- Acinar and ductal cells produce intrapancreatic trypsin inhibitors.
- SPINK1 limits inappropriate trypsin activity before it propagates.
- Failure of inhibitory control amplifies susceptibility to injury.




A small amount of ectopic activation need not cause pancreatitis when control mechanisms remain intact.

Three safeguards oppose pancreatic autodigestion

Safeguard	Protective role
Inactive zymogens	Limit active protease inside acinar cells
Duodenal activation	Confines major proteolytic activation to intestinal lumen
SPINK1 inhibition	Restrains any prematurely activated trypsin

Breakdown of safeguards initiates pancreatitis




Several initiating pathways may coexist in a single patient.

Acute and chronic pancreatitis differ in outcome

Feature	Acute pancreatitis	Chronic pancreatitis
Dominant process	Acute injury and inflammation	Repeated/persistent injury with irreversible damage
Potential reversibility	Often complete recovery	Persistent functional and structural loss
Typical teaching emphasis	Enzyme activation, edema, necrosis	Fibrosis and permanent dysfunction

Acute pancreatitis spans mild to destructive disease



This is a spectrum of injury, not a mandatory stepwise progression for every patient.


Etiology is mechanistically heterogeneous

Etiologic class	Examples in Robbins
Mechanical	Gallstones, biliary sludge, perampullary obstruction, procedures
Toxic/metabolic	Alcohol, hypertriglyceridemia, hypercalcemia, implicated drugs
Vascular/infectious	Shock, embolism, vasculitis; mumps, coxsackievirus
Inherited predisposition	PRSS1, SPINK1, CFTR, CASR, CTRC, CPA1

Biliary disease and alcohol dominate common causes

- Biliary tract disease and alcohol as the principal contributors to cases in many settings.
- Etiologic proportions vary substantially by population and alcohol exposure.
- A biliary etiology is clinically important because obstruction can be transient or ongoing.
- Population percentages should not be extrapolated to an individual patient.

A migrating gallstone can obstruct pancreatic outflow



The ductal obstruction pathway connects an anatomical blockage to enzymatic tissue damage.

Duct obstruction creates an interstitial insult

- Increased intraductal pressure promotes leakage of enzyme-rich secretions.
- Lipase is secreted in an active form and can injure adjacent adipose tissue.
- Edema can further impair regional microvascular perfusion.

Outflow blocked



Enzymes enter interstitium

Mechanical obstruction and microcirculatory injury can reinforce each other.

Biliary sludge can have the same obstructive effect

- Pancreatic duct obstruction is not confined to a large radiologically obvious gallstone.
- Biliary sludge is included among common obstructive triggers.
- Physiological consequences depend on impaired outflow and pressure, not stone size alone.

Nonbiliary mechanical causes are clinically diverse

- Periapillary neoplasms may obstruct pancreatic drainage.
- Choledochoceles and congenital duct configurations may contribute.
- Parasites such as *Ascaris lumbricoides* or *Clonorchis sinensis* may block the outflow pathway.
- Ductal anatomy should be considered when common etiologies are absent.

Pancreas divisum is a predisposing ductal context


- The congenital duct arrangement alters the drainage route of pancreatic secretions.
- The source lists pancreas divisum among potential obstructive contributors.
- Its presence does not establish causality in every pancreatitis episode.

Duct configuration



Potential impaired drainage

Alcohol-related pancreatitis is multifactorial




That important aspects remain incompletely understood.

Alcohol alters pancreatic ductal secretions

- Chronic alcohol exposure can promote protein-rich pancreatic secretion.
- Inspissated protein plugs may obstruct small ducts.
- Transient sphincter of Oddi contraction is another proposed outflow influence.
- Together these effects can increase ductal pressure and acinar stress.

Alcohol also targets acinar-cell homeostasis



These mechanisms are proposed contributors, not a complete causal account.

Alcohol exposure alone is usually insufficient

- Most people with prolonged alcohol exposure do not develop pancreatitis.
- Clinical attacks commonly occur after years of heavy exposure.
- Susceptibility likely reflects multiple interacting cellular and host factors.
- The specific determinants remain incompletely resolved in the cited source.

Metabolic disturbances can initiate injury

Trigger	Textbook mechanistic connection
Hypertriglyceridemia	Included among metabolic causes of pancreatic acinar injury
Hypercalcemia	Intracellular calcium elevation favors inappropriate trypsin activation
Alcohol-associated metabolism	Can intensify oxidative and calcium stress


Drug associations require careful interpretation

- Drugs among suspected and established precipitants.
- Its examples include azathioprine, statins and selected antidiabetic agents.
- The strength of causal evidence varies by medication.
- Drug attribution requires a compatible clinical context and exclusion of competing causes.

Procedures and trauma can damage acinar cells

- Blunt abdominal trauma may directly injure pancreatic tissue.
- Operative injury and endoscopic retrograde cholangiopancreatography are iatrogenic settings.
- Direct cellular damage releases digestive enzymes and promotes inflammation.
- The trigger may be mechanical even without sustained duct obstruction.

Vascular insufficiency is another initiating event



Ischemia may be an initiating cause or a consequence of severe edema.


Infections may directly injure the pancreas

- Mumps is a named viral example in the Robbins etiologic table.
- Coxsackievirus is another infectious association.
- Direct injury of acinar cells can initiate the enzyme-mediated injury cycle.
- Infections are part of the differential when common causes are absent.

Inherited susceptibility often presents early or repeatedly


- Hereditary forms may cause recurrent severe attacks beginning in childhood.
- Repeated attacks can progress to chronic pancreatitis.
- Many predisposition genes ultimately alter trypsin activation, inhibition or ductal fluid composition.
- A family history supports—but is not required for—genetic evaluation.

PRSS1 gain of function undermines trypsin control




Autosomal-dominant inheritance for PRSS1-associated hereditary pancreatitis.

SPINK1 loss of function weakens trypsin inhibition



Recessive inheritance for the hereditary form associated with inadequate SPINK1 activity.

CFTR loss of function impairs ductal fluid chemistry



Heterozygous or homozygous variants can contribute, particularly with other susceptibility alleles.

CASR connects extracellular calcium sensing to injury

- The calcium-sensing receptor regulates calcium handling relevant to the pancreatic lumen.
- Loss-of-function variants may alter calcium concentrations.
- Enhanced calcium-associated trypsin activation increases pancreatitis susceptibility.

CASR altered



Trypsin activation favored

CTRC and CPA1 broaden the genetic mechanisms

Gene	Pathogenic effect in Robbins	Consequence
CTRC	Loss of function	Reduced protective degradation of trypsin
CPA1	Toxic gain of function	Abnormal carboxypeptidase-related zymogen regulation


Genetic risks converge on protease homeostasis

Gene	Dominant biological defect
PRSS1	Excessive / sustained trypsin activity
SPINK1	Loss of trypsin inhibition
CFTR	Reduced bicarbonate-rich ductal fluid; plugs
CASR	Disturbed calcium-mediated regulation
CTRC / CPA1	Defective trypsin degradation / altered zymogen regulation

Recurrent attacks create a cancer-risk context


- PRSS1-associated hereditary pancreatitis is linked to increased lifetime pancreatic cancer risk.
- A 40% lifetime estimate in the described hereditary population.
- The biological context is recurrent inflammation and chronic tissue injury.
- The estimate should be interpreted in its specific source population, not generalized to every pancreatitis patient.

Three major initiating pathways are proposed




These are conceptual mechanisms Figure 19.2; They may interact rather than act independently.

Obstruction begins with a pressure-and-leak sequence




Interstitial edema and ischemia can create feedback injury to acinar cells.

Lipase converts escaped secretion into fat injury




Lipase-associated fat injury explains why necrosis appears in and around the pancreas.

Fat necrosis can amplify interstitial inflammation




Local damage-associated signaling as a mechanistic hypothesis.

Edema compromises microvascular oxygen delivery



Once established, edema is both a morphological finding and an injury amplifier.

Direct acinar injury releases digestive machinery




Direct acinar injury does not require a preceding biliary obstruction.

Oxidative stress damages acinar membranes


- Free radicals trigger membrane lipid oxidation.
- Cellular injury disturbs organelle integrity and secretory pathways.
- Inflammatory transcription programs can be activated in parallel.
- Enzyme release and inappropriate activation follow acinar destabilization.

AP-1 and NF- κ B link injury to inflammatory recruitment




This pathway helps explain why enzymatic damage quickly recruits inflammation.

Intracellular calcium modulates trypsin behavior



The source contrasts high-calcium activation with low-calcium trypsin autoinhibition.

Defective proenzyme trafficking is a proposed pathway




Described predominantly in experimental animal models.

Trafficking evidence has an important limitation

- The source explicitly states that the human relevance of this animal-model pathway remains unclear.
- Cathepsin-associated lysosomal activation is therefore a mechanistic proposal.
- Avoid presenting the proposed trafficking route as established for all human cases.

Trypsin acts as a proteolytic amplifier



Trypsin activation is a central organizing concept, but not the only injurious process.


Different enzymes produce different lesions

Activated enzyme	Dominant tissue effect
Proteases / trypsin	Autodigestion and parenchymal necrosis
Lipase / phospholipase	Fat breakdown and fat necrosis
Elastase	Vessel wall injury with hemorrhage

Proteolysis destroys pancreatic parenchyma


- Activated digestive proteases injure acini and nearby structures.
- Severe injury extends to ducts and pancreatic islets.
- Proteolytic destruction distinguishes necrotizing pancreatitis from limited edema.
- The distribution of injury varies with attack severity and duration.

Lipase-mediated fat necrosis produces calcium soaps




This saponification process accounts for chalky gross foci and basophilic granular material microscopically.

Elastase-mediated vascular injury causes hemorrhage




Hemorrhage implies vascular destruction, not merely edema.

Inflammation recruits systemic plasma cascades



The interaction of enzyme activation and circulation increases severity.

Positive feedback sustains the acute attack



Bidirectionality: Injury both triggers and results from enzyme activation.


Five tissue lesions define the pathological spectrum

- Microvascular leakage and interstitial edema.
- Enzymatic fat necrosis.
- Acute inflammatory infiltration.
- Parenchymal autodigestion with necrosis.
- Blood vessel destruction and hemorrhage.

Interstitial pancreatitis preserves much of the parenchyma

Typical finding	Mechanistic explanation
Interstitial edema	Microvascular leakage
Mild acute inflammation	Mediator-mediated leukocyte recruitment
Focal intrapancreatic fat necrosis	Local lipase activity
Peripancreatic fat necrosis	Diffusion or leakage beyond gland

Edema is an early anatomical consequence of permeability




Edema itself can worsen tissue perfusion.


Microscopic fat necrosis reflects adipocyte injury

Simplified adipocyte morphology — not a clinical specimen

Preserved adipose tissue




Enzymatic fat necrosis



Schematic representation only; This is not a histology micrograph.

Saponification links fat injury to hypocalcemia



Hypocalcemia is a potential clinical correlate, not proof of a specific microscopic lesion.

Necrotizing pancreatitis destroys glandular elements

- Necrosis may involve acini, pancreatic ducts and islets.
- Parenchymal destruction reflects severe enzyme-mediated injury.
- Peripancreatic tissues may also undergo necrosis.
- Necrosis and organ failure are unfavorable prognostic findings.

Hemorrhagic pancreatitis adds vessel-wall destruction

- Extensive parenchymal necrosis accompanies vascular injury.
- Grossly the tissue may appear red-black with yellow-white chalky fat-necrosis foci.
- Blood extravasation reflects destruction of vessel-wall integrity.

Elastase injury



Hemorrhage

Fat necrosis can extend beyond the pancreatic gland

- Adjacent omentum and mesentery may be involved.
- Systemically released lipase can injure distant adipose deposits.
- Subcutaneous fat may occasionally be involved.
- Peritoneal fluid may contain visible fat globules.

Gross severity parallels the dominant tissue lesion

Pattern	Gross / microscopic emphasis
Limited interstitial	Swelling, edema, modest focal inflammation
Necrotizing	Destruction of gland and peripancreatic fat
Hemorrhagic	Necrosis plus vascular destruction and hemorrhage

Severe pancreatitis may produce abnormal peritoneal fluid

- Pancreatic inflammatory exudation may enter the peritoneal cavity.
- Serous, slightly turbid, brown-tinged fluid.
- Fat globules indicate local digestion of adipose tissue.
- Fluid appearance is a clinicopathologic clue, not a stand-alone diagnosis.


Abdominal pain is the cardinal clinical manifestation

- Pain is classically constant and intense.
- The upper abdomen is predominantly involved.
- Referral to the upper or middle back is characteristic.
- Occasional radiation toward the left shoulder may occur.

Gastrointestinal symptoms reflect the acute attack

- Anorexia, nausea and vomiting are common.
- The clinical spectrum ranges from discomfort to incapacitating pain.
- The pattern should be interpreted with the abdominal examination and laboratory evidence.

Acute-phase laboratory evidence reflects acinar leakage



A raised enzyme concentration is evidence of acinar injury, not a direct grade of tissue necrosis.

Lipase and amylase differ in diagnostic persistence

Feature	Serum amylase	Serum lipase
Early elevation	First 4–12 hours in source	First 4–12 hours in source
Persistence	May normalize in 3–5 days	May remain high for 8–14 days
Textbook interpretation	Less specific than lipase	Preferred pancreatic enzyme marker


CT imaging demonstrates structural consequences

- Computed tomography may visualize pancreatic enlargement and inflammation.
- Anatomical imaging complements clinical findings and serum enzymes.
- Imaging can help assess the extent and distribution of destructive change.
- The supplied excerpt does not specify contemporary imaging-timing criteria.

Metabolic laboratory changes reflect tissue injury

Finding	Clinicopathologic explanation
Hypocalcemia	Calcium can be consumed in fat saponification
Glycosuria	Can accompany acute pancreatitis and acinar/islet dysfunction
Leukocytosis	Systemic inflammatory response to acute injury

Enzyme and cytokine release can become systemic




Severe pancreatitis can be a systemic inflammatory illness, not only abdominal pain.

Acute respiratory distress is a remote complication


- Systemic mediator release can damage pulmonary microvasculature.
- Increased alveolar-capillary permeability promotes respiratory failure.
- Acute respiratory distress syndrome as an ominous complication.
- Respiratory compromise signals severe systemic involvement.

Renal injury may follow shock and inflammation



Renal dysfunction in this setting is a marker of severe systemic disease.

Coagulation abnormalities add to multiorgan risk



The process can evolve into disseminated intravascular coagulation.

Necrotic tissue can become secondarily infected

- Necrotizing pancreatitis can leave extensive acellular tissue debris.
- Gut-derived gram-negative organisms may infect necrotic material.
- Infection further complicates an already destructive attack.
- Anatomical necrosis and infected necrosis are related but distinct observations.


Local sequelae may persist after an acute episode

- Pancreatic pseudocysts and abscess-like sequelae.
- Collections reflect residual damaged tissue and inflammatory fluid.
- The cited text does not supply current standardized collection definitions.
- Precise classification of collections should not be inferred from the excerpt alone.

Organ failure and necrosis carry prognostic weight

Unfavorable indicator	Why it matters
Pancreatic necrosis	Evidence of extensive enzyme-mediated destruction
Systemic organ failure	Circulatory and inflammatory injury beyond the gland
ARDS / renal failure	Potentially life-threatening remote-organ dysfunction
Infected necrotic debris	Complicates local injury and systemic inflammation

A focused diagnostic synthesis integrates three domains




These are complementary clinical domains; Formal criteria are not specified in the provided section.

Clinical vignette: Biliary mechanism

- A patient develops sudden constant epigastric pain radiating to the back, with vomiting.
- Serum lipase is elevated, and gallstones are identified.
- Question: Explain the initiating mechanical event and subsequent tissue injury.

Discuss the mechanism before advancing.

Clinical reasoning: Obstruction progresses to edema




The clinical vignette illustrates mechanism; It does not establish the exact stone position.

Clinical vignette: Severe systemic injury

- A patient with acute pancreatitis develops hypotension, respiratory distress and renal dysfunction.
- Structural evaluation shows extensive pancreatic necrosis.
- Question: Identify the local and systemic processes explaining deterioration.

Discuss the mechanism before advancing.

Clinical reasoning: Local injury becomes multisystem disease




Remote-organ failure can drive morbidity even when the initiating injury is intra-abdominal.

Key distinction: Reversible attack versus permanent injury

Acute episode	Persistent/recurrent injury
Potentially reversible edema and acute inflammation	Repeated attacks can damage parenchyma permanently
Some attacks include necrosis	Progression may culminate in chronic pancreatitis
Immediate issue is acute injury and systemic severity	Long-term issue includes functional loss

Integrated mechanism of acute pancreatitis



The central bridge from etiology to morphology is aberrant enzymatic activity.

Five concepts to retain

- Pancreatic defenses separate zymogen synthesis, activation and inhibition.
- Duct obstruction, acinar injury and abnormal trafficking are proposed initiating pathways.
- Trypsin amplifies proteases; Lipase injures fat; Elastase injures vessels.
- Interstitial edema, fat necrosis, necrosis and hemorrhage form a severity spectrum.
- Severity is driven by pancreatic destruction and systemic organ dysfunction.

References

Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th edition (2026). Chapter 19, The Pancreas: Pancreatitis and Acute Pancreatitis, pp. 801–804.

Primary topic: pancreatic protection, etiologies, pathogenesis, morphology and clinical features.

All diagrams are original editable educational schematics; no source textbook images were reproduced.