

Alcohol-Associated Liver Disease

Pathogenesis · morphology · clinicopathologic correlation

Dr Abu Suraih Sakhri


MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Connect alcohol metabolism to lipid accumulation and hepatocyte injury.
- Recognize the morphologic spectrum from steatosis to cirrhosis.
- Explain clinical presentation, laboratory patterns, and important diagnostic pitfalls.

Three overlapping patterns of liver injury



These patterns may coexist; Progression is not obligatory or uniform.

Spectrum is not an inevitable linear pathway

- Hepatic steatosis is an early, often reversible response.
- Only a subset develops inflammatory injury and progressive fibrosis.
- Some patients first present with cirrhosis without a recognized preceding syndrome.

Alcohol exposure: Dose is important, not sufficient

Short-term exposure

Even 80 g of ethanol over 1–several days may induce reversible steatosis.

Sustained high intake

Daily consumption around ≥ 80 g increases the risk of severe hepatic injury.


Long-duration exposure

About ≥ 160 g/day over 10–20 years is frequently associated with severe disease.

Variable outcome

The source reports cirrhosis in only ~10–15% of patients with alcohol use disorder.

Susceptibility varies among drinkers




Consumption alone does not determine which individuals develop advanced scarring.

Sex modifies susceptibility

- The source describes higher dose-for-dose susceptibility in females.
- Variation in ethanol pharmacokinetics and metabolism may contribute.
- Estrogen-related responses to intestinal endotoxin are a proposed additional mechanism.

The gut–liver axis provides a second insult



An inflammatory amplifier may operate alongside direct ethanol toxicity.

Genetic susceptibility is multifactorial

- Twin studies support a genetic component, confounded by environment.
- Polymorphisms in alcohol-processing enzymes may influence exposure to metabolites.
- Variation in inflammatory mediators may modify the response to injury.

Low-activity ALDH variants alter ethanol tolerance

Ethanol oxidation



Alcohol dehydrogenase forms acetaldehyde.

ALDH activity



Low activity limits conversion of acetaldehyde to acetate.

Clinical response



Acetaldehyde accumulation causes flushing, nausea, and intolerance.

Comorbid injury amplifies alcohol-related damage

Iron overload

A second injury source that may worsen the hepatic response.

Steatotic metabolic disease

Steatosis and steatohepatitis may reflect mixed mechanisms.

Hepatitis B and C

Concurrent viral injury can aggravate severity and scarring.


Clinical implication

Do not attribute every abnormal liver test to alcohol.

MetALD acknowledges overlapping etiologies

- A category of MASLD with increased alcohol intake.
- The source gives alcohol thresholds of >140 g/week in females or >210 g/week in males.
- Overlapping metabolic and alcohol-related injury complicates etiologic attribution.

Hepatocyte ethanol metabolism generates reducing equivalents



Alcohol dehydrogenase and aldehyde dehydrogenase increase the hepatic NADH burden.

An altered NADH/NAD⁺ balance favors fat storage

High NADH



Alcohol oxidation shifts the hepatic redox state.

Fatty-acid oxidation



β -oxidation is suppressed, reducing lipid disposal.

Lipid accumulation




Triglyceride stores expand in hepatocytes.

Suppression of fatty-acid oxidation promotes steatosis

- Alcohol-induced redox changes reduce fatty-acid oxidation.
- Less oxidative disposal leaves more fatty-acid substrate available.
- Excess substrate contributes to intracellular triglyceride droplets.

De novo lipid synthesis is also increased



The source identifies increased expression of lipogenic enzymes as another route to steatosis.

Impaired lipid export adds to steatosis

Hepatocellular lipid



Triglycerides require lipoprotein packaging for export.

**Alcohol-associated
dysfunction**




Lipoprotein assembly and secretion may be impaired.

Net effect



Retention of lipids exacerbates hepatic fatty change.

Peripheral lipid flux can augment hepatic lipid uptake



The liver receives lipids in addition to those synthesized internally.

Several metabolic effects converge on steatosis

Oxidation

Fatty-acid β -oxidation decreases.

Synthesis

Fatty-acid synthetic pathways are increased.

Export

Lipoprotein assembly/secretion may be impaired.

Delivery

Peripheral lipid mobilization increases substrates.

Acetaldehyde directly injures hepatic structures

Acetaldehyde generation



Alcohol dehydrogenase forms a reactive aldehyde.

Protein adduct formation



Binding to proteins alters their function and may create neoantigens.

Structural injury



Cytoskeletal and membrane disturbances follow.

Lipid peroxidation propagates injury

- Acetaldehyde is associated with lipid-peroxidation reactions.
- Membrane lipids undergo oxidative damage, disturbing integrity.
- Protein injury and membrane dysfunction add to hepatocellular stress.

CYP2E1 induction increases toxic oxidative products

Heavy alcohol exposure



Microsomal CYP2E1 becomes induced.

Oxidation of ethanol




The pathway generates reactive oxygen species (ROS).

Cellular damage



ROS injure proteins, membranes, and mitochondria.

Mitochondrial dysfunction adds to cell death



Mitochondria are important targets of oxidant injury in affected hepatocytes.

Glutathione depletion reduces antioxidant defenses

Alcohol exposure



Methionine metabolism becomes impaired.

Glutathione ↓




Antioxidant capacity is reduced.

**Oxidative susceptibility
↑**



ROS-mediated hepatocyte injury becomes more likely.

Homocysteine can trigger ER stress



A distinct stress mechanism complements oxidant injury.

Alcohol induction can change drug hepatotoxicity

- Alcohol-induced microsomal enzymes may accelerate formation of toxic drug metabolites.
- The textbook uses acetaminophen as an example.
- Drug exposure and alcohol history should be considered together in a liver-injury assessment.

Intestinal products amplify hepatic inflammation

Intestinal barrier

→ Alcohol-related permeability permits greater endotoxin passage.


Portal delivery

→ Bacterial components reach Kupffer cells.

Innate immune activation

→ Inflammatory cytokines and chemokines are released.

CD14 and TLR4 link endotoxin to Kupffer cells



This pathway is proposed in the source as one modifier of inflammatory disease.

PAMPs and DAMPs arise from different sources


Pathogen-associated molecules

- Microbial endotoxin arrives from the gut.
- These signals reflect intestinal-derived products.

Damage-associated molecules

- Dying hepatocytes release intracellular signals.
- These signals reflect local tissue damage.

Inflammasome activation amplifies liver inflammation



Increased hepatic IL-1 β and caspase-1 in alcohol-associated disease.

A multifactorial injury model

Metabolic perturbation

Steatosis results from altered lipid synthesis, oxidation, export, and supply.

Chemical toxicity

Acetaldehyde and reactive oxygen species injure cellular constituents.


Innate immunity

Gut-derived and hepatocyte-derived signals sustain inflammation.

Tissue repair

Repeated injury elicits fibrosis and architectural distortion.

Why zone 3 is a typical starting point




Sinusoidal blood flow: Portal/periportal → centrilobular

Alcohol-associated lesions characteristically begin in zone 3.

Morphologic sequence: Injury begins centrally

- Typical changes begin in the centrilobular region (acinar zone 3).
- With greater injury, abnormality may extend toward portal tracts.
- This zonal pattern helps organize interpretation of steatosis and fibrosis.

Steatotic hepatocytes accumulate lipid vacuoles



Droplets may enlarge and coalesce, displacing nuclei to the cell margin.

Schematic: Simplified hepatocytes, not a diagnostic histology image

Macrovesicular steatosis is predominant

Macrovesicular pattern

- Large fat vacuoles distend hepatocytes.
- The nucleus is often displaced to the side.

Microvesicular pattern

- Smaller droplets occupy cytoplasm.
- This is not the predominant pattern in typical disease.

Gross fatty liver: Enlarged, yellow, greasy

- The source describes a soft, yellow, greasy organ with marked enlargement.
- Chronic severe steatosis may produce a very heavy liver.
- Gross appearance expresses widespread hepatocellular triglyceride accumulation.

Alcoholic foamy degeneration is an atypical pattern

- A relatively uncommon steatotic pattern can occur with prolonged heavy exposure.
- Diffuse small-droplet macrovesicular to microvesicular change may be seen.
- Endoplasmic reticulum and mitochondrial injury are implicated in the source.

Steatosis can reverse with abstinence


Ethanol exposure withdrawn



Hepatic fat accumulation may resolve

Simple fatty change generally permits substantial reversal when the injurious exposure stops.

Ballooning signals significant hepatocyte injury



Swollen hepatocytes have pale cytoplasm and disrupted cytoskeletal organization.

Schematic: Simplified hepatocytes, not a diagnostic histology image

Mallory hyaline is an abnormal protein aggregate

Cytoskeletal injury



Keratin intermediate filaments are damaged.

Incomplete clearance



Altered keratins accumulate with ubiquitin.

Mallory hyaline



Eosinophilic cytoplasmic aggregates may become conspicuous.

Keratin 8/18 highlight cytoskeletal alterations

- Mallory hyaline contains partially degraded, ubiquitinated keratin filaments.
- Keratins 8 and 18 may be demonstrated immunohistochemically.
- The staining pattern illustrates cytoskeletal collapse in injured ballooned cells.

Neutrophils may cluster around injured hepatocytes

- Neutrophils are commonly prominent in alcohol-associated hepatitis.
- They may surround ballooned cells, especially with Mallory hyaline.
- This distribution provides a useful morphologic clue, not a standalone diagnosis.

Inflammation involves several immune populations

Neutrophils

Often conspicuous around affected hepatocytes.

Lymphocytes

Lobular infiltrates are common; Portal lymphocytes may appear.

Macrophages

Can increase in both lobules and portal tracts.

Hepatocyte death

Spotty necrosis/apoptosis is usual; Confluent injury can occur.

Ballooning and Mallory hyaline are not equivalent


Ballooning

- An essential morphologic feature of steatohepatitis.
- May be present without Mallory hyaline.

Mallory hyaline

- A cytoplasmic inclusion in some injured cells.
- Not specific to alcohol-associated disease.

Pericellular fibrosis begins in zone 3




Perisinusoidal collagen forms a network around hepatocytes

Schematic — not a diagnostic tissue image

“Chicken-wire” fibrosis describes a network of scar


- Early perisinusoidal/pericellular fibrosis surrounds groups of hepatocytes.
- The resulting fine lattice produces a “chicken-wire” appearance.
- Collagen stains help demonstrate this matrix pattern.

Progressive scarring extends beyond zone 3



The original distribution expands with repeated, persistent injury.


Bridging fibrosis interrupts the lobular framework



Parenchymal groups are separated by broad scar

Schematic — not a diagnostic tissue image

Micronodular cirrhosis is a classic late pattern



Fibrous septa enclose regenerative parenchymal nodules

Cirrhosis also changes hepatic vascular structure

Perivenular injury



Fibrosis develops around terminal venules.

Vascular remodeling



Phlebosclerosis and veno-occlusive lesions may occur.

Functional effect



Distorted perfusion compounds parenchymal dysfunction.

“Burned-out” cirrhosis can lose obvious steatosis

- A cirrhotic liver need not retain fatty droplets or active steatohepatitis.
- Advanced scarring may dominate the biopsy or gross appearance.
- A negative steatosis finding does not exclude alcohol as an earlier cause.

Reversibility depends on the stage of disease

Steatosis

Typically reversible with cessation of alcohol exposure.

Early fibrosis

Some scar regression can occur after abstinence.

Established cirrhosis

Full restitution is less likely owing to structural and vascular alterations.

Rare exception

Complete regression of cirrhosis has been reported but is unusual.

Clinical features of simple steatosis

- Hepatomegaly may be the principal manifestation.
- Serum bilirubin and alkaline phosphatase can be mildly increased.
- Severe hepatocellular functional impairment is uncommon in isolated steatosis.

Alcohol-associated hepatitis often appears acutely

- An episode frequently follows a period of heavy drinking.
- Patients may have malaise, anorexia, weight loss, and upper abdominal discomfort.
- Tender hepatomegaly is a common examination finding.
- The syndrome may coexist with previously established cirrhosis.

Characteristic biochemical and hematologic findings

Bilirubin

Hyperbilirubinemia is common in clinically apparent hepatitis.

Transaminases

Serum AST and ALT increase.

Cholestasis markers

Alkaline phosphatase may be elevated.

Peripheral blood

Neutrophilic leukocytosis is often present.

AST tends to exceed ALT in alcohol-associated injury

$$\text{AST} : \text{ALT} \geq 2 : 1$$

This pattern as a useful clue, especially when alcohol exposure is not disclosed; It is not by itself diagnostic.

The clinical presentation can mimic other disorders

Cholestatic presentation

- An acute cholestatic syndrome can resemble extrahepatic obstruction.
- Differential assessment remains necessary.

Severe hepatitic presentation

- Advanced injury may resemble acute liver failure.
- Severity is not determined from the AST:ALT ratio alone.

Episodes of hepatitis have variable outcomes

- Resolution can follow total abstinence and adequate nutrition.
- Inflammation may persist even after exposure stops.
- Repeated episodes can lead to fibrosis and cirrhosis.
- The severity of any one episode is unpredictable.

Cirrhosis adds hepatic synthetic dysfunction

Albumin / globulins

Abnormalities reflect altered hepatic synthetic capacity.

Coagulation proteins

Reduced clotting-factor synthesis contributes to coagulopathy.

Bilirubin

Hyperbilirubinemia may reflect advanced dysfunction.

Blood counts

Anemia may accompany advanced liver disease.


Cirrhosis can remain clinically silent

- Advanced architectural injury may precede a dramatic clinical presentation.
- Infection, trauma, or another stressor can precipitate decompensation.
- Diagnosis may therefore be delayed until complications appear.

Alternative causes should still be investigated

- A history of heavy drinking does not eliminate other liver etiologies.
- In a subset of presumed alcohol-associated cirrhosis, another disease is identified.
- Biopsy may help resolve uncertainty in selected patients.
- Interpret histology together with clinical history and competing risk factors.

The pathologic consequences of cirrhosis are systemic



Loss of functional parenchyma accompanies circulatory remodeling.

Major complications drive late mortality

Hepatic failure

Encephalopathy or hepatic coma.

Portal hypertension

Severe gastrointestinal hemorrhage.

Systemic complications

Infection and hepatorenal syndrome.

Cancer risk

Hepatocellular carcinoma may develop in a cirrhotic liver.

Cirrhosis increases hepatocellular carcinoma risk

- Alcohol-associated cirrhosis provides a chronically injured, regenerative environment.
- The source reports an annual HCC risk of 1%–6% in cirrhosis.
- This is a textbook-era estimate, not an individualized risk prediction.

Abstinence is the principal modifiable determinant

- Stopping ethanol exposure permits resolution of many early morphologic changes.
- Adequate nutrition is an important supportive component of recovery.
- Established scarring and perfusion abnormalities may limit reversibility.
- Continuing exposure increases the risk of recurrent injury.

Compare three major clinicopathologic patterns

Feature	Steatosis	Steatohepatitis
Dominant lesion	Intracellular lipid droplets	Ballooning, inflammation $\pm$ Mallory hyaline
Early anatomic focus	Often centrilobular	Often zone 3 with lobular injury
Typical clinical expression	Hepatomegaly; minimal dysfunction	Acute illness, jaundice, inflammatory laboratory findings
Reversibility	Usually with abstinence	Variable; persistent disease can progress

Cirrhosis: Fibrosis and architectural distortion

Question	Pathology-based answer
Characteristic morphology	Fibrous septa surround nodules; often initially micronodular.
Residual fat required?	No: advanced “burned-out” cirrhosis may lack steatosis.
Functional problem	Loss of parenchyma plus altered vascular perfusion.
Recovery after abstinence	Improvement possible, complete regression uncommon.

Steatohepatitis has overlapping etiologies

Alcohol-associated

- More conspicuous neutrophils may favor alcohol-associated injury.
- Exposure history and distribution help interpretation.

Metabolic dysfunction–associated

- Ballooning and Mallory hyaline may also occur.
- Histologic findings overlap; No isolated lesion proves cause.

Case 1: Enlarged fatty liver

A patient with heavy ethanol exposure has an enlarged liver. Microscopy shows predominantly large intracellular lipid vacuoles, with little inflammation or fibrosis. What process predominates?

Interpretation

Alcohol-associated steatosis. In the absence of major inflammation or established scarring, cessation of exposure can permit substantial reversal.

Case 2: Ballooned cells and neutrophils

A patient develops jaundice after heavy drinking. Biopsy shows ballooned hepatocytes, Mallory hyaline, neutrophil-rich lobular inflammation, and pericellular fibrosis. How should the lesion be classified?

Interpretation

Alcohol-associated steatohepatitis is strongly supported by the combined exposure, clinical syndrome, and morphology. Individual features are not wholly specific.


Case 3: Scar without residual fatty change

A nodular cirrhotic liver shows broad fibrous septa and altered vascular architecture but no obvious hepatocyte steatosis. Does the lack of fat exclude previous alcohol-associated injury?

Interpretation

No. Advanced “burned-out” cirrhosis may lack fat or active inflammation. Etiologic assignment requires clinical and exclusionary evidence.

Synthesis: Dose, susceptibility, injury and scarring



Morphology and clinical disease arise from interacting rather than isolated mechanisms.

Textbook reference and morphologic figures

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. (2026), Chapter 18, pp. 766–769.
- Fig. 18.18: relation among steatosis, steatohepatitis, and cirrhosis.
- Figs. 18.19–18.21: steatosis and fibrosis, hepatocyte ballooning, and cirrhosis.