

Atrial Septal Defect

Development • morphology • hemodynamics • clinical significance

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)


www.abusuraihsakhri.com

What this lecture will explain

- Distinguish a true atrial septal defect (ASD) from a patent foramen ovale (PFO).
- Reconstruct atrial septation and localize three anatomical ASD types.
- Connect abnormal atrial communication to pulmonary overcirculation, clinical presentation, and closure rationale.


ASD belongs to congenital heart disease

- Congenital heart disease (CHD) comprises structural abnormalities of the heart or great vessels present at birth.
- Atrial defects belong to the septal-defect category, rather than outflow obstruction or great-vessel misrouting.
- A circumscribed lesion may be compatible with live birth but become clinically relevant later.



Timing matters: Cardiac morphogenesis


- Most CHD arises from errors in embryogenesis during gestational weeks 3–8.
- Atrial development requires coordinated cell movement, tissue reshaping, and septation.
- Disturbance of the septal program can leave an enduring interatrial opening.



Developmental or clinical stages — schematic

Building a four-chambered heart


- Cardiac progenitors converge and form a beating heart tube by about day 20.
- Chamber formation proceeds during the following days.
- By about day 50, septation of atria, ventricles, and AV valves yields a four-chambered heart.



Developmental or clinical stages — schematic

Heart-field contributions to the atria

- First-wave progenitors contribute to the developing left ventricle.
- The second heart field contributes to the outflow tract, right ventricle, and most atrial tissue.
- Errors in spatially coordinated development can produce defects of discrete cardiac structures.



Two cardiac progenitor lineages

Etiology is heterogeneous, not deterministic


- Genetic and environmental causes are established for CHD, but account for a minority of cases.
- Examples of environmental associations include congenital rubella, teratogenic exposures, and gestational diabetes.
- Most affected individuals have no identified single explanatory genetic lesion.

Contributors	Interpretation
Genetic variation	Can disrupt morphogenesis
Gestational exposures	Can alter development
Unexplained cases	Common in CHD

Native editable PowerPoint comparison table

Septal development requires transcriptional coordination

- Cardiac developmental programs use interacting transcription factors and signaling systems.
- GATA4, TBX5, and NKX2.5 as interacting factors in septal development.
- Variation in one component may disrupt their shared transcriptional targets.



Interacting transcription factors

Textbook-described interaction; Not a quantitative network

GATA4: Septal-defect association

- GATA4 among genes associated with ASD or ventricular septal defect.
- Its product is a transcription factor, not an anatomic structural component of the septum.
- A gene–phenotype association does not imply that all ASDs carry this variant.

GATA4 variation



Developmental transcription altered



Septal defect can result

Conceptual mechanism — editable schematic

NKX2.5: ASD and conduction phenotypes


- NKX2.5 is linked in the Robbins gene table to nonsyndromic ASD or conduction defects.
- The same developmental regulator may influence more than one cardiac phenotype.
- The precise relationship between a variant and its clinical manifestation can vary.

Affected phenotype	Shared regulator
Atrial septal defect	NKX2.5
Conduction defect	NKX2.5

Native editable PowerPoint comparison table


TBX5 and syndromic septal defects

- TBX5 in Holt–Oram syndrome, with ASD, VSD, or conduction defects among cardiac features.
- The gene product is a transcription factor.
- Clinical association is not equivalent to a molecular diagnosis from cardiac anatomy alone.



Complex developmental causes explain heterogeneity

- GATA4, TBX5, and NKX2.5 participate in common regulatory interactions.
- Different alterations may yield overlapping phenotypes; One alteration can yield different manifestations.
- That CHD pathogenesis remains incompletely understood.



Interacting transcription factors

Textbook-described interaction; Not a quantitative network

Three broad hemodynamic patterns in CHD


- Shunts permit flow between chambers or blood vessels.
- Left-to-right shunts usually increase pulmonary blood flow without initial cyanosis.
- Right-to-left shunts bypass the lungs; Obstructive lesions impede passage through the cardiovascular circuit.

Pattern	Principal consequence
Left → right	Pulmonary overcirculation
Right → left	Systemic desaturation
Obstruction	Restricted forward flow

Native editable PowerPoint comparison table

A true ASD is a fixed interatrial opening

- The defect is an abnormal, persistent opening within the atrial septum.
- It establishes direct communication between the right and left atria.
- Its defining anatomy is distinct from the physiologic fetal foramen ovale.



A patent foramen ovale is not a true ASD


- PFO represents postnatal failure of the normal fetal atrial channel to seal.
- The retained channel is flap-like and can open when right atrial pressure exceeds left atrial pressure.
- A true ASD instead involves a fixed abnormal opening in septal tissue.

Atrial septal defect	Patent foramen ovale
Fixed tissue opening	Potential flap-like channel
Abnormal septal development	Incomplete postnatal sealing
Usually left → right	May open transiently right → left

Native editable PowerPoint comparison table

Septation requires sequential, overlapping tissues

- The septum primum develops first and only partially divides the common atrium.
- New interatrial openings preserve fetal blood passage during septation.
- The later septum secundum overlies the channel, leaving the foramen ovale.



Developmental or clinical stages — schematic

Septum primum: Initial partial partition

- The first septal ingrowth is crescent-shaped and membranous.
- It develops posteriorly between the forming right and left atria.
- Its growth begins separation without immediately eliminating fetal atrial communication.

Developing common atrium




Septum primum begins ingrowth

Original schematic of developmental sequence; Not an anatomic section

Ostium primum: A transient opening

- The gap remaining during early septum primum growth is termed ostium primum.
- This opening allows interatrial blood movement during early development.
- It is normally obliterated as development progresses.



Ostium primum: Transient gap

Original schematic of developmental sequence; Not an anatomic section

Ostium secundum maintains interatrial transit

- Before ostium primum is completely obliterated, another opening develops in the septum primum.
- This second opening is the ostium secundum.
- Its formation prevents premature elimination of fetal interatrial communication.

Developing common atrium



Ostium secundum: New opening

Original schematic of developmental sequence; Not an anatomic section

Septum secundum creates an overlapping partition

- A second septal ingrowth develops to the right and anterior of the septum primum.
- It partially covers the ostium secundum.
- The remaining channel is the foramen ovale, connected with the ostium secundum.

Developing common atrium




Overlapping septa → fetal channel

Original schematic of developmental sequence; Not an anatomic section

The fetal foramen ovale is physiologic

- During intrauterine life, interatrial right-to-left passage is normal.
- Pulmonary vascular resistance is high while the lungs are not functioning for gas exchange.
- This pressure setting favors right atrial pressure over left atrial pressure.



The pressure gradient regulates fetal passage

- Flow through the fetal atrial channel depends on right-versus-left atrial pressures.
- When right atrial pressure is higher, fetal interatrial flow is right to left.
- The channel is not equivalent to an irreversible tissue defect.

High fetal pulmonary resistance



Right atrial pressure relatively high



Right → left interatrial passage

Conceptual mechanism — editable schematic

Birth reverses atrial pressure relationships

- Lung expansion lowers pulmonary vascular pressure after birth.
- Right atrial pressure falls below left atrial pressure.
- The foramen ovale valve is pressed closed: Normal fetal communication is no longer required.

Lungs expand



Pulmonary pressure falls



LA pressure > RA pressure



Functional closure

Conceptual mechanism — editable schematic

Functional closure and anatomic sealing differ


- Pressure-driven apposition occurs with postnatal circulatory adaptation.
- Permanent fusion of the atrial flap occurs subsequently in most individuals.
- Failure of permanent fusion leaves PFO; Deficient septal tissue formation produces an ASD.

Transition	Structural result
Functional postnatal closure	Pressure-dependent apposition
Permanent anatomic fusion	Sealed normal channel
Failure of sealing	Patent foramen ovale

Native editable PowerPoint comparison table

Atrial septation: Developmental checkpoints

- First partition grows without immediately closing the atrial route.
- A secondary perforation preserves passage as the first gap disappears.
- Overlapping septa permit fetal flow and eventual postnatal closure.



Developmental or clinical stages — schematic

Where development fails determines morphology

- Deficient central septal development yields a secundum-type defect.
- A defect near the atrioventricular valve region is termed a primum ASD.
- A defect near the superior vena caval entry is termed sinus venosus ASD.

SECUNDUM

Center of septum

90%

PRIMUM

Near AV valves

5%

SINUS VENOSUS

Near SVC entry

5%

Robbins morphological classification

Three anatomical ASD categories

- Secundum: Central atrial septum.
- Primum: Adjacent to the atrioventricular valves.
- Sinus venosus: Near the entrance of the superior vena cava.

SECUNDUM

Center of septum

90%

PRIMUM

Near AV valves

5%

SINUS VENOSUS


Near SVC entry

5%

Robbins morphological classification


Secundum ASD is the predominant type

- Approximately 90% of ASDs to the secundum category.
- These arise from deficient septum secundum formation near the septal center.
- They are usually not associated with other congenital anomalies.




Secundum ASD can vary markedly in geometry

- A single defect may be small or large.
- Multiple perforations may coexist in the same septum.
- A fenestrated defect is therefore a recognized variation, not a separate main category.




Secundum location determines its anatomy

- The central atrial septum is the key anatomic landmark.
- The fixed communication connects both atria rather than ventricles or major arteries.
- Position is central; Multiplicity and size provide additional morphologic description.



Primum ASD is a low septal defect

- Approximately 5% of ASDs are described as primum anomalies.
- They occur next to the atrioventricular valves.
- Their location distinguishes them from a centrally situated secundum defect.



Primum ASD often accompanies other lesions

- Primum lesions with abnormalities of atrioventricular valves.
- They may also occur together with a ventricular septal defect.
- A primum finding warrants attention to nearby atrioventricular structures.

Low atrial septal defect



Adjacent AV valves




Look for AV valve anomaly ± VSD

Conceptual mechanism — editable schematic

Sinus venosus ASD lies near the SVC

- Approximately 5% of ASDs are sinus venosus defects.
- The defect is located near entry of the superior vena cava.
- Its anatomical position differs from central secundum and low primum defects.



Sinus venosus defects: Venous association

- Association with anomalous pulmonary venous return to the right atrium.
- The septal defect and venous drainage pattern are therefore linked morphologic observations.
- The association is not stated as obligatory for every sinus venosus defect.

Sinus venosus defect



Assess pulmonary venous connections




Possible return into RA

Conceptual mechanism — editable schematic

Relative frequencies of ASD variants

- Secundum dominates the distribution reported.
- Primum and sinus venosus each comprise a small minority.
- Percentages describe morphological categories, not severity or closure eligibility.



Proportions as, not population incidence

Locate the defect before describing its consequences


- Central defect: Secundum; Lower AV-valve region: Primum.
- High septal region near SVC: Sinus venosus.
- Record multiplicity and associated anomalies separately from type.

ASD type	Location	Key association
Secundum	Center	Usually isolated
Primum	Near AV valves	AV valve $\pm$ VSD
Sinus venosus	SVC entrance	Possible anomalous PV return

Native editable PowerPoint comparison table

Postnatal ASD flow is usually left to right

- The fixed opening permits blood to enter the right atrium from the left atrium.
- Low-resistance pulmonary circulation facilitates this direction.
- The resulting volume burden is largely on the right ventricle and pulmonary pathway.



Pulmonary versus systemic vascular resistance

- Pulmonary vascular resistance is much lower than systemic vascular resistance after birth.
- A route from the left side to the right side of the circulation therefore permits recirculation through the lungs.
- This is one major determinant of ASD shunting.

Low pulmonary resistance



LA → RA shunting



Pulmonary overcirculation

Conceptual mechanism — editable schematic

Relative ventricular compliance matters


- The right ventricle is more compliant (distensible) than the left ventricle.
- This favors filling and accommodation of additional right-sided volume.
- ASD shunt flow therefore reflects compliance as well as vascular resistance.

Physiologic feature	Effect on shunt
Lower PVR	Facilitates pulmonary recirculation
Higher RV compliance	Accommodates added volume
Fixed opening	Allows atrial communication

Native editable PowerPoint comparison table

Trace the extra blood volume through the circuit

- A portion of oxygenated left atrial blood returns to the right atrium.
- The right ventricle ejects this additional volume into the pulmonary trunk.
- The lungs receive more than the normal forward-flow volume.



Right-ventricular volume load is characteristic

- Robbins specifically distinguishes ASD from defects generating a strong right-ventricular pressure load.
- ASD principally augments right-ventricular and pulmonary outflow volumes.
- The remodeled right side may remain compensated for many years.

Left → right shunt



Extra RV filling



Increased RV outflow

Conceptual mechanism — editable schematic

Pulmonary blood flow may rise substantially

- Pulmonary flow volumes of two to eight times normal in ASD.
- This is a possible range described by the source, not a patient-specific measurement.
- High flow helps explain audible flow findings despite an initially well-tolerated lesion.

Atrial communication



Increased right-heart throughput




Pulmonary flow: 2–8× normal (textbook)

Conceptual mechanism — editable schematic

Why the pulmonary valve can produce a murmur

- Excess pulmonary outflow passes through an anatomically separate valve.
- A murmur may reflect increased flow through the pulmonary valve and/or the atrial defect itself.
- The sound is a hemodynamic clue, not a direct measure of defect diameter.



ASD differs from VSD and PDA hemodynamics

- ASD primarily raises right-ventricular and pulmonary flow volumes.
- VSD and PDA generally increase pulmonary blood flow and pressure.
- These differences influence downstream pulmonary vascular stress.

Defect	Pulmonary volume	Pulmonary pressure
ASD	Increased	Less dominant early feature
VSD	Increased	Also increased
PDA	Increased	Also increased

Native editable PowerPoint comparison table

Severity varies among left-to-right shunts

- The hemodynamic consequences depend on anatomy, size, and location.
- Left-to-right lesions may range from clinically silent to substantial hemodynamic disease.
- ASD is often tolerated longer than major pressure-loading lesions.

Defect anatomy



Shunt magnitude



Cardiac / pulmonary burden



Clinical impact

Conceptual mechanism — editable schematic

Why an uncomplicated ASD is acyanotic

- An initially left-to-right shunt does not introduce systemic venous blood directly into systemic arterial outflow.
- The additional circulating volume instead traverses the pulmonary circuit.
- Cyanosis implies another mechanism, such as a significant right-to-left component.

Direction	Immediate consequence
Left → right	Pulmonary overcirculation
Right → left	Desaturated blood enters systemic circulation

Native editable PowerPoint comparison table

Chronic shunt flow can remodel lung vessels

- Sustained pulmonary circulatory overload may elicit arteriolar medial hypertrophy and vasoconstriction.
- Longstanding exposure can generate obstructive intimal changes.
- This general left-to-right shunt pathway explains concern about irreversible pulmonary vascular disease.

Prolonged pulmonary overload



Medial hypertrophy + vasoconstriction



Obstructive intimal remodeling

Conceptual mechanism — editable schematic

Eisenmenger physiology is a late complication

- In severe longstanding left-to-right CHD, pulmonary resistance may approach systemic resistance.
- The shunt can reverse to right-to-left flow and produce hypoxemia and cyanosis.
- That irreversible pulmonary hypertension is unusual in ASD specifically.

High pulmonary resistance



Shunt reversal




Systemic desaturation

Conceptual mechanism — editable schematic

ASD is often clinically silent for decades

- ASDs are usually asymptomatic until adulthood.
- Symptoms commonly do not emerge before approximately age 30.
- Clinical quiescence should not be confused with an absence of ongoing shunt flow.



Developmental or clinical stages — schematic

Why ASDs commonly appear in adult practice

- Ventricular septal defects are more common at birth, but many close spontaneously.
- ASDs are less likely to close spontaneously in the comparison made by Robbins.
- ASDs consequently constitute an important defect detected in adults.

VSD	ASD
More common initially	Less common initially
Many close spontaneously	Less likely to close
Fewer persist to adult diagnosis	Often diagnosed in adulthood

Native editable PowerPoint comparison table

The clinical clue can be a flow murmur

- A murmur may be present despite absent symptoms.
- Increased pulmonary valve flow is a mechanism rather than proof that the valve is malformed.
- An observed murmur alone does not establish defect subtype.

Pulmonary overcirculation



Audible flow phenomenon



Evaluate underlying shunt anatomy

Conceptual mechanism — editable schematic

Why heart failure can eventually occur

- The atrial communication perpetuates right-sided volume overload.
- A chronically stressed cardiac circulation can eventually become unable to maintain compensation.
- Preventing heart failure is one stated rationale for ASD closure.

Long-term shunt



Right-sided volume burden




Potential cardiac failure

Conceptual mechanism — editable schematic

Paradoxical embolization requires a right-to-left route

- Venous embolic material normally encounters the pulmonary circulation.
- Interatrial right-to-left passage can provide access to systemic arterial blood.
- Avoiding paradoxical embolization is an explicit rationale for ASD closure.



Pulmonary vascular disease is a preventable concern

- Chronic abnormal pulmonary flow can produce structural vascular adaptation.
- Late obstructive vascular remodeling may be irreversible.
- Pulmonary hypertension as unusual in ASD but relevant to closure timing.

Abnormal pulmonary flow



Vascular remodeling




Risk of irreversible disease

Conceptual mechanism — editable schematic

Closure targets future harm, not just current symptoms

- ASD closure is undertaken to prevent later heart failure, embolic events, and permanent pulmonary vascular injury.
- Surgical and intravascular closure approaches.
- Clinical benefit is assessed in the context of the defect and its physiologic consequences.



Two closure routes are described


- Surgical repair and intravascular approaches both seek to eliminate the abnormal communication.
- Anatomical type and associated lesions are relevant to procedural evaluation.
- Robbins does not provide a detailed device-selection or procedural eligibility algorithm in this excerpt.

Approach	Textbook scope
Surgical closure	Named option
Intravascular closure	Named option
Patient selection	Not specified in excerpt

Native editable PowerPoint comparison table

Reported outcome after closure is favorable

- Robbins characterizes mortality after ASD closure as low.
- Postoperative survival is reported as comparable to an unaffected population.
- These are general textbook statements; Individual outcomes depend on clinical circumstances.



PFO: Postnatal sealing is frequently incomplete

- Permanent foramen ovale closure in approximately 80% by age two.
- In about 20%, an unsealed flap-like channel persists.
- This prevalence refers to PFO, not to the frequency of a true ASD.

Closed by age ~2	Unsealed channel persists
Approximately 80%	Approximately 20%
Normal completed fusion	Patent foramen ovale

Native editable PowerPoint comparison table

PFO can open during transient pressure increases

- Coughing, sneezing, or straining can raise right-sided pressure transiently.
- A nonfused flap can open when right atrial pressure exceeds left atrial pressure.
- Brief right-to-left shunting is then possible.

Transient RA pressure rise



PFO flap opens



Brief right → left flow

Conceptual mechanism — editable schematic

PFO and ASD: Separate mechanisms for embolic risk

- PFO creates an intermittent right-to-left channel under suitable pressure conditions.
- ASD is a fixed opening with a usual left-to-right shunt, but paradoxical embolization remains a recognized concern.
- The presence, direction, and hemodynamic context of an interatrial passage determine risk mechanisms.

PFO	ASD
Flap opens under RA pressure	Fixed abnormal defect
Transient right → left episodes	Usually left → right flow
May permit paradoxical embolus	Closure partly prevents embolic risk

Native editable PowerPoint comparison table

Case: An adult with an incidental murmur

- A person in early adulthood feels well but has a persistent flow murmur on examination.
- The main question is whether a congenital left-to-right communication increases pulmonary output.
- ASD can remain silent for decades; Lack of symptoms does not negate the possibility.

Why might a substantial ASD be detected only in adult life?

Interpretation

A compliant right ventricle tolerates additional volume, and irreversible pulmonary hypertension is uncommon early.

Case: A low atrial septal opening

- A fixed opening is seen near the atrioventricular valve plane.
- This location favors a primum defect rather than a centrally placed secundum ASD.
- The nearby valves and ventricular septum require careful assessment for associated anomalies.

Which structures are most relevant to inspect next?

Interpretation

Atrioventricular valves and the ventricular septum.

Case: A high septal communication

- A defect lies near the superior vena cava entrance.
- This localizes to the sinus venosus category.
- The possibility of pulmonary veins draining anomalously to the right atrium should be considered.

Which additional anatomical connection matters?

Interpretation

Pulmonary venous return.

Case: Brief right-to-left passage during straining

- The interatrial communication appears flap-like rather than a fixed septal tissue gap.
- A transient pressure rise on the right opens the passage.
- This is characteristic of PFO physiology, not evidence of secundum ASD.

What distinguishes this lesion from an ASD?

Interpretation

The channel is a nonfused fetal flap, not a fixed deficiency of septal tissue.

Check understanding: Shunt reversal

- A longstanding left-to-right lesion develops progressively elevated pulmonary vascular resistance.
- When pulmonary resistance becomes sufficiently high, right-to-left passage may occur.
- This defines the physiology of Eisenmenger syndrome; It is unusual in ASD.

Would this be an expected early ASD presentation?

Interpretation

No. ASD generally causes a tolerated left-to-right shunt; Irreversible pulmonary hypertension is unusual.

Diagnostic interpretation: Read anatomy and physiology together


- Location classifies the defect; Flow direction explains its immediate hemodynamic effects.
- Associated atrioventricular or pulmonary venous anomalies can change the anatomical context.
- Clinical manifestations cannot be inferred reliably from subtype alone.

Question	What it establishes
Where is the opening?	ASD morphological type
Which way does blood move?	Physiologic shunt direction
Which structures are associated?	Accompanying anomalies

Native editable PowerPoint comparison table

ASD: Integrated mechanism in one sequence

- A developmental septal deficiency produces a fixed communication between the atria.
- Low pulmonary resistance and greater RV compliance favor left-to-right shunting.
- Pulmonary overcirculation is initially tolerated but motivates preventive closure.



High-yield anatomical synthesis

- Secundum is central and represents about nine of ten ASDs.
- Primum is adjacent to the AV valves and may accompany AV anomalies or VSD.
- Sinus venosus is close to the SVC and may coexist with anomalous pulmonary venous return.

SECUNDUM

Center of septum

90%

PRIMUM

Near AV valves

5%

SINUS VENOSUS

Near SVC entry

5%

Robbins morphological classification

High-yield clinical synthesis

- Most ASDs are well tolerated until adult life; A flow murmur may be present.
- Increased RV and pulmonary volumes are more characteristic than an early high-pressure shunt.
- Closure aims to prevent failure, paradoxical embolization, and irreversible pulmonary vascular disease.

Often silent



RV volume overload + pulmonary flow



Prevent future complications

Conceptual mechanism — editable schematic

Reference and further reading

- Robbins, Cotran & Kumar: Pathologic Basis of Disease, 11th edition (2026).
- Read the sections on congenital heart disease, atrial septal defect, and patent foramen ovale together.
- Review the textbook diagrams and gene-association tables alongside the conceptual drawings in this lecture.