
Autoimmune Enteropathy

Immune dysregulation, intestinal injury, and clinicopathologic reasoning



Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning goals

- Define the syndrome and its relationship to IPEX.
- Explain how impaired regulatory T cells permit multisystem autoimmunity.
- Interpret the reported intestinal inflammatory pattern.
- Distinguish major mimics and connect findings to therapeutic principles.


Disease concept: Persistent diarrhea with autoimmunity

Severe, persistent diarrhea is the defining gastrointestinal presentation.

Other autoimmune manifestations provide the systemic context.

The Robbins account emphasizes presentation in young children.

A syndrome, not simply isolated bowel inflammation



Mechanistic schematic

Clinical timing: Early life is a major clue

- Young children are especially emphasized in the source description.
- The combination of early onset and severe persistence raises the importance of inherited immune dysregulation.
- Age is contextual—not a stand-alone diagnostic criterion.

Why “persistent” diarrhea matters

Transient illness

A short-lived episode alone cannot establish the AIE syndrome.

Persistent syndrome

Ongoing diarrhea plus autoimmunity prompts a broader evaluation.

The intestine is one site of systemic autoimmunity

**Gastrointestinal
involvement**

Immune dysregulation

**Other autoimmune
disease**

Conceptual grouping

IPEX expands the clinical phenotype

I

Immune dysregulation

P

Polyendocrinopathy

E

Enteropathy

X

X-linked

IPEX is the severe familial form highlighted by Robbins

- A particularly severe inherited form of autoimmune enteropathy.
- Associated with germline loss-of-function variants in FOXP3.
- Intestinal manifestations are part of a broader autoimmune syndrome.

The causal gene in IPEX is FOXP3



X-linked inheritance: What the source establishes

Genetic location

FOXP3 is located on the X chromosome.

Disease relationship

A germline loss-of-function mutation is the basis of the IPEX form.

FOXP3 is a transcription factor

The encoded protein regulates gene expression.

It is expressed in CD4+ regulatory T cells.

Deficiency compromises development and function of this regulatory lineage.

Regulatory T cells protect against self-reactivity



Phenotype of the best-defined regulatory T cells

CD4+ T cell

High CD25

FOXP3 expression

Conceptual grouping

Regulatory T-cell ontogeny

Thymus

Regulatory T cells develop mainly in the thymus during recognition of self antigens.


Periphery

Additional regulatory T cells can be induced in peripheral lymphoid tissues.

IL-2 is required for regulatory T-cell maintenance



CTLA-4 limits T-cell costimulation



Mechanistic schematic

Regulatory cytokines suppress effector responses

IL-10

An immunosuppressive cytokine implicated in regulatory T-cell function.

TGF- β

Also helps inhibit lymphocyte activation and effector functions.


Immune tolerance is an active process

Regulatory T cells restrain immune reactions to self antigens.

Loss of restraint can permit multiorgan inflammation.

Intestinal injury in IPEX is one manifestation of this failure.

From FOXP3 loss to enteropathy



Not all cases are attributed to FOXP3

- FOXP3 loss of function identifies the IPEX-associated form.
- Robbins also links other regulatory T-cell defects to less severe autoimmune enteropathy.
- Do not equate the entire disease spectrum with one genotype.

A genotype is not identical to a histologic pattern

Molecular level

FOXP3 mutation establishes a particular etiologic pathway.

Tissue level

Biopsy demonstrates mucosal changes that are not gene-specific.

Autoantibodies show the breadth of autoreactivity

E

Enterocyte antibodies

G

Goblet-cell antibodies

P

Parietal-cell antibodies

I

Islet-cell antibodies

Enterocyte-directed autoantibodies

- Antibodies directed against enterocytes are common in AIE.
- The target corresponds to the intestinal absorptive epithelial compartment.
- An associated autoantibody finding must be integrated with clinical and morphologic context.

Goblet-cell-directed autoantibodies

- Anti-goblet-cell antibodies are also reported commonly.
- Goblet cells are part of the intestinal epithelial compartment.
- This finding supports an autoimmune phenotype but is not given as a stand-alone diagnostic criterion.

Extraintestinal autoantibodies

Parietal cells

Autoantibodies may be detected in some patients.

Islet cells

Autoantibodies may also be detected in some patients.


Autoantibody patterns are supportive observations

Associations, not formal antibody-based diagnostic criteria.

Antibody detection alone does not explain every clinical manifestation.

Keep the phenotype, histology, and inherited-regulation pathway separate.

Small-intestinal mucosal compartments



Epithelial mantle



IELs inside epithelial lining



Lamina propria in villous core

Schematic — normal compartments, not AIE histology

Intraepithelial lymphocytes may increase

- IELs are lymphocytes within the epithelial layer.
- In AIE, an increase may be present.
- This rise is typically less extensive than in celiac disease.

Neutrophils offer a contrasting clue

- Neutrophilic infiltration of intestinal mucosa is often reported in AIE.
- The source explicitly contrasts this with celiac disease.
- Interpret the neutrophils as part of a pattern, never an exclusive marker.

Interpret both immune-cell populations together

Feature	Autoimmune enteropathy	Celiac disease comparison
Intraepithelial lymphocytes	May increase	Generally more extensive
Neutrophils in mucosa	Often seen	Contrasted as less typical
Specificity	Neither cell type proves AIE	Neither single finding proves celiac

Histology: Reported findings and limitations


Describes

Possible IEL increase and frequent mucosal neutrophils.

Does not establish

A required villous grade, a numeric histologic cutoff, or a pathognomonic lesion.

Tissue findings require clinical interpretation




The physiology of malabsorption

Malabsorption is defective absorption of nutrients, electrolytes, and water.

Stool and nutritional effects may converge across distinct etiologies.

Symptoms alone therefore identify the problem imperfectly.

Four physiological stages of nutrient absorption



Diarrhea has more than one mechanism

Category	Textbook descriptor
Secretory	Persists during fasting
Osmotic	Unabsorbed luminal solutes; improves with fasting
Malabsorptive	Generalized absorptive failure; often steatorrhea
Exudative	Inflammatory, sometimes purulent or bloody

Why malabsorptive conditions look similar

- Different defects can converge on diarrhea, weight loss, and abdominal distention.
- Failure in one absorption stage can coexist with defects in another.
- The clinical pattern must be interpreted alongside disease-specific findings.

Systemic consequences of chronic malabsorption

Absorptive deficit	Potential clinical consequence
Iron, folate, or vitamin B12	Anemia or mucosal abnormalities
Vitamin K	Bleeding tendency
Calcium, magnesium, vitamin D	Bone or neuromuscular consequences
General nutrient deficit	Weight loss / muscle wasting

Why the autoimmune context changes interpretation

Intestinal physiology

Diarrhea and nutritional failure can follow many mucosal disorders.

Immune phenotype

Coexisting autoimmune disease and Treg defects narrow mechanistic hypotheses.

Celiac disease is a crucial immune-mediated comparator


Celiac disease

Gluten-driven enteropathy in a genetically susceptible host.

Autoimmune enteropathy

Regulatory immune dysfunction, with IPEX as a major example.

Celiac disease has a defined antigenic trigger



Celiac mucosa has a characteristic combination

Increased IELs

Crypt hyperplasia

Villous atrophy

Conceptual grouping

Keep celiac-specific morphology distinct

Celiac disease is characterized by increased IELs, crypt hyperplasia, and villous atrophy.

The AIE paragraph emphasizes a less extensive IEL increase and frequent neutrophils.

Shared inflammatory changes must be interpreted with disease-specific clinical context.

Celiac serology addresses a different pathway

- Anti-TG2 IgA and antiendomysial antibodies are described for celiac disease.
- Celiac evaluation integrates serology with histology.
- These are not the enterocyte/goblet-cell autoantibodies listed under AIE.

Autoimmune enteropathy versus celiac disease

Dimension	Autoimmune enteropathy	Celiac disease
Main context	Autoimmunity / Treg defects	Dietary gluten exposure
Established gene example	FOXP3 in IPEX	HLA-DQ2 / HLA-DQ8 predisposition
Mucosal emphasis	Neutrophils; variable IELs	Prominent IELs, crypts, villi
Therapy in source	Immune suppression; HSCT	Gluten-free diet

Lactase deficiency: An osmotic comparator

Mechanism

Absent / reduced brush-border
lactase retains unabsorbed lactose.

Distinguishing context

Dietary lactose association;
Histology generally unremarkable.

Microvillus inclusion disease: A transport comparator

MYO5B-associated
trafficking defect



Defective brush-border
assembly



Severe early diarrhea

Abetalipoproteinemia: A lipid-transport comparator

MTP deficiency



Defective apolipoprotein B
lipoprotein assembly



Fat malabsorption and
steatorrhea

Environmental enteric dysfunction: Different context

Exposure

Linked to poor sanitation, repeated enteric exposure, and mucosal dysfunction.

Evidence gap

No accepted single clinical, laboratory, or histologic criteria.

Infection can also produce epithelial injury

- Enteric infection is an important competing cause of diarrhea.
- The celiac section explicitly notes that IEL increases and villous atrophy are not exclusive to celiac disease.
- A shared mucosal abnormality cannot identify immune dysregulation by itself.

Intestinal graft-versus-host disease

Clinical setting

A recognized cause of malabsorption and diarrhea after allogeneic stem-cell transplantation.

Reasoning point

Treatment history can matter as much as the biopsy pattern in enteropathy.

A structured enteropathy differential

Cause	Mechanistic discriminator	Helpful context
IPEX/AIE	Treg immune dysregulation	Autoimmunity; FOXP3 form
Celiac	Gluten-directed response	TG2 serology, histology
Lactase deficiency	Osmotic carbohydrate defect	Lactose-related symptoms
Microvillus inclusion	Epithelial trafficking defect	MYO5B; early severe diarrhea
Abetalipoproteinemia	Lipoprotein assembly defect	Steatorrhea; lipid transport

Organize the diagnostic information by domain

Phenotype and age

Biopsy pattern

**Immune / genetic
context**

Conceptual grouping

Pattern recognition has limits

- Neither increased IELs nor neutrophils alone establishes a diagnosis.
- The source does not establish a mandatory numeric histologic cutoff for AIE.
- Do not substitute a celiac biopsy figure for evidence of AIE morphology.

AIE treatment: Immunosuppression

Regulatory failure and
autoimmunity



Immune-directed therapy



Example reported:
Cyclosporine

Allogeneic hematopoietic stem-cell transplantation

Listed by Robbins as a therapy for autoimmune enteropathy.

Represents an immune-system-directed intervention in selected contexts.

The excerpt does not provide selection criteria, risks, success rates, or timing.

Keep therapeutic context and differential distinct

Therapeutic role

Allogeneic transplantation is included among treatments for AIE.

Potential competing etiology


Graft-versus-host disease can cause post-transplant enteropathy.

Hypothetical case: Enteropathy plus systemic autoimmunity

Hypothetical teaching vignette

- A young child has severe persistent diarrhea and additional autoimmune manifestations.
- Genetic testing identifies a germline loss-of-function FOXP3 variant.
- Question: What unifies the gastrointestinal and systemic findings?

Case interpretation: Defective immune regulation



Hypothetical case: Intestinal lymphocytosis

Hypothetical teaching vignette

- A patient with chronic diarrhea has increased intraepithelial lymphocytes.
- No immune phenotype or disease-specific serology has yet been characterized.
- Question: Can this biopsy finding alone identify AIE or celiac disease?

Case interpretation: Seek discriminating evidence

- No: Intraepithelial lymphocytosis is not specific to either diagnosis.
- Celiac-specific history, serology, and mucosal features require assessment.
- AIE requires interpretation in the context of immune dysregulation and its reported inflammatory pattern.

Hypothetical case: Severe diarrhea very early in life

Hypothetical teaching vignette

- An infant develops intractable diarrhea during the first months of life.
- The main question is whether the defect is immune or epithelial.
- Which alternative textbook disorder involves defective brush-border membrane trafficking?


Case interpretation: MYO5B-related enteropathy

- Microvillus inclusion disease results from defective MYO5B-related trafficking.
- It may produce severe diarrhea in early infancy.
- An early age alone therefore cannot distinguish inherited epithelial disease from immune dysregulation.

Common diagnostic reasoning pitfalls

Pitfall	Correction
Every AIE case = FOXP3 mutation	IPEX is one severe familial form
IEL rise alone identifies disease	IEL increase occurs in multiple disorders
Celiac features automatically identify AIE	Keep entity-specific histology separate
Autoantibody = definitive diagnosis	Integrate clinical, tissue, immune context

Mechanism-to-phenotype synthesis



Mechanistic schematic

Four points to retain

- IPEX is a severe X-linked FOXP3-related form of AIE.
- FOXP3-dependent regulatory T cells restrain autoimmunity.
- AIE may show IEL increases and often mucosal neutrophils.
- Diagnosis is integrative; Treatment examples include cyclosporine and allogeneic HSCT.

Primary references and source boundaries

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th edition (2026).
- Chapter 17: The Gastrointestinal Tract — autoimmune enteropathy, celiac disease, and malabsorption sections.
- Chapter 6: Diseases of the Immune System — regulatory T cells and peripheral tolerance.
- Morphology and diagnostic reasoning require correlation with the full clinical and immunologic context.