

Autophagy

Cellular self-digestion, stress adaptation and the paradox of cancer survival

Dr Abu Suraih Sakhri


MBBS, MD (Pathology)

www.abusuraihsakhri.com

What this lecture will establish

- Describe the formation and lysosomal maturation of an autophagosome.
- Explain why recycling supports cells during nutrient deprivation and hypoxia.
- Distinguish tumor-suppressive from tumor-supportive effects of autophagy.
- Relate autophagy to growth metabolism, dormancy and therapeutic resistance.

Autophagy is a cellular survival decision



- Starved cells arrest growth while salvaging carbon and essential metabolites.
- If the adaptation cannot sustain viability, the cell may die.

Definition: Lysosome-mediated self-digestion

- Autophagy delivers cytoplasmic material to lysosomes for degradation.
- “Self-eating” describes utilization of intracellular contents, not engulfment of another cell.
- The resulting breakdown products re-enter cellular metabolism.

What can enter the autophagic pathway?

Structural material

Damaged or obsolete organelles

Membrane material

Cytoplasmic constituents

Macromolecular cargo

Aggregated proteins

Protein complexes

Intracellular debris


Recovered value

Smaller metabolites

Substrates for maintenance

Carbon for energy production

Nutrient deprivation is a major trigger



- Autophagy is evolutionarily conserved as an adaptive response to starvation.
- Its survival benefit is especially apparent when external substrates are scarce.

Hypoxia creates parallel metabolic pressure

- Cells under inadequate oxygen supply experience metabolic stress.
- Hypoxia as a pathologic trigger of autophagy.
- Autophagy can help sustain function by recovering intracellular substrates.

Autophagy also responds to nonnutritional stress

Physiologic settings

Aging

Exercise

Turnover of aging cell structures


Pathologic settings

Oxidative stress

Organelle or membrane damage

Hypoxia and limited nutrients

Stress response is not unlimited rescue



- Recycling delays injury when external nutrients cannot meet cellular needs.
- Autophagy does not guarantee indefinite survival under persistent severe stress.

Autophagy is one route to the lysosome

Autophagy

Intracellular organelles and aggregates

Double-membrane autophagosome

Fusion with lysosome

Heterophagy

Material acquired from outside the cell

Endosome or phagosome

Fusion with lysosome

Autophagy and proteasomal degradation differ

Autophagic–lysosomal route

Captures organelles or larger assemblies

Membrane enclosure precedes digestion

Lysosomal enzymes degrade contents


Ubiquitin–proteasome route

Processes many cytosolic proteins

Ubiquitin marks target proteins


Proteasomal complexes generate peptides

Three core compartments organize the process



- A forming membrane sequesters cargo before lysosomal digestion.
- The final vesicle is a processing compartment rather than a permanent store.

Phagophore: The isolation membrane




Growing isolation membrane

- The phagophore is an early isolation membrane that curves around cytoplasmic cargo.
- It enlarges before closing into a complete vesicle.
- Schematic: Membrane arrangement, not a histologic specimen.

The endoplasmic reticulum is a membrane source

- The endoplasmic reticulum as a probable source of isolation membrane.
- Plasma membrane and mitochondrial membranes may also contribute.
- The phagophore therefore originates within the cell's membrane system.

ATG proteins regulate assembly in sequence




- More than 30 autophagy-related (ATG) genes.
- Their products coordinate creation of an autophagosome.

Initiation responds to environmental cues


- Nutrient deprivation and reduced growth-factor stimulation activate autophagy initiation.
- An initiation complex recruits the proteins required to assemble the isolation membrane.
- This is a regulated program, not uncontrolled membrane fragmentation.

Nucleation creates a membrane platform



- Nucleation establishes the membrane region that will grow around material.
- Subsequent expansion requires additional ATG-dependent reactions.


Elongation progressively sequesters cargo



Growing isolation membrane

- The open membrane expands around organelles and cytosolic structures.
- Continued elongation eventually enables membrane closure.
- Enclosure separates future digestive cargo from surrounding cytoplasm.

LC3 lipidation helps build the autophagosome




- Two ubiquitin-like conjugation systems participate in membrane expansion.
- Phosphatidylethanolamine (PE) is covalently linked to LC3.

LC3 has a structural and tracking role

- Lipidated LC3 contributes to enclosure of the forming autophagosome.
- The amount of PE-linked LC3 increases during autophagy.
- LC3 as a useful marker for recognizing autophagy-related structures.


Closure forms a double-membrane autophagosome




Sequestered cytoplasmic cargo

- After enclosure, captured cytoplasm lies within a sealed double-membrane vesicle.
- Cytoplasmic organelles and other structures can be enclosed.
- The autophagosome must mature before its contents are degraded.

Lysosome fusion is necessary for maturation




Autophagosome



Lysosome

- The newly formed autophagosome fuses with the lysosomal compartment.
- The resulting autophagolysosome gains access to hydrolytic enzymes.
- Fusion connects sequestration to actual cargo destruction.


Lysosomal hydrolases digest the contents



Lysosomal hydrolase action


- The inner autophagosomal membrane and sequestered contents are degraded.
- Lysosomal enzymes convert complex cellular material into smaller products.
- The process supports recovery rather than simple intracellular storage.

Recycled metabolites support continued function



- Cells can use recovered material when supply from the environment is limited.
- This is a catabolic contribution to cellular energy and substrate balance.

The complete pathway is a staged process



- Sequestration and degradation are mechanistically distinct stages.
- The pathway depends on organized membrane trafficking and lysosomal function.

Autophagic cargo selection is not wholly random

- That autophagy can selectively capture cytoplasmic targets.
- Lipidated LC3 participates in targeting aggregates and obsolete organelles.
- Target selection helps remove constituents that threaten cellular integrity.

Autophagy supports organelle quality control

Examples of turnover

Endoplasmic reticulum

Mitochondria

Lysosomes


Functional consequence

Removes worn or damaged structures

Maintains intracellular organization

Supports cellular stress tolerance

Intracellular aggregates can be removed



- Stress and aging can increase intracellular protein aggregates.
- Autophagic clearance is one mechanism that limits this burden.

Atrophic cells use nutrient salvage

- Autophagy is prominent in cells undergoing severe nutrient deprivation.
- Self-digestion supports maintenance even while cell mass is being lost.
- Atrophy and adaptive recycling may therefore coexist.

LC3 positivity requires cautious interpretation

What the source supports

LC3 is linked to autophagosome membranes

Lipidated LC3 increases during autophagy

It is useful as an autophagy marker


What a static observation cannot establish by itself

How much material was fully degraded

Whether fusion with lysosomes succeeded

Whether autophagy benefits a particular tumor

Prolonged severe stress can overwhelm adaptation



- Excessive autophagy settings to apoptosis, necroptosis and ferroptosis.
- These associations do not prove autophagy directly caused cell death.

Autophagy-associated death remains mechanistically debated

- Cell death can accompany marked or prolonged autophagic activity.
- The responsible insult may be the original stressor rather than the autophagic process.
- Do not equate the observation of autophagosomes with an established cause of death.


Disease associations extend beyond neoplasia

Disease context	Autophagy-related concern
Neurodegenerative disorders	Defective proteostasis and turnover
Infectious diseases	Microbial degradation and host defense
Inflammatory bowel diseases	Genetic susceptibility associations
Cancer	Both suppression and survival benefit

Neurodegeneration links clearance to proteostasis


- Impaired organelle turnover can aggravate neuronal dysfunction.
- Abnormal protein handling becomes more important in long-lived cells.
- Defective autophagy with several neurodegenerative disorders.

Alzheimer disease: Maturation can be impaired



- Robbins specifically describes impaired autophagosome maturation in Alzheimer disease.
- Genetic autophagy defects accelerate neurodegeneration in cited mouse models.

Autophagy contributes to control of intracellular pathogens



- The source cites mycobacteria, Shigella species and HSV-1 as examples.
- Autophagy can deliver microbial material to antigen-presentation pathways.

Inflammatory bowel disease has an ATG16L1 association

- Genome-wide association studies link ATG16L1 variants with Crohn disease and ulcerative colitis.
- The precise mechanism promoting intestinal inflammation is unknown.
- Genetic association does not automatically establish a complete causal mechanism.

Normal autophagy integrates catabolism and quality control

During limited nutrients

Recovers carbon and metabolites

Buffers acute metabolic deprivation

Helps maintain cellular viability

During cellular damage

Eliminates damaged organelles


Disposes of aggregates

Supports intracellular homeostasis

Cancer cells confront an unstable microenvironment

- Tumor cells often encounter limited nutrients and hypoxia.
- Their growth demands biosynthetic material while stress threatens survival.
- Autophagy may be helpful, dispensable or suppressed depending on tumor context.

Some tumors appear to evade autophagy induction




- Tumors that grow under marginal conditions without triggering autophagy.
- This raises the possibility of impaired autophagy-inducing pathways.

Certain pro-autophagy genes act as tumor suppressors


- The source identifies a subset of genes promoting autophagy as tumor suppressors.
- Loss of such functions may enhance tumor growth.
- The excerpt does not justify claiming that every autophagy gene is tumor suppressive.

Reduced autophagy can favor tumor development



- Autophagy loss as one plausible cancer-promoting mechanism.
- This is a context-dependent model, not a universal causal sequence.

Autophagy can also be a tumor survival mechanism




- Cancer cells can use the same nutrient-salvage program as normal cells.
- This survival function complicates the concept of autophagy as purely tumor-suppressive.

Tumor dormancy creates a prolonged survival state

- Dormancy as hibernation-like survival during adverse conditions.
- Autophagy may sustain quiescent cancer cells through prolonged hardship.
- Persistence need not be accompanied by active tumor-cell division.

Dormant cells may evade proliferation-targeting treatment



- This is a proposed explanation for some treatment failures in the source.
- The textbook uses cautious language: Such cells are believed to be resistant.

Autophagy may be a friend or a foe of a tumor

Autophagy in cancer

```
graph TD; A[Autophagy in cancer] --> B[Tumor-suppressive setting]; A --> C[Tumor-supportive setting];
```

Tumor-suppressive setting

- Protective genes oppose tumor development
- Loss can support growth
- Effect varies across tumors

Tumor-supportive setting

- Recycling under metabolic stress
- Support of long-term dormancy
- Potential treatment persistence

Tumor context determines the net effect

- The consequences depend on the molecular wiring of regulatory pathways.
- The same autophagic machinery may have opposing roles at different tumor stages.
- A binary “autophagy always promotes cancer” claim contradicts Robbins.

Two hypothetical tumors can behave differently

Tumor model A

Protective autophagy induction is defective

Loss may favor tumor growth

Restoring protection is conceptually relevant

Tumor model B

Severe nutrient stress persists

Autophagic recycling sustains viability

Blocking adaptation could have different effects

A static LC3 result cannot assign tumor prognosis

- LC3 is useful for recognizing autophagy-related structures, not a stand-alone cancer behavior score.
- A tumor's clinical phenotype depends on pathway context.
- The supplied excerpt provides no validated diagnostic cutoff or predictive threshold.

Pro-growth metabolism is the adjacent context

Growth-driven state

Uptake of glucose and glutamine

Biosynthesis of nucleotides, proteins and lipids

Oncogenic growth signaling


Nutrient-stressed state

Reduced available external substrates

Growth arrest / adaptation

Autophagic salvage of intracellular carbon

Aerobic glycolysis supports biosynthesis




- The Warburg effect is aerobic glycolysis in growing cells.
- This program is not unique to malignant cells.

Growing cells need carbon skeletons, not only ATP


- A dividing cell must duplicate nucleic acids, proteins, lipids and organelles.
- Glycolytic and mitochondrial intermediates supply biosynthetic precursors.
- This helps explain the metabolic program of rapidly proliferating cells.

PI3K–AKT signaling amplifies anabolic metabolism



- PI3K–AKT signaling to increased glucose transport and glycolysis.
- The same pathway supports lipid and protein synthesis.

MYC reinforces the growth-metabolism program



- MYC promotes gene expression supporting glycolysis and glutamine metabolism.
- These effects support the biomass requirements of growing cells.

Tumor suppressors oppose progrowth metabolism

Pro-growth drivers

RTK / PI3K / AKT

MYC

Oncogenic signaling

Counterbalancing factors

PTEN opposes PI3K–AKT

P53 restrains anabolic routes

STK11 antagonizes metabolic changes

Krebs-cycle enzyme alterations can contribute to cancer

- Fumarate hydratase (FH) and succinate dehydrogenase (SDH) as tumor suppressors.
- Inherited loss-of-function is associated with predisposition to selected tumors.
- The mechanisms remain incompletely resolved and may involve abnormal metabolites.

Autophagy and oncometabolism are distinct concepts

Autophagy

Catabolic processing of intracellular material

Recycles metabolites under stress

May restrain or support tumors


Oncometabolism

Oncogenic effects of altered metabolite pathways

Enzyme mutations can create oncometabolites


Changes gene regulation and tumor behavior

Mutant IDH produces an abnormal metabolite



- Mutant IDH acquires a neomorphic enzymatic function.
- The aberrant product 2-hydroxyglutarate (2-HG) is an oncometabolite.

2-hydroxyglutarate perturbs epigenetic control



- The text highlights TET2 as an affected α -ketoglutarate–dependent enzyme.
- Abnormal methylation patterns provide one route to oncogenesis.

IDH mutation creates a therapeutic target

- Mutant IDH proteins differ structurally and functionally from normal IDH.
- Drugs designed to inhibit mutant IDH selectively.
- This example illustrates mechanism-driven cancer therapy, not a direct autophagy-targeted treatment.

Two adaptations answer different cellular pressures

Cell seeking proliferation

Turns external glucose and glutamine into precursors

Requires oncogenic growth signaling

Builds biomass for division

Cell seeking persistence

Recycles internal proteins and organelles

Responds to nutrient and oxygen stress

Conserves viability while growth slows

Case exercise: Starved nonmalignant cell

A nonmalignant cell is deprived of nutrients but remains viable for a limited period despite growth arrest.

Discussion prompts

- Explain how autophagosomes and lysosomes contribute to viability.
- Identify what is recycled and why this cannot sustain the cell forever.
- Predict the consequence if this adaptation fails.

Case exercise: Nutrient-limited residual tumor

Following treatment, a small population of cancer cells persists in a nutrient-limited environment without active proliferation.

Discussion prompts

- Describe the putative role of autophagic recycling.
- Explain why nondividing cells might escape treatments directed at proliferation.
- State which part of this interpretation remains hypothetical.

Interpretation of the dormant-tumor case

- Autophagy may provide carbon sources that help dormant cells remain viable.
- Low proliferation could reduce susceptibility to drugs acting mainly on dividing cells.
- This as a plausible contributor to therapeutic failure, not an established universal rule.

Common interpretive errors

Overstatement

More LC3 proves stronger recycling

Autophagy always kills tumor cells

All cancers have autophagy disabled

Dormancy guarantees drug resistance

Evidence-concordant statement


LC3 supports membrane-pathway assessment

Effects on tumors can be bidirectional

Regulatory wiring varies by tumor

Dormancy is a proposed persistence mechanism

Mechanistic checkpoints for analysis



- Different questions address initiation, execution and biological impact.
- A descriptive marker cannot by itself answer all four questions.

Self-assessment: Resolve the paradox

Statement: “Autophagy is tumor suppressive; Therefore, inducing it always improves cancer control.”

Discussion prompts

- Identify the part supported by Robbins.
- Identify the missing tumor-specific qualifier.
- Explain how dormancy changes the inference.

Resolution: The outcome depends on context

Autophagy in cancer

```
graph TD; A[Autophagy in cancer] --> B[Potentially restrains cancer]; A --> C[Potentially sustains cancer];
```


Potentially restrains cancer

- Some pro-autophagy genes are suppressors
- Loss may favor tumor growth
- Early protective role is plausible

Potentially sustains cancer

- Metabolic salvage during hardship
- Persistence of dormant cells
- Possible contribution to treatment failure

Integrated mechanism: Stress, recycling and fate



- Autophagy is a controlled recycling program.
- In cancer, the biological consequence depends on regulatory wiring and state.

Five statements to retain

- Autophagy encloses cytoplasmic material and delivers it to lysosomes.
- ATG-dependent membrane formation and LC3 lipidation organize sequestration.
- Recycled metabolites support cells under nutrient deprivation or hypoxia.
- Cancer may be restrained by autophagy loss-sensitive pathways or supported by stress recycling.
- Dormancy makes simple treatment predictions unsafe.

Primary references and source boundaries

Primary

Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. (2026). Chapter 7: Neoplasia, p. 269 — Autophagy.

Foundational

Same textbook, Chapter 2: Cell Injury, Cell Death, and Adaptations, pp. 34–35 — Autophagy and Figure 2.5.

Context

Same textbook, Chapter 1 — Intracellular Catabolism (Figure 1.10); Chapter 7, pp. 267–270 — Growth-Promoting Metabolic Alterations and Oncometabolism.