

Cancer Risks From Exposures to Radiation

Molecular mechanisms, human evidence, and clinical risk interpretation

Ionizing exposure



DNA damage



Surviving mutated cell



Delayed cancer

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Trace ionizing radiation from DNA injury to delayed malignant transformation.
- Interpret evidence at high doses and uncertainty at lower exposures.
- Identify cancer patterns after fallout, radiotherapy, imaging, and radon.
- Explain why pediatric exposure and exposure context influence risk.

Radiation-associated cancer is a delayed outcome

- Ionizing radiation can produce mutations in cells that remain viable and capable of division.
- Such cells can contribute to neoplastic transformation long after the exposure.
- Population studies demonstrate excess leukemias and solid tumors after major exposures.
- Immediate tissue injury and late cancer development are distinct outcomes.

A viable mutated cell creates the biological possibility




Mutation is necessary for this pathway, but exposure does not make every altered cell malignant.

Radiation can injure DNA directly

- Ionization can disrupt atoms and molecular bonds within cellular material.
- DNA lesions include base damage, strand breaks, and DNA–protein crosslinks.
- The biological effect depends on the damage and the cellular response.
- An unrepaired lesion is not equivalent to a fixed mutation unless the cell survives and propagates it.

Water radiolysis generates reactive intermediates



Indirect radical-mediated injury is a major mechanism of radiation-induced DNA damage.

DNA lesions differ in their consequences

Lesion	Why it matters
Single-base damage	May be corrected by cellular DNA repair systems.
Single-strand break	Often repairable; contributes to the damage burden.
Double-strand break	Especially serious; joining errors can alter chromosomes.
DNA–protein crosslink	Interferes with normal DNA structure and processing.

Double-strand breaks are particularly consequential

- They disrupt both strands of the DNA double helix.
- Surviving cells require a mechanism to restore chromosomal continuity.
- Faulty repair may cause deletions, duplications, inversions, or translocations.
- The resulting chromosomal damage can persist when cell-cycle controls fail.

Two routes repair double-strand breaks


Homologous recombination

- One of two major mammalian double-strand break repair pathways.
- Uses homologous sequence information to guide repair.

Nonhomologous end joining

- as the more common DSB repair pathway.
- End joining can introduce sequence changes or rearrangements.

End joining may restore continuity with a genomic cost




Physical restoration of a chromosome does not guarantee preservation of its original sequence.

Chromosomal rearrangements may persist after repair

- Inversions change the orientation of a chromosomal segment.
- Translocations move material between chromosomal locations.
- Short deletions or duplications can modify genomic content.
- These outcomes among mutations associated with NHEJ.

Cell-cycle checkpoints provide a crucial safeguard



A damaged cell becomes a lasting hazard when altered genetic material survives checkpoint control.

Cancer risk extends beyond the exposure event

- Damage arises during irradiation; Malignant transformation need not be immediate.
- Persisting chromosomal abnormalities can contribute to carcinogenesis years later.
- Additional transforming events may follow the initial genetic lesion.
- Clinical latency therefore does not exclude an earlier radiation contribution.

Tissue destruction and carcinogenesis are not the same

Early tissue injury

- Cell loss follows DNA damage and impaired replication.
- Often conspicuous in rapidly proliferating normal tissue.

Late cancer risk

- Requires surviving cells with retained or acquired oncogenic changes.
- May emerge after a prolonged interval without obvious acute injury.


A plausible mechanism is not a patient-level diagnosis

- Radiation is a demonstrated human carcinogen in the settings discussed by Robbins.
- A specific tumor cannot automatically be assigned to a particular exposure.
- Interpretation requires exposure history, latency, organ involved, and population evidence.
- A cancer diagnosis alone does not reveal its etiologic origin.

Biological effects depend on exposure context

Modifier	Biologic relevance
Dose delivery	Intervals may allow cellular repair.
Field size	Changes the volume and organs exposed.
Cell proliferation	Actively dividing populations are vulnerable to DNA injury.
Oxygenation	Influences free-radical-mediated injury.
Vascular response	Can drive fibrosis and chronic ischemic tissue change.

Spacing radiation doses changes repair opportunity



Fractionation modifies biological damage; It does not erase all accumulated exposure effects.

Field size determines the exposed tissue population


Localized field

- Only a restricted anatomical region is deliberately irradiated.
- Tissues outside the field may be shielded.

Broad exposure

- More organs and proliferating cell populations receive energy.
- Consequences can differ even if a simple dose number looks similar.

Oxygen participates in radical-mediated injury



This describes a determinant of biological effect, not a patient-specific numerical cancer-risk estimate.

Fibrosis is a separate late radiation complication

Vascular pathway

- Endothelial damage may produce luminal narrowing or occlusion.
- Chronic ischemia impairs tissue function and healing.

Cancer pathway

- Surviving genetically altered cells can continue dividing.
- Neoplastic transformation may develop after years.

Absorbed dose and biologic dose answer different questions

Measure	Meaning
Gray (Gy)	Energy absorbed per unit tissue mass.
Sievert (Sv)	Dose measure weighted for biological effect.
Millisievert (mSv)	One-thousandth of a sievert; used for small exposure summaries.
Interpretation	A single dose figure is insufficient to specify individual cancer outcome.

An mSv value requires careful interpretation

- The number summarizes an exposure estimate rather than a cancer diagnosis.
- Type of radiation, anatomical distribution, and exposure context still matter.
- Radiological risk discussions should distinguish absorbed from weighted dose.
- Comparisons are most defensible when measures refer to the same dose quantity.

Radiation-associated cancer is a probability question

- The endpoint of concern is a change in future cancer incidence.
- An exposed group can show excess cancers without every individual becoming ill.
- An unexposed person may develop the same tumor type for unrelated reasons.
- Population evidence and individual attribution are different inferential tasks.

Higher-dose exposure has established carcinogenic consequences

- Exposures producing doses greater than 100 mSv as clearly concerning.
- Major human exposure settings show later leukemias and solid tumors.
- The existence of excess cancer is well supported; Precise risk varies with exposure circumstances.
- This statement is not a claim that doses below 100 mSv are safe.

The 5–100 mSv range creates a detection problem

- Expected increases in cancer incidence may be small relative to background incidence.
- Dose reconstruction may not be precise in every exposed individual.
- Substantial difficulty measuring the long-term risk in this interval.
- Uncertain estimation must not be interpreted as absence of biological effect.

Do not turn epidemiologic dose ranges into thresholds

- Levels of statistical evidence, not a biologically established safe cutoff.
- Different populations and designs can yield different statistical precision.
- A measured association is not identical to a universal minimum carcinogenic dose.
- Use the word “evidence” rather than “no risk below” when teaching low-dose effects.


Acute x-ray and gamma-ray evidence above 50 mSv

- Good evidence for a statistically significant excess risk at acute doses greater than 50 mSv.
- This statement concerns population-level cancer incidence.
- The radiation type and acute-exposure context are material qualifiers.
- It should not be generalized automatically to every organ or delivery pattern.

Very low acute doses: Evidence is less certain

- “reasonable evidence” at acute x-ray and gamma-ray doses greater than 5 mSv.
- Low-dose risk estimates require careful interpretation of uncertainty.
- The textbook does not give an individual probability of malignancy for a single low-dose examination.
- Do not invent a numerical probability from an exposure value alone.

Small excess risks require large study populations



That studies may require roughly 50,000–5,000,000 participants to quantify risk in the 5–100 mSv range.

Population associations are not causal labels for each tumor

- An exposed cohort supplies information about frequency, not a unique signature in an individual cancer.
- The same tumor type occurs through multiple carcinogenic pathways.
- Causation is assessed with exposure details and cumulative evidence.
- Use “radiation-associated” when attribution in a single patient is uncertain.

Radiologic exposure estimates are context dependent

- Reported dose examples are illustrative, not values for every machine or protocol.
- Anatomical target and exposure geometry differ between radiography and CT.
- A contemporary clinical estimate requires the particular examination and setting.
- Robbins values are retained here as textbook-era reference examples only.

Chest imaging: Examples reported by Robbins

Examination	Reported dose
Posteroanterior chest radiograph	0.01 mSv
Lateral chest radiograph	0.15 mSv
Chest CT	10 mSv

Values quoted from the textbook as lung-related effective-dose examples; Not universal current protocol estimates.

A dose estimate is not a cancer-risk calculation



Avoid translating “one CT” directly into a fabricated individual cancer percentage.

Evidence settings described

Exposure setting	Reported tumor signal
Atomic-bomb survivors	Leukemias and solid tumors in several organs.
Chernobyl nuclear accident	Increased thyroid cancers.
Marshall Islands fallout	Thyroid tumors and elevated leukemia frequency.
Therapeutic irradiation	Second hematologic and solid malignancies.
Radon exposure	Lung cancer risk.

Hiroshima and Nagasaki: A foundational observation

Atomic-bomb survivors

EXPOSURE

Major ionizing radiation exposure

OBSERVED FINDING

Increased leukemias and several solid tumors

INTERPRETATION

Supports human carcinogenicity across different tissue types.

Atomic-bomb survivors: Example tumor spectrum

Malignancy group	Example reported by Robbins
Hematologic	Leukemias
Endocrine	Thyroid tumors
Breast	Breast cancers
Respiratory	Lung cancers

Leukemia is one observed radiation-associated malignancy

- An increased incidence of leukemia is documented among atomic-bomb survivors.
- The Marshall Islands account also reports elevated leukemia frequency.
- Radiotherapy survivors may develop acute myeloid leukemia.
- These observations come from different exposure contexts and should not be pooled as one dose-response estimate.

Radiation-associated solid tumors involve multiple organs

- Thyroid, breast, and lung tumors among atomic-bomb survivor outcomes.
- Tumor distribution depends on which tissues were exposed and the exposure setting.
- An observed association does not imply every tissue has equal sensitivity.
- Organ-specific evidence is more informative than a universal “cancer risk” label.

Chernobyl: Thyroid cancer after a nuclear accident

Chernobyl accident

EXPOSURE

Population exposure following reactor accident

OBSERVED FINDING

Increased thyroid cancers among survivors

INTERPRETATION

A radiation-associated tumor pattern can be prominent in a particular organ.

Marshall Islands: Cancer risk after nuclear fallout

Marshall Islands inhabitants

EXPOSURE

Exposure to nuclear fallout

OBSERVED FINDING

Thyroid tumors and elevated leukemia frequency

INTERPRETATION

The textbook also mentions birth defects, a distinct non-cancer outcome.

Exposure route and anatomy influence observed outcomes

Exposure description

- External whole-body or regional irradiation differs from internal exposure.
- The responsible radionuclide or radiation type can matter.

Outcome interpretation

- Assess organs receiving the biologically relevant exposure.
- Keep tumor outcomes separate from other radiation injury.

Radiotherapy raises a long-term survivorship question

Cancer-treatment survivors

EXPOSURE

Radiation therapy for a prior malignancy

OBSERVED FINDING

Subsequent myeloid and solid malignancies

INTERPRETATION

The immediate therapeutic aim and later carcinogenic hazard must be considered separately.

Acute myeloid leukemia and MDS are reported late effects

- Acute myeloid leukemia among secondary cancers after radiotherapy.
- Myelodysplastic syndrome is also explicitly listed.
- These entities represent distinct marrow disorders following cancer treatment.
- A specific therapy-related diagnosis requires the individual clinical and pathological context.

Solid second cancers can follow regional irradiation

- Solid tumors as another delayed neoplastic consequence of radiotherapy.
- The exposure field provides anatomical context for evaluating later lesions.
- A new solid tumor is not automatically recurrence of the original malignancy.
- Long-term follow-up must be interpreted without assuming causation for every new tumor.

Therapeutic benefit and late risk coexist

Immediate treatment objective

- Deliver a tumoricidal radiation dose to the cancer.
- Exploit differences in damage recovery between tumor and normal tissue.

Delayed complication

- Normal surviving cells may retain radiation-associated genomic changes.
- Secondary malignancy risk enters long-term survivorship assessment.

Acute injury and secondary cancer require different timelines

Feature	Acute injury	Second malignancy
Principal process	Direct tissue damage and cell loss	Survival and propagation of altered cells
Clinical pattern	Early dysfunction in sensitive tissues	New cancer after a long interval
Biologic endpoint	Tissue depletion or inflammation	Clonal neoplastic proliferation

Pediatric exposure requires special attention

- Children a group with particularly high secondary-cancer concern.
- Exposure occurs while many tissue cell populations remain proliferative.
- Long intervals after childhood exposure permit late outcomes to become clinically apparent.
- The practical interpretation still depends on actual exposure and clinical need.

Repeated pediatric CT: Small measured associations

Pediatric CT epidemiology

EXPOSURE

Children receiving at least two CT examinations

OBSERVED FINDING

Very small but measurable increases in leukemia and malignant brain tumors

INTERPRETATION

A measurable population association does not imply a large absolute risk for an individual child.

Relative and absolute increases answer different questions

Relative change

- Compares incidence between groups.
- Can appear important even when the baseline event is uncommon.

Absolute change

- Counts added events over a defined population and period.
- Is necessary for balanced communication of a “very small” increase.

Adolescent chest irradiation: Later breast cancer

Historical radiotherapy cohorts

EXPOSURE

Chest irradiation during adolescence

OBSERVED FINDING

Increased subsequent breast cancer in females exposed at that age

INTERPRETATION

Age at the time of exposure can matter for the later tumor spectrum.

Timing of exposure changes the reasoning problem


Childhood / adolescence

- Different developmental and proliferative tissue context.
- Potential for extended follow-up after exposure.

Later-life exposure


- Different tissue state and competing clinical risks.
- Do not assume identical risk at the same reported dose.

Radon originates in the natural decay of uranium



Polonium-214 and polonium-218 as carcinogenic radon progeny.

Radon progeny cause localized alpha exposure



The carcinogenic exposure arises largely from alpha-emitting daughters rather than a generic “radon dose.”

Lung localization explains the primary tumor outcome



The causal chain links environmental exposure to the tissue in which cancer is observed.

Uranium miners: Occupational lung-cancer risk

Uranium-mining exposure

EXPOSURE

Chronic inhalation of radon decay products

OBSERVED FINDING

Elevated risk of lung carcinoma

INTERPRETATION

A prolonged workplace exposure can produce a recognizable organ-specific malignancy.

Residential radon can also present a carcinogenic hazard

- Cancer risk in homes with very high radon levels.
- Some household concentrations can approach levels found in mines.
- Lower residential concentrations are described as suspected contributors to lung cancer.
- A household address does not substitute for an actual exposure assessment.

Radon and tobacco require combined risk thinking

Radon

- Alpha-emitting daughter products expose pulmonary tissues.
- Lung carcinoma is the reported associated malignancy.


Tobacco co-exposure

- Particular concern among people who smoke.
- The excerpt does not quantify interaction or supply a joint risk ratio.

Textbook mortality estimates are time-bound

- Nearly 3,000 annual U.S. Nonsmoker deaths to radon, citing a CDC estimate.
- This is a reported textbook figure, not independently verified current surveillance.
- It highlights public-health importance without resolving individual cancer attribution.
- Do not present the figure as an up-to-date count without checking its original source and year.

Risk interpretation requires more than knowing a hazard



The same carcinogenic agent can be associated with different risk profiles across exposure settings.

A radiation-risk study needs interpretable inputs

Question	Why it affects inference
Which population?	Baseline disease frequency and age may differ.
What exposure?	Radiation type, intensity, dose and distribution matter.
How measured?	Dose misclassification can obscure small effects.
What outcome?	Leukemia and organ-specific solid tumors are distinct endpoints.
What follow-up?	Malignancies may appear long after exposure.

Exposure settings: Compare before generalizing

Setting	Exposure structure	Cancer signal
Bombings	Acute population exposure	Leukemias and several solid cancers
Fallout	Environmental contamination	Prominent thyroid and hematologic outcomes
Radiotherapy	Targeted therapeutic fields	Late hematologic and solid second cancers
CT imaging	Medical diagnostic exposures	Small epidemiologic signals in childhood
Radon	Chronic inhalational exposure	Lung carcinoma

Low-dose interpretation should preserve uncertainty

- Lower-dose cancer risk as difficult to measure rather than absent.
- Population size, exposure reconstruction, and tumour background rates limit statistical certainty.
- Numeric estimates for a particular patient cannot be extracted from the textbook's dose examples.
- Clinically necessary irradiation and avoidable exposure should not be discussed as identical decisions.

Case reasoning: Cancer years after Hodgkin radiotherapy

A long-term survivor of Hodgkin lymphoma develops a new myeloid malignancy. Which exposure history is relevant?

Reasoning checkpoints

- Prior therapeutic irradiation is relevant to secondary malignancy assessment.
- AML and MDS are listed by Robbins among post-radiotherapy neoplastic outcomes.
- The diagnosis still requires clinical and pathological evaluation.

Case reasoning: Repeated childhood CT examinations

A child underwent multiple CT examinations. How should possible later cancer risk be explained?

Reasoning checkpoints

- A very small measurable association for leukemia and malignant brain tumors after at least two CT scans.
- Avoid translating scan count into a fabricated individual probability.
- Explain the evidence while preserving the medical context of the examinations.


Case reasoning: Residential radon and smoking

A person who smokes is concerned about radon exposure at home. What is the relevant malignant endpoint?

Reasoning checkpoints

- Radon daughter products can deposit in lung and emit alpha particles.
- Lung carcinoma is the important radiation-related cancer in the excerpt.
- Smoking co-exposure heightens concern, but no interaction ratio is provided.

A common causal framework links all exposure settings



Epidemiologic evidence is required to establish how often the late outcome occurs.

Five conclusions for clinical interpretation

- Ionizing radiation is a proven human carcinogen.
- Unrepaired or aberrantly repaired DNA damage creates a plausible transformation pathway.
- Strong evidence comes from bombings, nuclear fallout, radiotherapy, and radon.
- Children and irradiated survivors warrant special attention in long-term assessment.
- Low-dose uncertainty is not proof of zero risk, and dose alone is not an individual prognosis.

Principal source and scope

- Robbins, Cotran & Kumar: Pathologic Basis of Disease. 11th ed. (2026).
- Primary teaching section: Cancer Risks From Exposures to Radiation.
- Mechanistic context: DNA Damage and Carcinogenesis; Main Determinants of Biologic Effects of Ionizing Radiation; Radiation Units.
- No independent update to imaging-dose estimates or historical surveillance figures is claimed.