

Chronic pancreatitis

Irreversible injury · fibrosis · loss of pancreatic function

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning outcomes

- Explain how recurrent pancreatic injury produces irreversible fibrotic remodeling.
- Recognize major causes, molecular susceptibility, and defining morphology.
- Differentiate autoimmune pancreatitis phenotypes and carcinoma mimics.
- Link exocrine and endocrine injury to clinical findings and diagnostic clues.

Normal pancreas: Two functional compartments

EXOCRINE

Acinar cells synthesize digestive enzymes.

Duct epithelium conveys fluid and bicarbonate.

ENDOCRINE

Islets of Langerhans regulate blood glucose.

Endocrine loss generally occurs later in chronic disease.

This functional division predicts the sequence of clinical failure.

The defining lesion is irreversible parenchymal destruction

Persistent inflammation → exocrine tissue loss + fibrosis


Structural / mechanistic meaning

Acinar dropout and ductal remodeling are central structural changes.

Clinical / interpretive meaning

With progression, loss of islet cells produces endocrine insufficiency.

Recurrent injury drives a self-perpetuating process



The progression model links recurrent acute episodes to irreversible chronic damage.

Acute and chronic pancreatitis are not synonymous

Feature	Acute pancreatitis	Chronic pancreatitis
Principal injury	Potentially reversible acinar injury	Irreversible tissue loss
Dominant process	Enzyme activation and inflammation	Fibrogenic remodeling
Typical outcome	Resolution in many patients	Fibrosis and insufficiency
Microscopic clue	Edema ± necrosis	Acinar atrophy with fibrosis

Etiologic groups form a coherent framework

Alcohol exposure

Long-term heavy alcohol use is the commonest cause cited.

Duct obstruction

Calculi or neoplasms block pancreatic outflow.

Immune-mediated disease

Autoimmune pancreatitis has distinct morphologic variants.


Inherited predisposition

Genetic susceptibility may coexist with acquired injury.

Alcohol is the leading recognized association


- Sustained alcohol exposure underlies many cases of chronic pancreatitis.
- Ductal dilation, protein plugs, and calcifications are especially characteristic.
- Exposure alone is insufficient to explain every case: Susceptibility varies.
- Alcohol can also precipitate symptomatic pain attacks.

Alcohol-related disease remodels the duct system



Schematic links documented duct features; Mechanisms are multifactorial.

Obstructed ducts can sustain parenchymal injury



Evaluate an obstructive process rather than assuming alcohol as the cause.

Autoimmune pancreatitis is a separate pathogenic category

Type 1

IgG4-associated, part of systemic IgG4-related disease.

Type 2

Duct-centered neutrophilic lesions; May coexist with ulcerative colitis.

Type 3

Checkpoint-inhibitor–associated pancreatic injury.

Common pitfall

A focal or mass-like process can resemble pancreatic carcinoma.


Inherited susceptibility changes the threshold for injury

- Some recurrent acute attacks beginning in childhood lead to chronic fibrosis.
- Inherited mechanisms often increase or prolong intrapancreatic trypsin activity.
- PRSS1, SPINK1, and CFTR are emphasized in the adjacent Robbins section.
- Genetic and environmental factors can act together.

Pancreatitis-associated genes affect distinct safeguards

Gene	Alteration	Pathogenic effect
PRSS1	Gain of function	Persistent/greater trypsin activation
SPINK1	Loss of function	Weakened trypsin inhibition
CFTR	Loss of function	Viscous secretions and duct plugging
CTRC	Loss of function	Reduced trypsin degradation

PRSS1 gain of function sustains trypsin activity



Autosomal-dominant hereditary pancreatitis with PRSS1 variants.

SPINK1 deficiency weakens a local protective brake

Normal safeguard

SPINK1 encodes a trypsin inhibitor.


Adequate inhibitor limits inappropriate intrapancreatic activity.

Loss of function

Reduced inhibitor function favors sustained trypsin activity.

Recessive inheritance for this form.

CFTR dysfunction favors duct obstruction




CFTR-related pancreatitis may occur without classic cystic fibrosis findings.

Other susceptibility genes emphasize calcium and enzymes


Gene	Protein function	Mechanistic implication
CASR	Calcium-sensing receptor	Calcium changes may facilitate trypsin activation
CTRC	Chymotrypsin C	Impaired degradation of trypsin
CPA1	Carboxypeptidase A1	Toxic gain of function in a zymogen-regulating enzyme

Susceptibility and exposure are not mutually exclusive



A single patient may have more than one contributor; Avoid rigid etiologic labels.

The bridge from acute to chronic disease is fibrosis




A conceptual sequence, not a claim that every acute episode becomes chronic.

Acinar injury releases inflammatory mediators

- Repeated injury stimulates local inflammatory signaling.
- In chronic disease, profibrotic effects become especially important.
- Fibrogenic signals favor activated stellate cells and matrix deposition.
- Progressive replacement by scar compromises exocrine tissue.

Pancreatic stellate cells are central effectors



Stellate cells behave as periacinar myofibroblasts in the injured gland.

TGF- β and PDGF promote fibrotic remodeling

TGF- β

Profibrotic signaling is prominent in chronic pancreatitis.

Supports extracellular-matrix accumulation.


PDGF

Supports stellate-cell activation and proliferation.


Promotes persistence of the fibroblastic response.

Extracellular matrix replaces functioning acini


Editable architectural schematic



Intact



Remodeling



Scarred

Schematic — not a histologic specimen

Structural features

Intact lobules contain enzyme-producing acini.

Injury induces collagen deposition around lobules.

Extensive fibrosis leaves scattered residual glands.

Duct distortion reinforces injury



Ductal pathology and parenchymal scarring can reinforce one another.

Exocrine and endocrine injury follow different trajectories

Earlier / prominent

Acinar atrophy and dropout

Reduced digestive enzyme reserve


Later / progressive

Relative early preservation of islets

Diabetes when endocrine tissue is lost

Gross morphology: A hard, scarred gland

Editable architectural schematic



Duct with concretions

Schematic — not a histologic specimen

Structural features

Firmness reflects replacement of parenchyma by fibrosis.

Ducts may be visibly dilated.

Calcified intraductal concretions may be present.

Calcification points to chronic ductal disease

Ductal concretions are a structural clue, not the sole diagnostic criterion.

Structural / mechanistic meaning


They are particularly emphasized in alcohol-associated chronic pancreatitis.

Clinical / interpretive meaning

CT and ultrasonography can demonstrate intrapancreatic calcifications.

Histology: Fibrosis and acinar loss coexist

Editable architectural schematic



Schematic — not a histologic specimen

Structural features


Broad fibrous areas partition residual parenchyma.

Acinar cell atrophy and dropout are defining.

Chronic inflammatory cells accompany the fibrotic process.

Lobules and ducts attract chronic inflammation

Editable architectural schematic



Schematic — not a histologic specimen

Structural features


Inflammatory infiltrates surround lobules and ducts.

Inflammation accompanies rather than replaces fibrotic change.

Interpret architecture together with cell populations.

Relative islet preservation creates an apparent paradox

Editable architectural schematic



Schematic — not a histologic specimen

Structural features

Exocrine acini may be extensively lost.

Residual islets persist in the scarred stroma.


Islet aggregates can appear fused or enlarged.

Pancreatic duct epithelium may undergo adaptation

Ductal change	Morphologic meaning
Dilation	Altered outflow and remodeling
Atrophy	Loss of epithelial volume
Hyperplasia	Reactive epithelial proliferation
Squamous metaplasia	Phenotypic adaptation under chronic injury

Alcohol-associated ducts contain protein plugs

Editable architectural schematic



Schematic — not a histologic specimen

Structural features

Proteinaceous material accumulates within duct lumina.

Concretions may calcify.

Correlate ductal change with acinar atrophy and fibrosis.

Morphologic recognition uses a constellation


Compartment	Expected observation	Interpretive relevance
Stroma	Collagen-rich fibrosis	Irreversible remodeling
Exocrine acini	Atrophy/dropout	Loss of secretion capacity
Ducts	Variable dilation $\pm$ plugs	Obstructive/secretory effects
Islets	Relative preservation early	Delayed endocrine failure

Type 1 autoimmune pancreatitis: IgG4-related disease

- Type 1 AIP is a pancreatic manifestation of IgG4-related disease.
- Dense lymphoplasmacytic inflammation accompanies fibrosis.
- Abundant IgG4-positive plasma cells support the tissue pattern.
- The full histologic constellation matters more than one isolated feature.

Storiform fibrosis is a distinctive tissue pattern

Editable architectural schematic



Schematic — not a histologic specimen

Structural features


Fibrous bands form a whorled or storiform configuration.

Lymphoplasmacytic inflammation is prominent.

The pattern suggests type 1 autoimmune pancreatitis in context.

Obliterative phlebitis is a second defining clue

Editable architectural schematic



Nearly obliterated vein

Schematic — not a histologic specimen

Structural features

Inflammation and fibrosis involve venous walls.

The venous lumen may be obliterated.

Search for this in a mass-forming fibrosing pancreatic lesion.

IgG4-positive plasma cells require pattern correlation

What supports type 1 AIP

Storiform fibrosis

Obliterative phlebitis

Rich lymphoplasmacytic infiltrate

Interpretive caution

IgG4 immunostaining provides supportive tissue information.


Do not reduce diagnosis to plasma-cell presence alone.

Type 2 autoimmune pancreatitis centers on ducts

- Type 2 AIP shows neutrophils invading pancreatic duct epithelium.
- Neutrophils may also occupy lumina of medium-sized ducts.
- These are granulocytic epithelial lesions.
- Abundant IgG4-positive plasma cells typical of type 1 are absent.

Granulocytic epithelial lesions define type 2 histology

Editable architectural schematic



Schematic — not a histologic specimen

Structural features

Neutrophils infiltrate ductal epithelium.

Neutrophils can accumulate in the duct lumen.

The injury has a conspicuously duct-centered distribution.

Autoimmune pancreatitis: Compare types 1 and 2

Attribute	Type 1 AIP	Type 2 AIP
Systemic context	IgG4-related disease	Mainly pancreatic
Inflammatory feature	IgG4-rich plasma cells	Duct-associated neutrophils
Characteristic lesion	Storiform fibrosis / phlebitis	Granulocytic epithelial lesion
Clinical association	Systemic IgG4 disease	Sometimes ulcerative colitis

Type 3 is described as treatment-related

- Robbins 11th ed. Uses “type 3 AIP” for checkpoint-inhibitor–associated injury.
- Described as uncommon and often mild, particularly with combination therapy.
- Diagnosis is commonly based on imaging and enzyme abnormalities.
- Histologic features remain incompletely established in the supplied source.

A pancreatic head mass may be inflammatory

Autoimmune pancreatitis

Can produce a mass-like enlargement.

Often responds to glucocorticoid treatment.

Pancreatic carcinoma

Can also appear as a pancreatic head mass.

Distinguishing neoplasia before therapy is essential.

Avoid treating every mass as autoimmune disease

Mass-forming pancreatitis and pancreatic carcinoma are consequential mimics.

Structural / mechanistic meaning

Histology, distribution, serology and imaging require integrated interpretation.


Clinical / interpretive meaning

Steroid responsiveness does not replace an appropriate malignancy assessment.

Clinical symptoms may be episodic or persistent

- Many patients have recurrent mild-to-moderately severe abdominal pain.
- Others have persistent pain with radiation to the back.
- Attacks may follow alcohol consumption or excessive food intake.
- Opiates and drugs that increase sphincter-of-Oddi tone may precipitate attacks.

Recurrent pain reflects persistent pancreatic injury



Pain severity varies; There is no obligatory single symptom trajectory.

Some patients present only after functional loss

Symptomatic pathway

Painful recurrent episodes


Investigation reveals a chronically injured gland

Silent pathway

Little or no preceding pain

Later malabsorption or diabetes reveals disease

Exocrine failure disrupts intestinal digestion




Exocrine damage provides a mechanistic explanation for nutritional decline.

Weight loss is a consequence of malabsorption


- Exocrine deficiency impairs nutrient digestion.
- Malabsorption contributes to loss of body mass.
- The clinical picture may be dominated by nutritional rather than pain symptoms.
- Weight loss should be correlated with pancreatic structure and function.

Hypoalbuminemia can lead to edema



Edema is a secondary systemic consequence, not direct pancreatic swelling.

Endocrine tissue loss produces diabetes



Relative islet sparing on early histology does not rule out late diabetes.

Serum amylase has limited sensitivity in chronic disease

During an attack

Mild-to-moderate amylase elevation may occur.

Mild fever may accompany active inflammation.

Advanced disease


Extensive acinar loss may blunt enzyme release.

A normal value does not exclude chronic pancreatitis.

Imaging provides structural evidence

- Computed tomography can visualize pancreatic calcifications.
- Ultrasonography can also demonstrate calcified intrapancreatic concretions.
- Correlate imaging with symptoms, functional loss, and likely etiologic group.
- Ductal distortion and pancreatic fibrosis help explain chronicity.

Biliary obstruction changes the presentation



Alkaline phosphatase elevation and jaundice with obstruction.

Diagnosis requires convergent evidence

Clinical context

Recurrent upper abdominal/back pain or silent progressive insufficiency.

Etiologic clues

Alcohol, obstruction, hereditary predisposition, or autoimmunity.

Structure

Scarring, duct dilation, concretions or mass-like inflammation.

Function/laboratory

Malabsorption, diabetes; Amylase may be normal.

Pseudocysts are a recognized complication

- Pseudocysts may follow pancreatitis, particularly attacks on chronic alcoholic disease.
- They arise when pancreatic/peripancreatic necrotic injury is walled off.
- Pseudocysts in about 10% of chronic pancreatitis cases.
- Larger lesions may compress or erode adjacent structures.

Pseudocyst walls lack an epithelial lining

Pseudocyst

Fibrous and granulation-tissue wall

No true epithelial lining

True cyst (conceptual contrast)

An epithelial lining is present.

Cystic lesions require etiologic distinction.

Long-term morbidity extends beyond abdominal pain

Exocrine dysfunction

Chronic malabsorption and nutritional consequences.

Endocrine dysfunction

Diabetes mellitus in late-stage damage.

Pain

Persistent or recurrent severe pain.

Local sequelae

Pancreatic pseudocyst formation.

Interpret historical survival figures cautiously

Approximately 50% mortality over 20–25 years.

Structural / mechanistic meaning

The figure describes long-term prognosis in its textbook context; No cohort or era is specified in the excerpt.

Clinical / interpretive meaning

Do not interpret this figure as an individualized prognosis or as a contemporary risk estimate.

Case: Painful calcific pancreatic disease

Clinical vignette

A middle-aged patient with prolonged heavy alcohol exposure reports recurrent epigastric/back pain. CT demonstrates intrapancreatic calcifications.

Reason through the findings

- Which morphologic features are predicted?
- Why might serum amylase be nondiagnostic?

Expected findings: Fibrotic acinar loss, distorted/dilated ducts and concretions; Extensive acinar dropout may blunt enzyme release.

Case: Mass-forming autoimmune pancreatitis

Clinical vignette

Imaging shows a focal pancreatic head enlargement. Tissue examination demonstrates dense lymphoplasmacytic inflammation and storiform fibrosis.

Reason through the findings

- Which autoimmune subtype is most consistent?
- What additional tissue feature should be sought?

Type 1 AIP is favored; Assess obliterative phlebitis and IgG4-positive plasma cells while excluding carcinoma.

Case: Recurrent pancreatitis beginning in childhood

Clinical vignette


A patient experienced repeated episodes of acute pancreatitis beginning in childhood and now has pancreatic fibrosis.

Reason through the findings

- Which pathogenic axis suggests hereditary disease?
- Which three genes deserve initial conceptual emphasis?

Failure of normal trypsin regulation or duct secretion; PRSS1, SPINK1 and CFTR.

Morphology, mechanism and symptoms reinforce each other



A unified causal chain explains most important clinicopathologic observations.

Five findings to retain

- Chronic pancreatitis means irreversible acinar loss and fibrosis.
- Alcohol, obstruction, genetic predisposition and autoimmune injury are key causes.
- Stellate-cell activation under TGF- β /PDGF signaling produces scar.
- Type 1 AIP has storiform fibrosis and phlebitis; Type 2 has ductal neutrophils.
- Chronic pain, malabsorption, diabetes and pseudocysts reflect structural injury.

Reference and source boundaries

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. Elsevier; 2026.
- Textbook discussion: chronic pancreatitis — pathogenesis, morphology, and clinical features.
- Also covered: inherited predisposition to pancreatitis and major implicated genes.
- Also covered: postpancreatitis pseudocysts and their lack of an epithelial lining.
- All glandular and microscopic illustrations are original editable schematics, not diagnostic tissue images.