

Congenital Iodine Deficiency and Myxedema

A clinicopathologic study of hypothyroidism across the lifespan

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Explain the dependence of fetal neurodevelopment on thyroid hormone availability.
- Distinguish nutritional iodine deficiency, developmental defects, and dysharmonogenesis.
- Link the childhood phenotype and adult myxedema findings to hormone deficiency.
- Interpret TSH and free T4 patterns in primary, central, overt, and subclinical disease.

One hormone deficit, two age-dependent phenotypes

Early-life onset

- Fetal/neonatal nervous-system development is highly vulnerable.
- Growth failure and intellectual impairment may become prominent.
- Coarse facial features, tongue protrusion, and umbilical hernia may occur.

Older child / adult onset


- Physical and mental activity progressively slow.
- Cold intolerance, constipation, and reduced cardiac output emerge.
- Extracellular matrix accumulation produces nonpitting edema.

Timing of hormone deficiency determines which tissues and functions are most affected.

Hypothyroidism: Functional definition

- A structural or functional abnormality interferes with thyroid hormone production.
- The defect may reside in thyroid tissue or upstream regulatory structures.
- Clinical manifestations depend on severity, duration, and developmental age.
- Primary disease is far more common than central disease.


Hypothalamic–pituitary–thyroid axis



Pathophysiologic consequence

Thyroid hormone restrains hypothalamic TRH and pituitary TSH secretion by negative feedback.


How TSH activates thyroid follicles



Diagnostic interpretation

The pituitary signal acts on follicular epithelial cells; Failure of thyroid response can produce primary hypothyroidism.

The follicle is the structural unit of thyroid hormone production




Schematic: Native editable shapes; Not a histologic specimen

Thyroid hormones circulate mainly bound to plasma proteins

- Thyroid follicular cells release predominantly thyroxine (T4) and smaller amounts of triiodothyronine (T3).
- Thyroxine-binding globulin, transthyretin, and albumin reversibly bind circulating hormone.
- The free hormone fraction is available to tissues and supports physiologic signaling.

Peripheral activation of thyroid hormone




Pathophysiologic consequence

T3 binds the nuclear receptor more avidly than T4; Downstream effects involve regulation of transcription.

Why thyroid hormone deficiency affects many organs

- Thyroid hormone regulates carbohydrate and lipid catabolism and protein synthesis.
- Its net physiologic effect includes increasing basal metabolic rate.
- Fetal and neonatal brain development critically depends on adequate availability.
- Adult deficiency affects cognition, thermoregulation, skin, heart, bowel function, and lipids.

Negative feedback explains the diagnostic TSH response



Diagnostic interpretation

An elevated TSH indicates an intact pituitary responding to inadequate thyroid output; Upstream disease behaves differently.

Hypothyroidism is classified by the level of dysfunction

Pattern	Primary	Central
Anatomic site	Thyroid gland	Pituitary or hypothalamus
Hormone deficit	Reduced thyroid output	Inadequate TSH or TRH drive
Typical TSH	Increased	Low or inappropriately normal
Role in this lecture	Common background for congenital and adult cases	Important interpretation trap

Two distinct etiologic pathways in congenital hypothyroidism

Insufficient substrate

- Dietary iodine deficiency reduces the substrate available for hormone synthesis.
- Historically endemic in iodine-poor regions.
- Food iodization has greatly reduced this preventable burden.

Thyroid developmental / synthetic defect

- Thyroid agenesis or hypoplasia reduces functioning tissue.
- Dyshormonogenesis interrupts a step in hormone biosynthesis.
- Germline developmental abnormalities can involve PAX8, FOXE1, or the TSH receptor.


Congenital iodine deficiency: Textbook definition

- Hypothyroidism appearing in infancy or early childhood in the context of deficient iodine availability.
- Historically described using the term “cretinism,” now considered outdated terminology.
- The central consequence is thyroid hormone insufficiency during a vulnerable developmental period.
- Rare inherited hormone-synthesis defects may produce a similar early-life syndrome.

Endemic iodine deficiency is a nutritional setting

- Mountainous and inland regions historically affected by low dietary iodine.
- Illustrative regions named in the text include the Himalayas, inland China, and parts of Africa.
- The location is a historical epidemiologic context, not a diagnostic criterion.
- Thyroid hormone availability, rather than residence alone, is the causal biological concern.

How low iodine intake impairs hormone synthesis




Pathophysiologic consequence

Widespread iodine supplementation of foods is associated with markedly lower prevalence of the historical endemic syndrome.

Why iodine supplementation changed the epidemiology

- Endemic iodine deficiency was historically a frequent cause of early-life hypothyroidism.
- Iodine supplementation of foods addresses the nutritional bottleneck before severe deficiency develops.
- A major reduction in the prevalence of congenital iodine-deficiency syndromes.
- Inherited thyroid defects remain possible when dietary iodine is adequate.

Hormone biosynthesis has several vulnerable steps



Pathophysiologic consequence

Inherited failure of a single step can cause dyshormonogenetic hypothyroidism; Separate mechanisms need different etiologic interpretations.

Dyshormonogenesis: What can fail?

Step	Normal process	Consequence of a defect
Iodide transport	Iodide enters follicular epithelial cells	Insufficient substrate inside the cell
Organification	Iodine binds tyrosine residues in thyroglobulin	Iodinated hormone precursors cannot form normally
Further processing	Iodinated precursors yield active T3/T4	Reduced hormonally active product

Thyroid developmental defects differ from synthetic defects

Agenesis / hypoplasia

- Thyroid parenchyma is absent or severely reduced.
- Fewer functioning follicles are available for hormone production.
- Some cases to thyroid developmental gene defects.

Dyshormonogenesis

- Thyroid tissue may be present but synthetic activity is impaired.
- The block can involve transport, organification, or hormone formation.
- A goitrous phenotype may accompany some biosynthetic disorders.

Examples of congenital thyroid developmental genes

- PAX8, FOXE1, and TSH receptor mutations among rare causes.
- These genetic causes should not be confused with nutritional iodine insufficiency.
- Different defects converge on low thyroid hormone availability.
- A molecular diagnosis cannot be inferred from the clinical phenotype alone.

Why a goiter may be present—or absent

Goitrous hypothyroidism

- Persistent pituitary stimulation may enlarge functioning thyroid epithelium.
- The source table associates iodine deficiency and certain biosynthetic defects with goiter.
- Goiter reflects tissue response, not proof of preserved hormone synthesis.


Small or absent thyroid

- Congenital agenesis implies absent parenchyma.
- Hypoplasia implies reduced parenchyma.
- Normal thyroid enlargement cannot be assumed in developmental absence.

Maternal thyroid hormone is developmentally important

- Fetal brain development depends on adequate maternal thyroid hormone supply.
- Deficiency early in gestation may precede functional maturation of the fetal thyroid.
- The timing of insufficiency is strongly related to severity of neurologic consequences.
- Congenital hypothyroidism therefore cannot be understood only as a postnatal disorder.

Fetal timing changes the consequences of hormone deficiency



Schematic developmental phases — not a gestational-age scale

Early versus later gestational thyroid deficiency

Before fetal thyroid function

- Maternal deficiency removes a crucial early hormone supply.
- This timing with particularly severe intellectual impairment.
- Critical developmental processes may be compromised.

After fetal thyroid function develops

- Fetal hormone synthesis provides an additional source.
- Less adverse neurodevelopmental effect from later isolated maternal deficiency.
- The central distinction is developmental timing, not a single fixed gestational cutoff.

Age of onset is central to neuropathologic consequences

- The fetal and neonatal brain is a critical thyroid hormone target.
- Early hormone insufficiency can alter later intellectual development.
- Neurodevelopmental consequences are not interchangeable with adult cognitive slowing.
- The main disease variable is developmental timing of deficient hormone exposure.

Intellectual impairment is a major early-life consequence

- The classic syndrome includes severe intellectual disability.
- The text links greater severity to thyroid deficiency earlier in fetal life.
- This clinical effect reflects developmental vulnerability rather than only reversible adult-style fatigue.
- The clinical phenotype alone cannot establish the timing or cause of fetal hormone deficiency.

Stature and somatic development may be impaired

- Short stature is part of the classic congenital presentation.
- Growth and neurodevelopmental manifestations may occur together.
- Clinical examination should connect stature to additional features rather than use it alone.
- Growth findings must be interpreted alongside neurodevelopmental and thyroid abnormalities.

Craniofacial and soft-tissue manifestations

Head and face

- Coarsening of facial features is part of the historical presentation.
- Appearance alone is not disease-specific.
- Neurodevelopmental and growth history provide important context.

Tongue and abdominal wall

- Protruding tongue is listed as a characteristic feature.
- Umbilical hernia is also described.
- A constellation of signs is more informative than a single visual feature.

The childhood phenotype is a clinical constellation

- Severe intellectual impairment.
- Short stature with coarse facial features.
- Protruding tongue and umbilical hernia.
- A maternal/early-life history consistent with thyroid hormone insufficiency.

Pediatric case: Connect development and thyroid function


Clinical vignette

A young child from a historically iodine-poor setting has poor linear growth, substantial developmental delay, a protruding tongue, and an umbilical hernia.

Reasoning prompt

What unifying hormonal deficiency explains this pattern, and why does timing matter?

Pediatric case: The clinicopathologic explanation



Developmental implication

The source supports nutritional iodine deficiency as an important historical cause; The phenotype alone cannot exclude inherited thyroid defects.


Terminology: Use age- and mechanism-specific language

- “Cretinism” is an obsolete historical label recorded in the source, not preferred contemporary clinical wording.
- Congenital hypothyroidism describes deficiency from birth, while iodine deficiency specifies one cause.
- Dyshormonogenesis describes a biosynthesis defect; Thyroid agenesis describes missing gland tissue.
- Distinguishing the diagnosis from the cause avoids an etiologic assumption.

Myxedema: From historical label to severe hypothyroidism

- The term originally referred to hypothyroidism of older children and adults.
- Its historical description by William Gull in 1873.
- Today the term is often associated with severe hypothyroidism.
- Myxedema coma is a severe clinical endpoint, not a necessary feature of every patient with hypothyroidism.

Adult disease may evolve over years



Pathophysiologic consequence

The text emphasizes an insidious presentation that can delay clinical suspicion.

Early symptoms are often nonspecific

- Generalized fatigue may be the first reported complaint.
- Apathy and mental sluggishness may accompany tiredness.
- This triad can resemble depressive illness.
- Overt clinical signs become more recognizable with increasing severity.


Mental and speech slowing reflect systemic hypothyroidism

- Speech may become slow.
- Intellectual functioning can be slowed in severe disease.
- Patients may appear listless or apathetic.
- This is distinct from the enduring neurodevelopmental impairment of severe early-life deficiency.

Cold intolerance and increased body weight

- Cold intolerance is frequently noted in more advanced adult disease.
- Patients may become overweight.
- Decreased thyroid hormone action reduces the metabolic drive that normally raises basal metabolic rate.
- Neither weight gain nor cold intolerance alone confirms hypothyroidism.

Reduced sympathetic effects and autonomic symptoms




Pathophysiologic consequence

Constipation and diminished sweating to decreased sympathetic activity in hypothyroidism.

Skin manifestations: Cool and pale

- Cool skin is a reported physical sign.
- Pallor is linked in the source to reduced cutaneous blood flow.
- Decreased sweating may coexist.
- Skin texture and tissue swelling also reflect extracellular matrix changes.

Cardiac output falls in hypothyroidism




Cardiometabolic implication

The source describes the transcriptional explanation as a potential contributor rather than the only cardiac mechanism.

Cardiovascular consequences are functional and clinical

- Diminished cardiac output contributes to exertional limitation.
- Shortness of breath and reduced exercise capacity may develop.
- The absence of prominent cardiac symptoms does not exclude milder hypothyroidism.
- Clinical interpretation should integrate the cardiovascular findings with thyroid studies.

Thyroid hormone and myocardial gene expression




Cardiometabolic implication

This is a qualitative molecular-to-organ model directly based on the listed gene-expression example.

Lipid abnormalities are part of the endocrine phenotype

- Adult hypothyroidism can increase total cholesterol.
- Low-density lipoprotein (LDL) levels may also rise.
- The resulting lipid profile is proatherogenic.
- Robbins relates these abnormalities to increased cardiovascular mortality.

How thyroid insufficiency links to vascular risk




Cardiometabolic implication

An association with higher cardiovascular mortality, not a patient-specific risk estimate.

The substrate of myxedematous swelling is extracellular

- Accumulation of matrix substances in skin and soft tissue.
- The principal examples are glycosaminoglycans and hyaluronic acid.
- Deposition may affect skin, subcutaneous tissue, and visceral sites.
- This provides a morphologic basis for nonpitting edema.

Myxedema is an extracellular matrix phenomenon



Extracellular glycosaminoglycan / hyaluronic-acid accumulation → nonpitting swelling

Schematic of matrix-rich connective tissue; Not microscopy

Why edema in myxedema is described as nonpitting

- The altered interstitium contains increased matrix material.
- Clinical swelling is therefore described as nonpitting rather than simple dependent fluid accumulation.
- Histologic matrix accumulation and the bedside examination sign describe related features.
- Biochemical evidence remains important because nonpitting edema is not a stand-alone diagnosis.

Myxedematous tissue changes broaden facial features

- Extracellular matrix accumulation contributes to a broad, coarse facial appearance.
- Soft-tissue expansion is not limited to the face.
- This morphology is linked to tissue material rather than a thyroid tumor.
- Connect the appearance to the overall clinical and laboratory pattern.

Tongue enlargement and voice change reflect tissue infiltration

- Enlargement of the tongue is a classic adult manifestation.
- Deepening of the voice is described in the same pathologic setting.
- Underlying matrix accumulation provides the histopathologic correlate.
- These features should be distinguished from the protruding tongue described in early-life disease.

Myxedema coma is a severe endpoint, not a synonym

- The textbook uses myxedema especially in severe adult hypothyroidism.
- Its most extreme clinical expression can be myxedema coma.
- The term denotes a much more severe clinical state than uncomplicated nonpitting edema.
- A diagnosis of myxedema coma should not be inferred from edema alone.


Childhood onset versus adult-onset hypothyroidism

Domain	Early-life deficiency	Adult myxedema
Critical biology	Brain development and somatic growth	Established-organ physiology and metabolic function
Neurologic	Severe intellectual impairment in classic syndrome	Mental and speech slowing, apathy
Somatic	Short stature, coarse features, protruding tongue	Cold skin, nonpitting edema, coarse features
Systemic	Developmental phenotype	Constipation, lower output, LDL elevation

Laboratory testing is essential when symptoms are nonspecific

- Adult symptoms can be subtle, slow, and clinically overlapping with other conditions.
- Unexplained weight gain and hypercholesterolemia as reasons to assess thyroid function.
- Serum TSH is the most sensitive screening test for primary hypothyroidism.
- The interpretation changes when pituitary or hypothalamic disease is suspected.

Primary hypothyroidism: The feedback logic



Diagnostic interpretation

An increased TSH is a pituitary response to insufficient thyroid hormone; It is not itself proof of a particular thyroid etiology.

Why TSH alone may mislead in central hypothyroidism

Primary thyroid failure

- The hypothalamic–pituitary axis responds to low thyroid output.
- TSH rises because negative feedback decreases.
- A high TSH favors primary disease.

Pituitary / hypothalamic failure

- TSH stimulation is insufficient for the low thyroid hormone state.
- TSH may be low or inappropriately normal.
- A seemingly normal TSH does not exclude central hypothyroidism.

Overt hypothyroidism includes low free T4

- Decreased T4 in overt hypothyroidism from any origin.
- Primary overt disease commonly has elevated TSH with low free T4.
- Central overt disease may have low free T4 with low or inappropriately normal TSH.
- The distinction identifies the site of endocrine dysfunction, not its specific cause.


Subclinical hypothyroidism is a biochemical definition

- Serum TSH is elevated in subclinical primary hypothyroidism.
- Free T4 remains within the normal range.
- Robbins also describes free T3 as normal.
- This laboratory pattern can occur without prominent clinical symptoms.

Interpreting TSH and free T4 patterns

State	TSH	Free T4
Primary overt hypothyroidism	High	Low
Central overt hypothyroidism	Low or inappropriately normal	Low
Subclinical primary hypothyroidism	High	Normal

Laboratory approach to suspected hypothyroidism



Diagnostic interpretation

Do not regard normal TSH as adequate exclusion if central disease is clinically plausible.

Adult case: A slowly evolving multisystem disorder


Clinical vignette

An adult reports fatigue, constipation, cold intolerance, poor exercise tolerance, and progressive facial coarsening. Examination reveals nonpitting soft-tissue swelling.

Reasoning prompt

Which mechanism accounts for edema, and which thyroid measurements distinguish overt primary disease?

Adult case: Pathology and laboratory interpretation



Diagnostic interpretation

The edema is a connective-tissue matrix phenomenon; Fatigue and cold intolerance are systemic physiologic effects of thyroid deficiency.

Interpretation trap: TSH is not always increased


Clinical vignette

A person with hypothalamic–pituitary disease has clinical symptoms compatible with hypothyroidism. The serum TSH is not elevated, but free T4 is low.

Reasoning prompt

Does a non-elevated TSH exclude hypothyroidism? Where may the defect reside?

Central hypothyroidism: Interpret the axis, not TSH alone



Diagnostic interpretation

A TSH within the usual reference interval may be physiologically inappropriate when free T4 is reduced.

Synthesis: Three age-specific diagnostic questions

Child with developmental abnormalities

- Was thyroid hormone lacking during a critical fetal / neonatal interval?
- Could the cause be nutritional iodine deficiency?
- Could an inborn synthetic or developmental thyroid defect account for disease?

Adult with insidious decline

- Are slowing, cold intolerance, dyslipidemia and matrix-rich edema present?
- Do TSH and free T4 indicate overt or subclinical deficiency?
- Is the endocrine defect primary or central?

Five essential conclusions

- Early thyroid hormone deficiency threatens fetal neurodevelopment; Timing is central.
- Iodine deficiency, dyshormonogenesis, and thyroid dysgenesis are different etiologic pathways.
- Adult myxedema combines systemic metabolic slowing with matrix-related tissue swelling.
- Primary hypothyroidism raises TSH; Central disease may not.
- Free T4 distinguishes overt deficiency from subclinical primary hypothyroidism.

Suggested reading

- Robbins, Cotran & Kumar: Pathologic Basis of Disease. 11th edition. Elsevier; 2026.
- Relevant textbook discussion: thyroid physiology, hypothyroidism, and its age-dependent manifestations.
- Review the differences among iodine deficiency, thyroid dysgenesis, and dyshormonogenesis.
- Use authoritative clinical guidance separately for neonatal screening and treatment decisions.