


Conjunctivitis—Trachoma and Other Causes of Painful Blindness

Conjunctival scarring • tear-film failure • corneal injury



Dr Abu Suraih Sakhri

MBBS, MD (Pathology)


www.abusuraihsakhri.com

Learning outcomes

Trachoma and other causes of painful blindness

- Identify the conjunctival structures that maintain the ocular surface.
- Explain how trachoma causes scarring, trichiasis, severe dry eye, and visual loss.
- Compare infectious, chemical, immune-mediated, and iatrogenic pathways to cicatrization.
- Relate conjunctival anatomy to tissue-preserving surgical decisions.


Mechanism schematic



Why conjunctival injury threatens vision


- The conjunctiva contributes to tear-film delivery and maintenance.
- Chronic scarring can damage the fornix, goblet cells, or lacrimal ductules.
- Pain and visual impairment may emerge from loss of ocular surface protection.

Mechanism schematic



Conjunctiva: Three functional zones

- Palpebral conjunctiva lines the inner eyelid.
- Forniceal conjunctiva forms a recess between lid and globe.
- Bulbar conjunctiva covers the ocular surface up to the limbus.



Palpebral conjunctiva: Attachment matters


- The palpebral conjunctiva is firmly tethered to the underlying tarsus.
- Its normal covering is low-profile pseudostratified epithelium.
- Inflammation may produce fine papillary folds on this surface.

Mechanism schematic



Fornix: A specialized protective recess


- The fornix has pseudostratified columnar epithelium.
- Goblet cells are abundant in this region.
- Accessory lacrimal tissue and lacrimal duct outflow are associated with the fornix.



Goblet cells support surface mucin

- Forniceal goblet cells provide mucin to the ocular surface.
- Severe cicatrization may eliminate goblet cells by obliterating the fornix.
- Mucin deficiency contributes to failure of the protective surface film.

Functional loss • mucin



Intact goblet-cell population

**Fornix
cicatrization**




**Reduced mucin
Contribution**

Diagram is conceptual; No cells were imaged.

Lacrimal delivery depends on conjunctival ductules

- Accessory lacrimal tissue lies within the fornix.
- The main lacrimal gland empties through ductules in the superior-lateral fornix.
- Scarring of these ductules diminishes tear delivery.



Bulbar conjunctiva: A different epithelium

- The bulbar conjunctiva consists of nonkeratinizing stratified squamous epithelium.
- It covers the visible globe outside the cornea.
- Its histologic architecture differs from the goblet-cell-rich fornix.


Mechanism schematic



Limbus: Junction and transition


- The limbus marks the transition of conjunctival to corneal epithelium.
- Ocular surface stem cells reside at the limbus.
- Conjunctival pathology must be understood in relation to this junction.

Mechanism schematic




Regional anatomy predicts injury consequences

- Forniceal injury threatens mucin production and aqueous secretion.
- Palpebral contraction can alter eyelid–cornea contact.
- Corneal injury becomes a downstream consequence of conjunctival disease.



Two contributors to the ocular surface film


- Goblet-cell mucin and lacrimal aqueous output are distinct protective inputs.
- Cicatrization can compromise both inputs simultaneously.
- The net result is an inadequately maintained tear film.



Mucin loss: The first failure route

- Extensive fornix loss removes mucin-secreting goblet cells.
- An intact lacrimal gland does not compensate for lost conjunctival mucin.
- The ocular surface becomes vulnerable to chronic irritation.

Functional loss • mucin



Intact goblet-cell population

**Fornix
cicatrization**




**Reduced mucin
Contribution**

Diagram is conceptual; No cells were imaged.

Aqueous loss: The second failure route

- Fibrosis may obstruct the lacrimal ductules crossing the conjunctiva.
- Tear production reaching the ocular surface is then diminished.
- Concomitant goblet-cell damage produces a compound deficit.



Inflammation has region-specific patterns

- Palpebral conjunctivitis can create fine papillary folds.
- Viral conjunctivitis may enlarge lymphoid follicles, especially in the fornix.
- Papillary and follicular responses do not, by themselves, establish a scarring etiology.

Structural / mechanistic distinction

Papillary pattern

Palpebral mucosal folds

Described with allergic
or bacterial
conjunctivitis

Follicular pattern


Lymphoid follicle enlargement

Described with viral
conjunctivitis

Trachoma: The pathogen

- Trachoma is caused by *Chlamydia trachomatis*.
- The bacterium is obligately intracellular.
- Its important late pathology lies in the host tissue response and scarring.


Mechanism schematic



Trachoma: Global significance


- Trachoma as a leading cause of preventable blindness worldwide.
- Its significance arises from progression beyond conjunctivitis to ocular surface destruction.
- The preventable outcome is severe, painful visual loss.

Mechanism schematic




How trachoma spreads: Direct transfer

- Infectious ocular or nasal discharge can contaminate hands.
- Close personal contact allows transfer between individuals.
- The source also identifies clothing, bedding, and hard surfaces as vehicles.



How trachoma spreads: Flies


- Flies contacting infected eye or nose secretions may transport infectious material.
- This is an additional route, not a replacement for person-to-person transfer.
- Transmission occurs before the late fibrotic complications appear.



Infection and blindness are different phases

- Infection initiates conjunctival inflammation.
- Untreated conjunctivitis may progress to major tissue scarring.
- Blindness is a late consequence of accumulated structural injury.


Mechanism schematic



The central injury: Conjunctival cicatrization


- Persistent ocular surface disease may produce fibrous scars.
- Scar contraction alters conjunctival architecture rather than causing mere redness.
- Permanent anatomic distortion is the critical link to chronic surface failure.

Mechanism schematic



Why the fornix may disappear

- The normal fornix provides a recess between eyelid and globe.
- Severe scar formation can obliterate this recess.
- Its loss removes goblet-cell-rich epithelium and compromises tear function.



Tissue loss has functional specificity

- Goblet-cell loss chiefly affects mucin.
- Lacrimal ductule damage chiefly affects aqueous delivery.
- Both are described consequences of severe conjunctival scarring.

Structural / mechanistic distinction

Lost structure

Goblet cells

Mucin contribution


Compromised structure

Lacrimal ductules

Aqueous tear delivery


Trichiasis: A mechanical injury pathway

- Scarring can misdirect eyelashes toward the cornea.
- This abnormal lash orientation is termed trichiasis.
- The corneal surface is then subjected to repeated mechanical contact.



Three pathways converge on the cornea


- Loss of mucin destabilizes the ocular surface film.
- Decreased lacrimal outflow reduces aqueous protection.
- Trichiasis adds direct lash-related irritation.



The end-stage trachoma eye

- The conjunctival fornix may be severely shortened or obliterated.
- The eye is very dry and chronically irritated.
- Corneal damage and visual loss can make blindness painful.


Mechanism schematic



A causal chain, not a single lesion

- Chlamydial conjunctivitis is the initiating condition.
- Scarring damages protective architecture and eyelid mechanics.
- Surface injury is the pathway from conjunctival disease to loss of vision.

Mechanism schematic



Separate infection from structural damage

- The presence of conjunctivitis identifies inflammatory disease.
- Fornix loss, trichiasis, and ductule destruction indicate scarring complications.
- The distinction matters when interpreting advanced clinical findings.

Structural / mechanistic distinction

Inflammatory component

Conjunctivitis

Visible inflammatory reaction


Cicatricial component

Fornix loss ± trichiasis

Altered anatomy and surface function


Which anatomic change explains mucin loss?

- An obliterated fornix eliminates goblet-cell-rich mucosa.
- The associated mucin deficit has a structural basis.
- It cannot be explained solely by an inflamed but preserved palpebral epithelium.



Which change explains reduced tears?


- The main lacrimal gland communicates with the fornix through ductules.
- Conjunctival scarring can compromise this route of secretion.
- Functional loss may occur without primary pathology inside the gland.



Which change explains direct corneal abrasion?

- Conjunctival scarring can redirect eyelashes inward.
- Trichiasis brings lashes into contact with the corneal surface.
- This mechanism is independent of the two tear-film deficits.

Schematic • misdirected lashes



The conjunctiva–cornea interface


- The conjunctiva contributes to ocular surface maintenance.
- The cornea depends on its surface film for normal function.
- Conjunctival disease can therefore impair vision through secondary corneal injury.

Mechanism schematic



Conjunctival scarring has multiple causes

- Trachoma is one infectious cause of conjunctival fibrosis.
- Robbins also identifies caustic alkalis and ocular cicatricial pemphigoid.
- Drugs and surgery can cause iatrogenic conjunctival scarring.



Caustic alkali exposure

- Caustic alkalis are a recognized cause of conjunctival scarring.
- The key late finding is an injured, cicatrized ocular surface.
- Such a history points to chemical injury rather than trachoma.

Conjunctival scarring pathway

Caustic alkali



Conjunctival scar → tear-film injury

Chemical injury: Why the history matters

- An exposure history can identify an alternative scarring pathway.
- The final anatomic consequences may resemble other cicatrizing disorders.
- An observed dry scarred eye alone cannot specify its original cause.

Structural / mechanistic distinction

Historical clue

Caustic contact

Noninfectious onset

Shared late finding

Conjunctival scarring

Surface-film
compromise

Ocular cicatricial pemphigoid

- Ocular cicatricial pemphigoid as another cause of scarring.
- The key relevance here is progressive loss of normal conjunctival architecture.
- The excerpt does not provide immunopathologic or laboratory diagnostic criteria.

Conjunctival scarring pathway


Cicatricial pemphigoid



Cicatrization of the ocular surface

Similar end point, distinct etiologies

- Trachoma is linked to C. Trachomatis infection.
- Caustic exposure is chemical; Cicatricial pemphigoid is an immune-mediated condition.
- All can produce scarring that threatens the surface film.



Drug-related iatrogenic scarring

- Reaction to drugs may scar the conjunctiva.
- The source identifies this possibility without naming specific agents.
- A medication and ocular-treatment history is therefore relevant to attribution.

Conjunctival scarring pathway

Drug-associated reaction




Conjunctival scarring

Surgery-related conjunctival scarring

- Surgical manipulation or removal of conjunctival tissue can create a scarred surface.
- Resection may remove essential goblet cells or damage lacrimal ductules.
- Visual function can be impaired even after successful lesion removal.

Mechanism schematic



Why conventional cancer margins may harm the eye


- Wide excision normally aims to remove invasive disease and adjacent at-risk tissue.
- In the conjunctiva, an overly extensive excision may sacrifice protective surface structures.
- The functional cost may be chronic painful dry eye.

Mechanism schematic




Goblet-cell preservation in tumor surgery

- Goblet cells are concentrated in the fornix.
- Extensive resection of this mucosa can diminish mucin production.
- The extent of excision must be considered against ocular surface preservation.



Lacrimal ductules: Structures to spare

- Main-gland ductules traverse conjunctiva in the superior-lateral fornix.
- Damage can compromise tear delivery.
- The surgical field must therefore be interpreted anatomically, not simply by tumor size.



Invasive tumor versus intraepithelial involvement

- Removing the invasive components of conjunctival neoplasms.
- Intraepithelial disease may instead receive tissue-sparing treatment.
- The goal is to balance disease control with preservation of the ocular surface.

Structural / mechanistic distinction

Invasive component

Excision

Remove invasive tissue

Intraepithelial involvement

Tissue-sparing modality

Limit unnecessary tissue loss

Examples of tissue-sparing modalities

- Cryotherapy for intraepithelial involvement.
- Topical chemotherapy delivered as eye drops is another cited approach.
- These are examples of the pathology-driven preservation principle, not a treatment protocol.

Structural / mechanistic distinction

Invasive component

Local excision

Remove invasive tissue

Intraepithelial disease

Tissue-sparing options

Cryotherapy or topical chemotherapy

A functional model of ocular-surface surgery


- The lesion must be addressed without unnecessary loss of conjunctival function.
- Goblet cells and lacrimal ductules are structures of concern.
- The late adverse outcome is dry, painful, vision-compromising ocular surface disease.

Mechanism schematic



Causes compared: Initiating event


- Trachoma begins with infection and conjunctivitis.
- Alkali exposure begins with chemical injury.
- Pemphigoid, drugs, and surgery provide other routes to conjunctival scarring.



Causes compared: Shared consequence

- Whatever the trigger, scar formation may affect the fornix and ocular surface.
- Tear-film delivery or maintenance may then be compromised.
- Anatomy explains the symptom overlap better than the initiating category alone.

Mechanism schematic



Papillae and follicles are not synonyms

- Papillary folds may occur in palpebral allergic or bacterial conjunctivitis.
- Follicles reflect a lymphoid response; Viral conjunctivitis is an example.
- Neither reaction should be mistaken for the full cicatricial mechanism of blindness.


Structural / mechanistic distinction

Papillae	Follicles
Folded palpebral mucosa	Enlarged lymphoid aggregates
Epithelial surface response	Immune-cell tissue response

Recognizing a cicatrizing phenotype

- A history of chronic conjunctivitis or injurious exposure is relevant.
- Anatomic indicators include fornix alteration and abnormal lash direction.
- Tear-film compromise links these findings to pain and visual loss.

Mechanism schematic



A patient with chronic irritation and trichiasis

- History: Prolonged ocular inflammation.
- Finding: Eyelashes misdirected toward the cornea.
- Question: What scar-related mechanical change explains corneal irritation?

Clinicopathologic reasoning


**Observe the
Structural finding**



**Identify the impaired
Protective function**

Case analysis: The trichiasis mechanism

- Trichiasis is abnormal lash orientation resulting in corneal contact.
- The problem is mechanical even when conjunctival inflammation has subsided.
- Mucin or aqueous deficiency may coexist, but is a distinct pathway.



A patient with a shortened conjunctival fornix

- History: Chronic cicatrizing ocular surface disease.
- Finding: Marked fornix scarring with profound dryness.
- Question: Which two protective inputs are most directly jeopardized?

Clinicopathologic reasoning


**Observe the
Structural finding**



**Identify the impaired
Protective function**

Case analysis: Why fornix loss is decisive

- Fornix obliteration removes goblet-cell-rich mucosa.
- Scarred ductules can obstruct the route for lacrimal secretion.
- One lesion location can produce a dual tear-film failure.



Dry eye after conjunctival tumor excision

- History: Removal of a conjunctival neoplasm.
- Finding: New chronic ocular irritation after surgery.
- Question: How can tumor-directed surgery itself produce surface injury?

Clinicopathologic reasoning

**Observe the
Structural finding**




**Identify the impaired
Protective function**

Case analysis: Anatomical cost of excision

- Conjunctival resection removes epithelium and supporting tissue.
- Loss of goblet cells or lacrimal ductules may reduce surface-film function.
- Tissue-sparing approaches may be considered for intraepithelial involvement.


Mechanism schematic



The shared causal architecture


- Diverse insults produce conjunctival scarring.
- Scarring removes secretory structures or distorts eyelid mechanics.
- Tear-film failure and lash trauma converge on corneal damage and painful visual loss.

Mechanisms leading to corneal injury



Anatomic distinctions to retain


- Palpebral conjunctiva: Tarsal attachment and papillary response.
- Fornix: Goblet cells, accessory lacrimal tissue, and gland ductules.
- Bulbar conjunctiva and limbus: Ocular covering and epithelial transition.



Clinical-pathologic synthesis

- Trachoma is a preventable infectious pathway to conjunctival blindness.
- Late visual loss reflects scarring and failure of ocular surface protection.
- Alkalis, pemphigoid, drugs, and surgery can converge on the same structural end point.
- Conjunctival tissue preservation is central to the clinical logic.

Mechanism schematic



References and scope

Source and limitations

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. Chapter 29, “The Eye,” pp. 1190–1191.
- Conjunctival anatomy and ocular surface context are taken from the preceding “Conjunctiva” subsection.
- All drawn structures are original editable teaching schematics, not clinical photographs or microscopic evidence.
- No external treatment guideline, diagnostic grading system, or therapeutic dose is represented.