

Developmental Anomalies

Female genital tract — vaginal development, morphology and pathological correlations

An illustrated pathology lecture

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning objectives

- Trace Müllerian, mesonephric and urogenital-sinus contributions to the lower genital tract.
- Explain developmental mechanisms underlying a double vagina and vaginal adenosis.
- Recognize the morphology and embryological origin of Gartner and Müllerian cysts.
- Distinguish developmental anomalies from lesions that may simulate neoplasia.

Developmental anomalies in anatomical context

- The vagina joins the uterine cervix to the lower genital tract.
- Abnormal development may affect the vagina and uterus together.
- Anatomical relationships guide interpretation of congenital lesions.

Developmental and anatomical context



Development links uterine and vaginal abnormalities

Embryological contributions are distinct

- Müllerian ducts form the uterus and endocervix.
- Vaginal squamous epithelium advances upward from the urogenital sinus.
- Mesonephric remnants may persist after the ducts regress.

Embryological origins

Müllerian ducts

Urogenital sinus



Uterus and endocervix

Ascending squamous lining

Continuity of the genital tract

Why embryological origin matters

- An abnormal lumen may reflect disordered fusion of paired Müllerian structures.
- A mucosal glandular focus may reflect persistence of fetal epithelium.
- A cyst can indicate persistent embryological duct tissue.
- Anatomy, epithelial phenotype and lesion site therefore provide complementary clues.

Normal development involves coordinated events

- Müllerian structures establish the uterine and endocervical tract.
- Vaginal lining changes from columnar to squamous epithelium.
- Mesonephric ducts regress, although remnants can persist.

Parallel developmental processes

Duct formation and configuration



Epithelial replacement



Regression of residual ducts

Three mechanisms for developmental lesions


Process	Developmental outcome	Example
Incomplete fusion	Duplicated luminal structure	Septate / double vagina
Incomplete epithelial replacement	Persistent glandular mucosa	Vaginal adenosis
Persistent duct remnants	Submucosal cystic lesion	Gartner duct cyst

Müllerian fusion defects alter tract configuration

- A double vagina is associated with failure of Müllerian duct fusion.
- The malformation affects duct architecture rather than epithelial differentiation.
- The term septate or double describes the resulting vaginal configuration.

Fusion failure (conceptual)

Paired Müllerian structures




Separate paired tracts

Fusion and failed fusion: A structural comparison

Normal fusion (conceptual)


Paired Müllerian structures



Convergent tract

Fusion failure (conceptual)

Paired Müllerian structures



Separate paired tracts

Septate or double vagina

- An uncommon developmental anomaly.
- The source describes it as arising from a failure of Müllerian duct fusion.
- Structural duplication distinguishes it from glandular mucosal lesions.
- The associated uterine configuration is clinically important.

Why uterus didelphys accompanies duplication

- The Müllerian ducts contribute to both uterus and endocervix.
- Failure of appropriate fusion can affect anatomically connected regions.
- Double vagina with double uterus (uterus didelphys).

Source-described association

Paired Müllerian ducts



Failure of fusion



Double vagina + uterus didelphys

Structural duplication versus epithelial persistence

- Both represent developmental disturbances.
- They affect different tissue properties:
Configuration versus lining.
- One should not be inferred from the other
without morphologic evidence.

Discriminating features

Double vagina

Abnormal
configuration

Müllerian fusion defect

Associated uterus
didelphys

Vaginal adenosis

Abnormal lining

Residual glandular cells

Mucosal red areas

Genetic syndromes are one etiologic context

- Some anomalies of the external genitalia can be manifestations of genetic syndromes.
- A congenital anomaly is not, by itself, evidence of a specific syndrome.
- The category without assigning a named syndrome or diagnostic test.
- Interpret individual findings within the patient's overall phenotype.

Prenatal exposures can alter development

- In utero exposure to diethylstilbestrol (DES) is cited as an etiologic context.
- DES was historically used to prevent threatened abortions.
- The relevant period of use spans the 1940s to 1960s.
- Exposure pertains to fetal development, not postnatal exposure.

Not all developmental disturbances have a known cause

- Robbins also identifies unknown factors that perturb fetal development.
- Uncertain etiology should not be converted into a specific causal assertion.
- A morphologic phenotype can be characterized without establishing a cause.
- Etiology and anatomical diagnosis are separate levels of inference.

Reciprocal epithelial–stromal signaling

- Fetal tissue development requires interactions between epithelium and its stroma.
- Perturbation of reciprocal signaling as a mechanism.
- Altered tissue interactions may be associated with developmental anomalies.

Conceptual developmental mechanism

Developing epithelium and stroma



Reciprocal tissue signaling



Potential developmental disruption

Separate cause, mechanism and phenotype

- Cause: An exposure, genetic context or unknown influence.
- Developmental mechanism: Altered fusion, replacement or tissue interaction.
- Morphologic phenotype: Duplicated tract, retained glandular mucosa or duct remnant.
- Do not confuse an epidemiologic association with demonstration of the mechanism in an individual case.

Clinical vignette: Duplicated tract

Clinical teaching vignette

An anatomical examination identifies a duplicated vaginal canal. A double uterine configuration is also documented.

What is the best morphologic or developmental interpretation?

Interpretation: Shared Müllerian fusion defect

Interpretation

Double vagina associated with uterus didelphys


- The paired anomalies implicate Müllerian tract configuration.
- The double vagina with failure of duct fusion.
- This is not the epithelial persistence process responsible for adenosis.

The early vaginal lining is columnar


- Embryonal vaginal epithelium is initially columnar.
- Its phenotype resembles endocervical-type epithelium.
- Normal development subsequently changes the predominant lining.

Simplified epithelial phenotypes

Early vaginal lining



Mature vaginal lining



Columnar → squamous replacement

Simplified epithelium schematic — not histology

Squamous replacement occurs during development

- Columnar vaginal epithelium is normally replaced.
- Squamous epithelium advances upward from the urogenital sinus.
- This directional replacement explains the mature vaginal epithelial phenotype.

Normal epithelial differentiation

Early columnar mucosa



Upward squamous advance




Mature squamous lining

Direction of squamous epithelial advance


- The source specifically describes upward movement from the urogenital sinus.
- Columnar cells need not disappear simultaneously throughout the vagina.
- The normal endpoint is predominant squamous lining of the vagina.

Simplified epithelial phenotypes

Early vaginal lining



Mature vaginal lining



Columnar → squamous replacement

Simplified epithelium schematic — not histology

Columnar and squamous vaginal epithelium

Property	Early fetal phenotype	Adult predominant phenotype
Cell type	Columnar, endocervical type	Squamous
Developmental source	Initial vaginal lining	Upward advance from urogenital sinus
Normal relationship	Replaced during development	Predominant mature lining


Residual glandular epithelium may persist

- Small areas of the original columnar lining can remain.
- Persistence into adult life produces a focal epithelial phenotype.
- Robbins calls the retained glandular epithelial areas vaginal adenosis.
- This represents epithelial persistence, not vaginal duplication.

Vaginal adenosis: Histogenetic definition

- Adenosis reflects residual glandular epithelium in the vagina.
- It is characterized by endocervical-type columnar mucinous cells.
- The lining differs from the predominant mature squamous mucosa.

Columnar epithelium retained




Endocervical-type columnar mucinous cells

Simplified epithelium schematic — not histology


Adenosis: Gross and examination appearance

- The abnormal areas appear red and granular on examination.
- They contrast with surrounding pale-pink vaginal mucosa.
- A surface appearance is a clue, not a substitute for epithelial identification.

Surface appearance (conceptual)



Normal mucosa



Adenosis


Red, granular mucosal area

Conceptual appearance — not a clinical photograph

Adenosis: Microscopic epithelial phenotype

- Columnar mucinous epithelium is present.
- It is indistinguishable from endocervical mucosa.
- The distinction from surrounding squamous lining is the diagnostic morphological point.

Columnar epithelium retained



Endocervical-type columnar mucinous cells

Simplified epithelium schematic — not histology

Linking development to visible adenosis

- The developmental event is incomplete replacement of glandular epithelium.
- Persisting mucosal patches form the anatomic substrate of adenosis.
- A red granular examination finding corresponds to the abnormal mucosa.

Development-to-morphology correlation

Persistent columnar cells



Vaginal adenosis



Red, granular mucosal patch

Adenosis should not be mistaken for a cyst

- Adenosis is an epithelial lining abnormality.
- A cyst is a fluid-containing structure within tissue.
- Surface patch and submucosal mass are different morphologic patterns.

Discriminating features

Adenosis

Red granular mucosa

Columnar surface lining

Not defined by a cyst lumen

Ductal cyst

Submucosal fluid cavity

Embryological remnant

Not defined by red mucosa

DES exposure and vaginal adenosis

- Adenosis occurs in only a small percentage of females overall.
- Reports in females exposed to DES in utero show much higher frequencies.
- A reported range of 35% to 90% among exposed females.
- The range describes historical reports rather than a universal individual risk estimate.

Interpreting the reported DES association

Observation	Source-supported statement	Interpretation constraint
General population	Only a small percentage have adenosis	No exact baseline percent supplied
DES-exposed	Adenosis reported in 35%–90%	Broad range across reports
Individual case	Exposure may be relevant	Adenosis does not establish exposure

Why a broad frequency range needs caution

- A range, not a single population estimate.
- The supplied passage does not specify cohorts, methods or denominators.
- Do not calculate an effect size or relative risk from the excerpt.
- The supported conclusion is a strong historical association with prenatal DES exposure.

DES exposure has a historical context

- DES was administered to pregnant individuals for threatened abortion in past decades.
- Fetal exposure preceded recognition of some later vaginal findings.
- The observations of clear cell carcinoma in the 1970s and 1980s altered practice.
- Historical timing matters when interpreting the association.

Rare clear cell carcinoma in DES-associated adenosis

- Rare clear cell carcinomas arising in DES-related adenosis.
- Cases were recognized in teenagers and young females.
- Adenosis is not synonymous with carcinoma.

Association; Not inevitable progression

Prenatal DES exposure



Vaginal adenosis in some exposed females



Rare recorded clear cell carcinoma

Adenosis and clear cell carcinoma are distinct

- Adenosis denotes glandular epithelium in the vaginal lining.
- Clear cell carcinoma is a malignant epithelial tumor.
- The excerpt documents rare cancers in association with DES-related adenosis.
- No transformation probability or surveillance interval is provided.

Avoid overstating the DES–carcinoma link

- The reported association is clinically and historically important.
- Neither adenosis nor prenatal DES exposure means carcinoma is present.
- The textbook passage does not quantify the absolute malignancy risk.

Discriminating features

Supported

Historical association

Rare recorded
carcinomas

Affected young
females

Not established here

Universal progression

A precise lifetime risk

A screening interval

Clinical vignette: Red mucosal patch

Clinical teaching vignette

A vaginal examination shows an irregular red, granular mucosal area against otherwise pale-pink mucosa. The focus contains columnar mucinous epithelium.

What is the best morphologic or developmental interpretation?

Interpretation: Vaginal adenosis

Interpretation

Persistent endocervical-type glandular epithelium

- Columnar mucinous lining is the microscopic defining finding.
- The visible red granular surface matches the textbook description.
- Prenatal DES history is relevant but not required for the morphological diagnosis.

Mesonephric ducts normally regress

- Mesonephric (Wolffian) ducts are present during development.
- They regress as female genital tract differentiation progresses.
- Residual duct tissue can persist in the lower genital tract.

Embryological origins

Mesonephric (Wolffian) ducts



Most regress



Residual duct rests

A source for Gartner duct cysts


Embryonic remnants have predictable locations

- Robbins Figure 22.1 shows mesonephric remnants in ovary, adnexa and cervix.
- The developmental-anomalies section also identifies lateral vaginal wall duct rests.
- Location helps connect a lesion with its embryological origin.
- These remnants must be distinguished from Müllerian-derived epithelium.

Gartner duct cyst: A mesonephric remnant

- The lesion arises from Wolffian (mesonephric) duct rests.
- It is a fluid-filled submucosal cyst.
- The origin is distinct from persistent Müllerian columnar mucosa.

Submucosal cyst — cross-section




Mesonephric duct remnant

Gartner cysts favor the lateral vaginal wall

- The characteristic anatomic site is the lateral vaginal wall.
- “Lateral” identifies position, not epithelial lineage by itself.
- Combining location and cystic morphology strengthens embryologic interpretation.

Lateral wall: Anatomic location




The drawing identifies position, not histology

Gartner duct cyst: Gross dimensions

- Cysts typically measuring 1–2 cm.
- The fluid-filled cavity lies beneath the mucosa.
- The size range is a descriptive feature, not a universal diagnostic cutoff.

Submucosal cyst — cross-section




Mesonephric duct remnant

Why Gartner cyst is a submucosal lesion

- Embryonic duct rests are retained beneath the surface lining.
- Fluid accumulation creates a cystic cavity within the wall.
- The surface epithelium is not itself defined as glandular adenosis.

Submucosal cyst — cross-section



Mesonephric duct remnant


Cyst location and tissue of origin

Lesion	Embryological origin	Described location / morphology
Gartner duct cyst	Mesonephric / Wolffian duct rest	Lateral vaginal wall; 1–2 cm; Fluid-filled
Mucus cyst	Müllerian epithelium	Proximal vagina
Vaginal adenosis	Persistent fetal glandular epithelium	Red granular mucosal patch

Müllerian-derived mucus cysts

- Other vaginal cysts may arise from Müllerian epithelium.
- Robbins specifically mentions mucus cysts.
- These cysts are described in the proximal vagina.

Submucosal cyst — cross-section



Müllerian epithelium

Gartner versus Müllerian vaginal cyst

- The strongest textbook distinctions are embryological origin and site.
- A Gartner cyst is mesonephric-derived and lateral.
- The mucus cyst is Müllerian-derived and proximal.

Discriminating features

Gartner cyst

Mesonephric remnant

Lateral wall

1–2 cm reported

Müllerian mucus cyst

Müllerian epithelium

Proximal vagina

No size range supplied

Why cyst lineage should not be inferred from size

- A 1–2 cm size description applies to Gartner duct cysts in the excerpt.
- No corresponding size distribution is provided for Müllerian mucus cysts.
- A lesion's location and morphology may support but not independently prove its origin.
- Avoid treating a descriptive size as a hard exclusion criterion.

Vaginal endometriosis may simulate neoplasia

- Endometriosis is another Müllerian-derived lesion discussed by Robbins.
- It may occur within the vagina.
- Its clinical appearance can simulate a neoplasm.
- The present developmental-anomalies section does not detail histologic criteria for endometriosis.

Surface and submucosal pattern recognition

Finding	Main consideration	Source-based clue
Red granular mucosal patch	Vaginal adenosis	Columnar mucinous surface epithelium
Lateral wall fluid cyst	Gartner duct cyst	Mesonephric duct remnant
Proximal vaginal mucus cyst	Müllerian-derived cyst	Müllerian epithelial origin
Lesion resembling neoplasm	Vaginal endometriosis is a possibility	Potential clinical mimic

The three axes of morphological assessment

- Site: Lateral vaginal wall, proximal vagina or surface mucosa.
- Configuration: Duplicated canal, epithelial patch or submucosal cyst.
- Cell type: Squamous versus columnar mucinous epithelium.
- Context: Uterine duplication or documented prenatal DES exposure.

An anatomy-first diagnostic reasoning sequence

- First establish whether the anomaly alters structure, lining or forms a cyst.
- Then identify its anatomic distribution and epithelial phenotype.
- Finally interpret embryological origin and any relevant exposure context.

Teaching algorithm, not formal criteria

Define the morphologic pattern



Locate it anatomically



Link to developmental origin

Clinical vignette: Lateral submucosal cyst

Clinical teaching vignette

A small fluid-filled submucosal cyst is found on the lateral vaginal wall. It measures within the 1–2 cm range.

What is the best morphologic or developmental interpretation?

Interpretation: Gartner duct cyst

Interpretation

Persistent mesonephric (Wolffian) duct rest

- Lateral location and submucosal fluid-filled morphology are characteristic.
- The typical 1–2 cm description supports the pattern.
- This differs from superficial adenosis and proximal Müllerian mucus cysts.

Clinical vignette: Proximal mucus cyst

Clinical teaching vignette

A cystic lesion containing mucus is identified in the proximal vagina. The question is its likely embryological derivation.

What is the best morphologic or developmental interpretation?

Interpretation: Müllerian-derived cyst

Interpretation

Müllerian epithelium is the source-described origin

- Mucus cysts in the proximal vagina.
- Gartner duct cysts instead derive from mesonephric duct rests.
- The distinction concerns embryological lineage rather than malignancy.

Integrated comparison of developmental lesions

Entity	Developmental basis	Diagnostic phenotype
Double vagina	Failure of Müllerian duct fusion	Septate / duplicated canal; Uterus didelphys
Vaginal adenosis	Residual fetal glandular lining	Red granular mucosa; Columnar mucinous cells
Gartner duct cyst	Mesonephric duct remnant	Lateral, submucosal, fluid-filled cyst
Müllerian mucus cyst	Müllerian epithelium	Proximal vaginal cyst

High-yield pathology synthesis

- Müllerian duct fusion defects affect tract architecture and may yield a double vagina with uterus didelphys.
- Adenosis reflects retained columnar mucinous epithelium and produces red granular areas.
- Prenatal DES exposure is strongly associated with adenosis; Rare clear cell carcinomas were reported historically.
- Gartner cysts originate from mesonephric remnants; Müllerian cysts and vaginal endometriosis represent different processes.

Final applied question

Clinical teaching vignette

Classify four findings by developmental process: (1) duplicated vaginal canal, (2) red mucosal patch with columnar cells, (3) lateral submucosal fluid cyst, and (4) proximal vaginal mucus cyst.

What is the best morphologic or developmental interpretation?

Final applied question: Explanation

Interpretation

Each pattern points to a distinct developmental mechanism

- (1) Müllerian fusion failure; Associated uterus didelphys.
- (2) Retained fetal columnar glandular lining: Vaginal adenosis.
- (3) Mesonephric remnant: Gartner duct cyst.
- (4) Müllerian epithelial origin: Mucus cyst of proximal vagina.

References and source boundaries

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th edition (2026).
- Embryology and developmental-anomalies discussions in the female genital tract section.
- All anatomical drawings and microscopic-style diagrams in this lecture are original, simplified schematics—not patient images or textbook reproductions.
- The cited passage does not supply formal diagnostic algorithms, molecular testing guidance or follow-up recommendations.