

Dilated Cardiomyopathy

Intrinsic myocardial dysfunction with progressive dilation and impaired systolic contraction

Etiology • pathogenesis • morphology • clinical consequences

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning outcomes

- Distinguish primary DCM from secondary causes of a dilated, poorly contracting heart.
- Explain genetic, inflammatory, toxic, metabolic, and pregnancy-associated pathways.
- Connect chamber dilation and microscopic remodeling with heart failure and embolic risk.
- Apply gross morphology and clinical clues to differential diagnosis.

The defining DCM phenotype

- Progressive cardiac chamber dilation is accompanied by loss of effective contraction.
- Systolic dysfunction is the principal functional abnormality.
- Myocardial hypertrophy commonly coexists despite the dilated, flabby appearance.
- DCM describes a convergent phenotype—not a single etiologic disorder.

From myocyte dysfunction to a dilated heart



A shared anatomic and functional endpoint can arise from many different initiating injuries.

The three major cardiomyopathy patterns

Dilated

Enlarged chamber cavities and ineffective contraction.
Principal dysfunction: Systolic.

Hypertrophic

Marked muscular hypertrophy with impaired ventricular compliance.
Predominantly diastolic dysfunction.

Restrictive

Reduced ventricular compliance restricts filling; Chambers are often not dilated.

DCM among cardiomyopathy patterns

~90%

DCM as the most common
major cardiomyopathy pattern.

- The estimate concerns the textbook's three-pattern categorization.
- Classification by phenotype does not establish an etiologic cause.
- Overlap in clinical manifestations is possible.


Cardiomyopathy versus secondary myocardial dysfunction

Intrinsic cardiomyopathy	Conditions that can mimic it	Clinical implication
Primary myocardial disease	Coronary artery disease / ischemic injury	Assess whether ischemia explains dysfunction.
A DCM phenotype may be idiopathic or specific	Hypertension / valvular disorders	Determine whether loading or valve disease is causal.
Potential genetic or acquired myocardial insult	Congenital heart disease	Evaluate structural disease before labeling primary DCM.

The diagnostic category requires etiologic exclusion

- Poor contraction and dilation are nonspecific features of an end-stage failing heart.
- Primary DCM is considered after major ischemic, valvular, hypertensive, and congenital explanations are excluded.
- An idiopathic designation means that a specific cause has not been identified—not that none exists.

Functional impairment and clinical expression



Remodeling may initially sustain circulation but does not restore normal contractile function.

Functional reference point: Systolic dysfunction

<40%

The Robbins functional-pattern table associates DCM with reduced LV ejection fraction.

- This tabular pattern is not a stand-alone diagnostic criterion.
- LV ejection fraction must be interpreted alongside structural findings.
- Competing causes of low ejection fraction require evaluation.

Severely advanced DCM

<25%

Typical ejection fraction in end-stage DCM.

- This is a description of end-stage disease, not all DCM.
- Poor global contractility explains impaired cardiac performance.
- Clinical severity can exceed what gross histology predicts.

Convergent etiologic pathways

Inherited

Variants affecting myocyte force generation, transmission, or cellular integrity.

Inflammatory / toxic

Myocarditis, alcohol, cardiotoxic exposures, and chemotherapy-associated injury.

Physiologic / metabolic

Peripartum injury, iron excess, tachycardia, endocrine or catecholamine stress.

Familial DCM is increasingly recognized

Up to 50%


A familial contribution in as many as half of cases.

- Family history can guide suspicion but is not universally evident.
- Incomplete penetrance limits inference from unaffected relatives.
- Multifactorial progression complicates genotype–phenotype prediction.

Genes converge on myocyte structural integrity

- DCM-associated variants involve proteins at the cytoskeleton, sarcolemma, and nuclear envelope.
- The functions of affected proteins are diverse; Several cellular routes lead to impaired contraction.
- A single gross phenotype does not imply a single molecular mechanism.

Structural protein defects: Mechanical consequences



Pathogenic pathways may involve impaired force generation, force transmission, or cellular signaling.

Titin: The sarcomeric molecular spring

- TTN encodes titin, a giant protein spanning the sarcomere from Z band toward M band.
- Titin limits passive stretching and contributes to muscle elasticity.
- Titin abnormalities can undermine normal mechanics of the contractile apparatus.

TTN truncations in DCM

10–20%

Approximate proportion of all DCM cases attributed to TTN truncation mutations.

- TTN is identified as the most common genetic cause figure caption.
- The variant class matters: A TTN sequence change alone is not diagnostic.
- Interpretation requires clinical and familial context.

Genetic structures implicated in DCM

Sarcomere

Titin contributes passive elasticity and mechanics across the sarcomere.

Cell membrane / cytoskeleton

Dystrophin links intracellular cytoskeleton to extracellular matrix.

Nuclear envelope

Lamin A/C contributes to nuclear structural organization.

Dystrophin-associated cardiomyopathy

- Dystrophin couples the myocyte cytoskeleton to the extracellular matrix.
- Dystrophin gene defects can accompany Duchenne or Becker muscular dystrophy.
- Some dystrophin mutations manifest primarily as dilated cardiomyopathy.

Lamin A/C and the nuclear envelope

- Nuclear-envelope protein genes, including lamin A/C, are represented among DCM-associated genes.
- The nuclear envelope is one example of a non-sarcomeric cellular compartment implicated in disease.
- Mechanical integrity and intracellular signaling abnormalities may converge on myocyte dysfunction.


Inheritance patterns in familial DCM

Inheritance	Relative frequency	Teaching implication
Autosomal dominant	Predominant	Variable penetrance within families.
X-linked	Less common	Dystrophin-associated disease is an example.
Autosomal recessive	Less common	Recessive variants are among recognized causes.
Mitochondrial	Less common	Defects can impair oxidative phosphorylation.

Variable penetrance complicates family assessment

- Identical familial risk does not imply identical clinical expression.
- Variable penetrance suggests influences beyond the causal genetic variant.
- A clinically unaffected relative cannot, by itself, exclude a familial contribution.

Mitochondrial dysfunction as a DCM pathway



Selected mitochondrial gene deletions with cardiomyopathy, often presenting in pediatric patients.

Fatty acid β -oxidation disorders

- Some inherited DCM phenotypes involve genes encoding enzymes for fatty-acid β -oxidation.
- Impaired metabolic substrate use can compromise the energy-dependent myocardium.
- Metabolic causes are especially relevant in the early-life differential.

Age and genetics: Clinical clues

Childhood

Mitochondrial defects are described as often manifesting in pediatric patients.

Postpubertal onset

X-linked DCM frequently emerges after puberty or in early adult life.

Any age

Inherited disease remains plausible outside these characteristic patterns.

Conduction abnormalities may accompany DCM

- Congenital conduction abnormalities can coexist with the dilated phenotype.
- Robbins relates this association to shared developmental origins of myocytes and conduction elements.
- An arrhythmic presentation may precede obvious severe pump failure.


Myocarditis may evolve into DCM

- Sequential endomyocardial biopsy observations support progression from myocarditis to DCM.
- Inflammatory myocardial injury can produce a persistent contractile phenotype.
- The final dilated morphology may not reveal the initiating myocarditis.

Viral detection: Supportive, not definitive

- Viral genetic material, including Coxsackie B signatures, has been detected in DCM myocardium.
- These findings support a possible link between viral myocarditis and later dilation.
- Viral detection alone does not establish causality in every individual patient.

Alcohol-associated myocardial injury



Nutritional deficiency may be an additional contributor rather than an exclusive explanation.

Thiamine deficiency provides a second alcohol-related route

- Chronic alcohol use can be associated with inadequate thiamine status.
- Thiamine deficiency causes beriberi heart disease, discussed as a form of DCM.
- Concurrent direct toxic and nutritional contributions complicate attribution.

Alcoholic cardiomyopathy has no unique morphology

- Dilated chambers, cardiomyocyte remodeling, and fibrosis are shared with other causes.
- No specific morphologic feature distinguishing alcohol-associated DCM.
- Etiologic attribution requires clinical exposure history rather than histology alone.

Examples of cardiotoxic exposures

Alcohol

Ethanol / acetaldehyde: Direct toxicity and nutritional contributions.

Cobalt

Heavy-metal exposure has produced DCM in documented contamination settings.

Antineoplastic therapy

Cardiotoxic chemotherapeutic agents, including anthracyclines, are recognized contributors.

The differential extends beyond obvious toxins

- Infection, metabolic disorders, neuromuscular disease, and infiltrative processes among heart-muscle disease associations.
- The same source list includes catecholamines, carbon monoxide, and other exposures.
- An exposure list is a diagnostic framework—not proof of causality without case evidence.

Peripartum DCM: Defined clinical window

- The Robbins discussion places peripartum cardiomyopathy late in pregnancy or within 5 months postpartum.
- The ventricular phenotype may resemble DCM from other causes.
- Clinical context is essential because morphology alone cannot determine pregnancy association.

Potential contributors to peripartum cardiomyopathy

Predisposition

Genetic susceptibility and metabolic vulnerability may lower myocardial resilience.


Hemodynamic conditions

Pregnancy-associated hypertension and volume loading may contribute.

Other influences

Nutritional or subtler biochemical disturbances may participate.

Experimental antiangiogenic mechanism



The specific antiangiogenic pathways are supported principally by mouse models and mechanistic inference.


Peripartum pathogenesis: Evidence boundaries

- Antiangiogenic mediators, including sFLT1 and prolactin cleavage products, in experimental systems.
- The proposed mechanism links reduced angiogenesis to microvascular functional ischemic injury.
- This is a mechanistic model, not a validated universal explanation for every human case.

Iron overload can injure the heart

- Iron excess can arise from hereditary hemochromatosis or repeated red-cell transfusions.
- DCM is described as the most common cardiac manifestation of such iron overload.
- The affected myocardium may have dysfunction due to disturbed enzymes and oxidant injury.

Cardiotoxicity from excess iron



These are proposed injury routes; The histologic phenotype of established DCM remains nonspecific.

Supraphysiologic stress pathways

High heart rate

Persistent tachycardia can produce a dilated, hypocontractile phenotype.


Endocrine stimulation

Hyperthyroidism can create excessive cardiac demand.

Adrenergic stimulation

Catecholamine excess, cocaine, or vasopressor exposures can injure myocardium.

Persistent tachycardia: A remodeling pathway



Tachycardia should be considered both a trigger for dysfunction and a possible consequence of cardiomyopathy.


Catecholamine excess has several clinical settings

- Pheochromocytoma can expose the heart to high circulating catecholamines.
- Cocaine and adrenergic vasopressor agents can produce related effects.
- Severe neurologic injury or acute emotional stress may trigger intense autonomic discharge.

Takotsubo: A stress-associated contractile syndrome


- Intense psychological stress may be followed by LV contractile dysfunction.
- The LV apex is often affected, producing characteristic apical ballooning.
- It is in the supraphysiologic-stress discussion; It should not be assumed equivalent to chronic idiopathic DCM.

How catecholamine stress may impair myocytes




Uncertainty about the precise mechanism.

Cardiac dilation: The morphologic centerpiece



Reference configuration



Dilated phenotype

Gross DCM: Enlarged, heavy, flabby heart

- All four chambers can be dilated, and the heart is often markedly enlarged.
- Heart weights often two to three times normal.
- Hypertrophy may coexist with dilation; Wall thickness alone is not an adequate classifier.


A striking increase in heart weight

2–3×


The DCM heart may weigh two to three times the expected normal weight.

- Increased mass can reflect hypertrophy despite enlarged cavities.
- Gross appearance is characteristic but not etiologically specific.
- Degrees of enlargement vary across patients.


Four-chamber dilation: An editable schema



Stasis can produce intracardiac mural thrombi




Why mural thrombi form



A cardiac thrombus is a complication of impaired flow, not the initiating lesion in typical DCM.

Why mitral regurgitation may accompany DCM



The primary pathology is ventricular remodeling; Valvular leaflets need not be intrinsically diseased.

Gross differential: Exclude primary valve disease

- For a primary DCM classification, valve alterations must not themselves explain the myocardial dysfunction.
- Associated mitral or tricuspid regurgitation can be secondary to chamber enlargement.
- Distinguish a dilated heart caused by chronic primary valvular regurgitation.

Coronary artery disease remains a critical exclusion

- Significant coronary stenosis may generate an ischemic dilated phenotype.
- Robbins requires major coronary obstruction to be absent or insufficient to explain the dysfunction in primary DCM.
- The ventricular morphology by itself cannot establish a nonischemic etiology.

Microscopic DCM: Three recurring findings

Myocyte hypertrophy

Many fibers enlarge, with enlarged nuclei.


Attenuated myocytes

Other cells appear stretched, thin, or irregular.

Fibrosis

Variable interstitial and endocardial collagen deposition is common.

Subendocardial scarring: A proposed explanation



This as a probable explanation, not a required morphologic progression.

Microscopy does not identify a unique etiology

- Fibrosis, myocyte hypertrophy, and attenuation are morphologically nonspecific.
- Neither the extent of changes nor one biopsy field necessarily predicts global pump function.
- Histologic abnormalities may not correlate closely with patient prognosis.

“Ninja-star” nuclei and TTN-related DCM

≥5%


Markedly distorted,
hyperchromatic nuclei in this
proportion of myocytes are
described as highly suggestive.

- The finding is specifically associated with titin-truncating mutation cases.
- Morphologic association is not a substitute for genetic confirmation.
- The marker must not be generalized to all DCM phenotypes.

DCM presents across the lifespan

- DCM can present in childhood and adulthood.
- Presentation most often between 20 and 50 years of age.
- Clinical timing can guide etiologic investigation without proving a particular mechanism.

Pump failure explains progressive symptoms



The dominant clinical syndrome is slowly progressive congestive heart failure.

Clinical consequences of advanced DCM

Heart failure

Dyspnea, poor exertional capacity, fatigue, and ongoing pump dysfunction.

Arrhythmias

Abnormal rhythm is frequent and contributes to morbidity and sudden death.

Embolic disease

Mural thrombi may embolize to the systemic circulation.

Rhythm disturbances can be life-threatening

- Abnormal cardiac rhythms are common in the setting of DCM.
- Progressive pump failure and arrhythmias are leading modes of death account.
- Underlying structural or genetic disease can influence arrhythmia susceptibility.

The pathology–outcome relationship is imperfect

- Gross enlargement and fibrosis provide evidence of remodeling but do not specify prognosis on their own.
- Contractile impairment and clinical consequences define functional severity.
- The evaluation must integrate etiology, function, complications, and response to therapy.

Patterns that distinguish DCM from its mimics

Pattern	Dominant functional defect	Characteristic morphology
Dilated cardiomyopathy	Systolic dysfunction	Large chambers; Hypertrophy may coexist.
Hypertrophic cardiomyopathy	Diastolic filling dysfunction	Thick ventricular myocardium; Often no dilation.
Restrictive cardiomyopathy	Impaired diastolic filling	Nondilated ventricles; Atrial dilation common.
Ischemic / valvular disease	May also impair systole	Identify causal coronary or valve abnormality.

Therapeutic implications noted by Robbins

- Pharmacologic heart failure treatment may produce substantial clinical improvement in selected patients.
- Electrical resynchronization by biventricular pacing can benefit appropriately selected patients.
- These interventions address functional consequences but do not make the underlying etiology uniform.

Advanced support options

- Cardiac transplantation is an option in suitable patients with refractory end-stage failure.
- Long-term ventricular assist devices can support circulation.
- In some patients, shorter-term mechanical support permits durable improvement in cardiac function.


Case: A dilated heart with mitral regurgitation

Illustrative vignette

An adult develops progressive exertional dyspnea. Imaging shows an enlarged, globally hypocontractile left ventricle and mitral regurgitation without primary mitral leaflet damage.

What causal sequence explains the regurgitation?

Case reasoning: Secondary mitral regurgitation



The valve leak is a consequence of altered ventricular geometry, not necessarily primary valve pathology.

Case: Etiologic classification

Illustrative vignette

A patient with ventricular dilation, poor contractility, and mural thrombus also has severe coronary obstruction and a history of large myocardial infarctions.

Why is “primary idiopathic DCM” not yet justified?

Case reasoning: A phenotype is not an etiology

- Dilated systolic failure may arise from ischemic myocardial disease.
- Significant coronary disease sufficient to explain dysfunction favors an ischemic cause.
- Primary DCM is considered after competing explanations are excluded.

Case: A familial presentation

Illustrative vignette

Two relatives have a dilated, poorly contracting heart. Another older relative has no cardiac symptoms. A pathogenic TTN truncating variant is under investigation.

Which genetic concepts prevent overinterpretation of the pedigree?

Case reasoning: Penetrance and molecular interpretation

- Variable penetrance means unaffected relatives do not necessarily exclude inherited disease.
- TTN truncating variants are an important association, but any candidate variant needs context.
- Family history, phenotype, and molecular data should be assessed together.

High-yield synthesis: The DCM phenotype

- Intrinsic myocyte dysfunction culminates in chamber dilation and impaired systolic contraction.
- Inherited and acquired pathways may produce the same gross and microscopic appearance.
- Fibrosis and hypertrophy are common but not diagnostic of a specific etiology.
- Clinical hazards include failure, arrhythmias, functional regurgitation, and thromboembolism.

Three points to retain

- DCM is a functional–morphologic syndrome; Seek an explanation rather than assuming idiopathic disease.
- TTN, other structural genes, myocarditis, toxins, peripartum factors, iron, and stress all merit etiologic consideration.
- Chamber dilation explains secondary valve incompetence and thrombogenic blood stasis.

Self-check: Clinical-pathologic reasoning

Question	Answer	Reason
What dominates functionally?	Systolic dysfunction	Contractile failure with chamber dilation.
Why can thromboembolism occur?	Intracavitary blood stasis	Reduced emptying favors mural thrombi.
Does fibrosis identify the cause?	No	Histologic changes are nonspecific.
Does MR require primary valve pathology?	No	Dilation can produce functional regurgitation.

Integrating the DCM phenotype

- The combination of chamber dilation and impaired contraction requires an etiologic explanation.
- Inherited and acquired mechanisms can converge on similar gross and microscopic pathology.
- Fibrosis, hypertrophy, and stasis-related thrombi contribute to morphologic interpretation.
- Clinical assessment must address pump failure, arrhythmias, and embolic risk.