

Dysregulated Nutrient Sensing

Caloric restriction · IGF-1 / mTOR · Sirtuins

Mechanisms linking nutrient-responsive signaling to cellular aging

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning outcomes

- Connect nutrient-responsive signaling with the biology of cellular aging.
- Trace growth hormone, IGF-1, mTOR, and rapamycin model.
- Explain how sirtuins link metabolic adaptation with stress responses and DNA repair.
- Critically distinguish model-organism findings from evidence for human longevity.

Cellular aging is a mechanistic process

- Aging entails progressive loss of cellular function and viability.
- Accumulating molecular damage and genetic abnormalities contribute to the decline.
- Both cell-intrinsic changes and environmental insults influence the process.
- Conserved genetic regulation means aging is not merely passive “wear and tear.”

Nutrient sensing belongs within cellular aging




These mechanisms interact; Nutrient sensing is not a stand-alone cause of aging.

Nutrient sensing: The central concept

- Cells respond to nutrient availability by changing metabolism and growth-related programs.
- The growth-promoting state is beneficial for tissue function but carries metabolic demands.
- The aging literature asks whether reducing selected nutrient-responsive signals changes longevity.
- Insulin/IGF-1–mTOR signaling and sirtuins.

Here “dysregulated” refers to nutrient-response pathways in the wider framework of aging.

From nutrient input to cellular outcome



Pathway activity is a mediator—not a direct measurement of biological age.

Why study conserved pathways?

- Experiments from yeast to mammals identify genetic control of longevity.
- Evolutionarily conserved regulation supports a biological mechanism rather than coincidence.
- Conservation does not guarantee an identical effect size or safety profile in people.
- Interpretation must keep organism, intervention, and outcome distinct.

Different claims require different evidence

Mechanism

A pathway changes
metabolism or cellular
repair.

Animal longevity


An intervention extends
lifespan in tested models.

Human benefit

A clinical outcome improves
safely in people.

Stronger model-organism evidence than definitive human intervention evidence.

Caloric restriction: Proposed antiaging logic



The pathways are proposed mediators; This is not a prescription for patients.

What the textbook says about lifespan

- Caloric restriction has increased lifespan across tested eukaryotic experimental systems.
- Results in nonhuman primates are described as encouraging.
- Observations in humans do not establish a general causal longevity effect.
- Interventions intended to prolong human life remain largely experimental.

Interpret caloric-restriction evidence

Evidence base	Defensible interpretation
Tested eukaryotic models	Consistent experimental support for lifespan effects
Nonhuman primates	Encouraging—not equivalent to proven human benefit
Humans	Do not infer lifespan extension from limited observations

Avoid treating species-to-species extrapolation as direct clinical evidence.

Longevity and healthy aging are related, not identical

Lifespan

- Time until death is the outcome in lifespan experiments.
- Model-organism lifespan is a measurable experimental endpoint.

Age-related pathology

- Disease burden and functional decline represent different outcomes.
- MTOR inhibition was reported to reduce age-related pathologic changes in animal models.

Always identify which outcome a study actually measured.

Nutrient-sensing components

Insulin / IGF-1

Signals nutrient availability and favors anabolic growth.

MTOR


Downstream kinase associated with growth and autophagy control.

Sirtuins

NAD-dependent enzymes that coordinate stress adaptation and repair.


MTOR is a prominent target downstream of IGF-1, not a third independent endocrine circuit.

Caloric restriction engages two proposed routes



Parallel proposed pathways may jointly contribute to lifespan effects in models.

Growth hormone links pituitary to IGF-1



IGF-1 production in many cell types.


IGF-1 production is not confined to one tissue

- Multiple cell types can synthesize IGF-1 in response to growth hormone.
- IGF-1 is part of the growth hormone–responsive cellular signaling network.
- For this lecture, its relevance is the link between systemic nutrition and cell growth.
- Aging effects are inferred from altered signaling intensity rather than IGF-1 abundance alone.

IGF-1 resembles insulin at the signaling level

- IGF-1 activates intracellular signals similar to those activated by insulin.
- account, this informs cells about glucose availability.
- The resulting response favors anabolic metabolism.
- Anabolic signaling also supports cellular growth and replication.

What an anabolic IGF-1 signal accomplishes



Anabolism is an adaptive physiological program, not intrinsically pathological.

Growth and metabolism: Linked responses

Strong IGF-1 signaling

- Favors anabolic activity.
- Promotes cell growth and replication.
- Reflects a nutrient-available state.


Attenuated signaling

- Associated with less cell growth.
- Associated with lower metabolic activity.
- May reduce subsequent cellular damage.

Attenuation is not the same as elimination

- The longevity association concerns reduced intensity of IGF-1 signaling.
- Normal IGF-1 pathways are involved in legitimate growth and metabolic functions.
- The textbook does not propose total blockade as a human therapeutic goal.
- A lower-signal phenotype must be interpreted within the experimental context.

Why less growth signaling may favor longevity



Robbins explicitly qualifies reduced cellular damage as a possibility, not a settled mechanism.

IGF-1 evidence: Fact versus proposal

Textbook statement	Interpretation
IGF-1 responds to growth hormone	Established source relationship
IGF-1 favors growth and replication	Established physiologic role in this account
Attenuation lowers growth / metabolism	Model-supported response
Reduced damage explains longevity	Plausible; Stated cautiously

IGF-1 has several downstream effects




Do not simplify all IGF-1 effects into a single mTOR-only route.

MTOR is a prominent nutrient-responsive kinase


- MTOR stands for mechanistic target of rapamycin.
- It is a major downstream kinase in IGF-1 signaling.
- It connects the growth-response network with cellular maintenance pathways.
- The aging literature examines how reducing mTOR activity changes outcomes.

The IGF-1–mTOR axis



Schematic summary; Other downstream signaling targets are not shown.

Rapamycin: Experimentally perturbing mTOR



A drug effect in an animal experiment is not evidence of safe preventive use in humans.


Rapamycin and longevity in animal models

- Extended lifespan with mTOR inhibition in diverse animal models.
- It specifically notes benefit after administration to middle-aged mice.
- Rapamycin affects several cellular processes; Autophagy is among them.
- The observed lifespan result does not identify one exclusive mediator.

Rapamycin and age-associated pathology

- In animal models, mTOR inhibition can reduce age-related pathological changes.
- This is a different endpoint from lifespan extension.
- Interpret both benefits at the level of the organism and experimental design.
- The text does not establish a standard antiaging treatment for healthy people.

How mTOR inhibition connects to autophagy



The source states promotion of autophagy; Causal attribution of all longevity effects remains broader.

Protein homeostasis depends on two systems


Keep proteins folded

- Molecular chaperones maintain correct protein conformations.
- Heat-shock protein chaperones provide experimental aging examples.

Clear unsuitable proteins


- Autophagy–lysosome removes material for degradation.
- Ubiquitin–proteasome degrades selected unwanted proteins.

Chaperones protect the functional proteome



Impaired folding is one recognized contributor to aging-related proteostasis failure.

Two clearance routes remove unsuitable proteins




Parallel disposal systems contribute to protein homeostasis.

Proteostasis deteriorates with aging


- Both correct protein folding and disposal of damaged proteins become less efficient.
- Experimental chaperone deficiency accelerates aging in mice.
- Chaperone overexpression has been associated with longer life in mice.
- Similar model-based evidence implicates autophagy and proteasomal degradation.

When misfolded proteins accumulate



This is a cellular pathology consequence—not a histologic test for “nutrient-sensing disease.”

MTOR inhibition has multiple effects



Do not claim that autophagy alone accounts for all observed survival benefits.

Sirtuins are NAD-dependent deacetylases

- Sirtuins are a family of NAD-dependent protein deacetylases.
- They respond to environmental and metabolic stresses.
- Their roles extend beyond one substrate or one cellular compartment.
- In the source model, higher sirtuin activity is proposed to contribute to caloric-restriction effects.

Three defining properties of sirtuins

Enzyme family

At least seven types are described in mammals.

NAD dependent

Activity is coupled to a metabolic cofactor.


Distinct roles

Compartmental distribution and nonredundant functions.

Sirtuin diversity limits simple generalizations


- Mammalian sirtuins occupy different cellular compartments.
- Their functions are not interchangeable.
- An effect attributed to one sirtuin need not apply to all seven.
- The text highlights sirtuin-6 for combined metabolic and genome-maintenance functions.

Food deprivation recruits adaptive responses




This is a proposed framework for caloric-restriction adaptation.

DNA damage is also a sirtuin-related stimulus



Robbins particularly emphasizes sirtuin-6 in activating DNA repair enzymes.

Sirtuins influence several adaptive outputs



These are source-described effects of regulated gene expression—not unconditional effects in every tissue.


Sirtuins and metabolic restraint

- Sirtuin-dependent gene programs may inhibit metabolic activity.
- A lower metabolic program may aid adaptation to food limitation.
- This complements the decline in growth-related IGF-1 signaling.
- Such interactions are presented as candidate explanations of longevity.

Sirtuins and apoptosis


- Decreased apoptosis among sirtuin-associated adaptive effects.
- A stress-survival response can preserve viable cells during adverse conditions.
- This does not mean blocking apoptosis is universally beneficial.
- Cellular context remains essential when interpreting any antiapoptotic signal.

Sirtuins and protein folding



Connect this route to the parallel aging mechanism of defective protein homeostasis.

Counteracting oxygen-radical injury



The textbook does not provide a numerical ROS reduction or a human effect size.

Sirtuins and glucose handling

- The source states that sirtuins can increase insulin sensitivity.
- It also describes effects that increase glucose metabolism.
- These actions create therapeutic interest in diabetes.
- A plausible disease target is not the same as validated clinical efficacy.

Sirtuin-6 has two important roles

Nutritional adaptation


- Helps mediate metabolic changes during caloric restriction.
- Places nutrient availability in a cellular stress-response context.

Genome preservation

- Promotes genomic integrity.
- Activates DNA repair enzymes through deacylation.

This dual function is explicitly emphasized.

Sirtuin-6 connects enzyme activity to repair



The text specifies deacylation here; Do not replace it with an invented detailed molecular cascade.

Sirtuins bridge distinct aging mechanisms

Metabolic response

Adaptation during reduced nutrient intake.

Protein homeostasis

Support protein-folding responses.

DNA integrity


Activate repair machinery, particularly via sirtuin-6.

IGF-1/mTOR and sirtuins are complementary

Dimension	IGF-1 / mTOR axis	Sirtuins
Primary function	Signals nutrients and growth	Adapts to deprivation / stress
Restriction model	Signal intensity decreases	Activity increases
Main source outputs	Less growth, altered autophagy	Metabolic adjustment, folding, repair
Human clinical status	Experimental longevity strategies	Experimental longevity strategies


This is a model-specific comparison—not a binary “bad vs good” classification.

Integrated proposed response to caloric restriction




The branches are conceptual responses, not a demonstrated linear cascade.

Mechanisms of aging converge on cell function



Different proximal mechanisms can converge on cell dysfunction and cell loss.

Aging links cellular changes to organ disease



Atherosclerosis and type 2 diabetes as illustrative chronic-disease consequences.

Nutrient sensing has no unique diagnostic morphology

- IGF-1/mTOR and sirtuin activity are biochemical signaling states, not standalone histologic entities.
- Morphologic manifestations are indirect and depend on the tissue and broader disease process.
- Microscopy cannot, by itself, establish a causal nutrient-sensing defect.
- Interpret claims using pathway biology, model system, and relevant clinical endpoints.

No synthetic H&E or immunostain is presented as evidence for this abstract mechanism.

Match the claim to the evidence

Claim	Evidence
Caloric restriction prolongs life	Strong model-organism evidence; Primate findings encouraging
Reduced mTOR improves aging phenotypes	Animal lifespan and pathology outcomes
Sirtuins affect stress responses	Experimental molecular and model-system findings
Antiaging medication extends human life	Not established in the supplied section

The clinical translation remains unresolved

- The textbook cautions that longevity interventions remain largely experimental.
- Long-term human safety and efficacy are not fully understood.
- Rapamycin animal findings do not justify routine preventive administration.
- Sirtuin-based approaches are mechanistically interesting but not proven longevity therapies.

A popular claim is not a therapeutic conclusion

- The source mentions publicity about a red-wine constituent and sirtuin activation.
- A biochemical possibility does not establish a lifespan effect in humans.
- Food or supplement narratives cannot replace controlled clinical outcome studies.
- Keep proposed targets separate from validated interventions.

Application: Reduced IGF-1 signaling

An experimental model shows lower IGF-1 signaling and reduced rates of cell growth and metabolism. Which proposed aging mechanism is most consistent with this phenotype?

- A. Intensified anabolic drive
- B. Attenuation of nutrient-responsive growth signaling
- C. Loss of all growth hormone production
- D. A specific histologic pattern of nutrient-sensing failure

Application answer: Attenuation of growth signaling

- Correct: B. Lower IGF-1 activity attenuates nutrient-responsive growth signals.
- The text links this to lower growth and metabolic activity.
- Reduced cellular damage remains a proposed explanation, not a demonstrated necessity.
- The scenario alone says nothing about a diagnostic morphology or human treatment.

Application: Rapamycin and experimental aging


A mouse study reports longer lifespan after rapamycin exposure. What inference is best supported by the Robbins discussion?

- A. All human aging pathways are now explained
- B. MTOR inhibition can influence lifespan in animal models
- C. Autophagy is proven to be the only causal mediator
- D. Long-term preventive therapy is established in humans

Application answer: Identify the evidence level

- Correct: B. Animal models support a longevity effect of mTOR inhibition.
- Rapamycin has multiple biological effects, including increased autophagy.
- A single observed outcome does not establish a unique mechanism.
- Human safety, benefit, and suitable clinical application remain separate questions.

Synthesize the chapter model



The strongest argument is mechanistic coherence plus animal experiments; Clinical translation requires direct testing.

Five conclusions to retain

- Caloric restriction is linked to longer lifespan in experimental eukaryotic systems.
- Reduced insulin/IGF-1–mTOR signaling is a major proposed mediator.
- MTOR inhibition can extend lifespan and modify age-related pathology in animal models.
- Sirtuins coordinate adaptive metabolism, proteostasis, and genome maintenance.
- None of these results by itself proves a safe human antiaging treatment.

Selected source

Robbins, Cotran & Kumar

Pathologic Basis of Disease

Cell Injury, Cell Death, and Adaptations

Cellular Aging — Dysregulated Nutrient Sensing

Diagrams are original, editable teaching schematics.