

Entamoeba histolytica

Invasive amoebic colitis and extraintestinal dissemination

Etiology · pathogenesis · morphology · diagnosis

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Trace cyst transmission to colonic invasion and hepatic dissemination.
- Explain why amoebic ulcers have a narrow neck and broad base.
- Recognize major colonic and hepatic pathologic features.
- Select diagnostic approaches while acknowledging microscopy limits.
- Relate disease compartments to antimicrobial strategy.

Amebiasis: Invasive protozoal infection

- *Entamoeba histolytica* is an intestinal protozoan pathogen.
- Amebiasis includes colonic disease and extraintestinal spread.
- Invasive infection can cause dysentery and liver abscess.

Central theme: Tissue invasion explains the pathology.

Three major infection states

E. Histolytica intestinal colonization

```
graph TD; A["E. Histolytica intestinal colonization"] --> B["Luminal carriage"]; A --> C["Invasive colitis"]; A --> D["Extraintestinal"]; B --> E["No tissue invasion; May be asymptomatic"]; C --> F["Epithelial damage, ulcers and dysentery"]; D --> G["Vascular spread; Liver most important"];
```

Luminal carriage

No tissue invasion; May be asymptomatic

Invasive colitis

Epithelial damage, ulcers and dysentery

Extraintestinal

Vascular spread; Liver most important

Epidemiologic exposure is fecal–oral

- Contaminated food, water or hands transmit infectious cysts.
- Transmission is favored by inadequate sanitation.
- History of residence, travel and contaminated-water exposure can guide suspicion.

Sexual transmission is also possible

- Oral–anal exposure can transmit the organism.
- Transmission remains biologically fecal–oral.
- Risk assessment should be respectful and behavior-specific.

The cyst is the infectious form

Mature cyst

Protective cyst wall

Survives the gastric environment

Ingested to initiate infection

Survives outside host for transmission

Trophozoite


Motile, amoeboid stage

Replicates in intestinal lumen

Can invade colonic epithelium

Poorly survives gastric and external exposure

Cyst morphology: An editable guide




Chitin-rich protective wall

Four nuclei in mature cyst

Infectious environmental stage

Why cysts traverse the stomach



Physical wall protection enables fecal–oral transmission.


Excystation precedes colonization




Trophozoites proliferate in the colon

- Motile trophozoites multiply by binary fission.
- The large intestine is the major site of residence.
- They may persist in the lumen or invade mucosal tissue.

Encystation completes the transmission cycle



Integrated intestinal life cycle



The invasive branch—not mere luminal carriage—produces tissue injury.

Most frequent colonic sites

Cecum

Common

Ascending colon

Common

Sigmoid colon

Possible

Rectum

Possible


Appendix

Possible

Colonic predilection has a clinical correlate

- Cecum and ascending colon are most commonly affected.
- Sigmoid colon, rectum and appendix may also be involved.
- Distribution alone does not establish an etiologic diagnosis.


Invasion begins with epithelial adherence




Host epithelial apoptosis is induced

- After attachment, amoebae can induce epithelial apoptosis.
- Cell loss compromises mucosal epithelial integrity.
- The weakened surface permits progression into crypts.

The organisms invade colonic crypts



Lateral extension destroys lamina propria



Epithelial adherence


Entry → lateral lamina propria injury

Schematic colonic wall; Lateral spread within lamina propria.


The neutrophilic response contributes to injury

- Invasion recruits neutrophils to the involved mucosa.
- Neutrophil recruitment accompanies tissue destruction.
- Pathology reflects parasite-mediated damage and host inflammation.

From surface invasion to destructive ulcer



The flask-shaped ulcer is a tissue geometry



Schematic tissue cross-section; Not a histologic specimen.

Why the ulcer opening is narrow

- A small mucosal entry zone may overlie a larger damaged area.
- The organism then spreads laterally in lamina propria.
- Thus superficial inspection can underestimate deep tissue damage.

Why the ulcer base is broad

- Burrowing in underlying tissue creates lateral extension.
- The ulcer expands beneath relatively narrower surface loss.
- The cross-sectional appearance is classically flask shaped.

Amebic colitis: Morphologic synthesis

Luminal / surface events

Trophozoite adherence

Surface epithelial apoptosis

Crypt entry

Narrow mucosal ulcer mouth

Deep tissue events


Lateral lamina propria spread

Neutrophil recruitment and damage

Broad undermining ulcer base

Potential vascular access

Trophozoites: Conceptual morphology



Single nucleus; Central karyosome

Amoeboid trophozoite

Internal red discs: Ingested erythrocytes

Erythrophagocytosis is a useful clue

- Red cells may be seen inside tissue-associated trophozoites.
- Erythrophagocytosis in a colonic biopsy.
- A phagocytosed erythrocyte is an observation, not a standalone molecular species confirmation.

Amoebae can resemble macrophages

Trophozoites

May resemble macrophages in size

Amoeboid cytoplasm and nucleus

Internalized erythrocytes may be present

Look for organisms near ulcerated tissue

Histiocytes / macrophages

Host inflammatory cells

Similar overall silhouette in H&E

May contain phagocytosed debris

Require attention to cell identity and context

A tissue identification problem

- Inflammation and necrosis can obscure organism margins.
- Some amoebae resemble host histiocytes.
- Use nuclear details, tissue distribution and confirmatory tests when needed.

What a histology description must establish

- Where are putative trophozoites located?
- Are organisms associated with epithelial damage or ulceration?
- Is erythrophagocytosis visible?
- Could the cells instead represent host macrophages?

Stool trophozoites: CDC morphologic features

- Usually a single nucleus with a small karyosome.
- Fine, relatively uniform peripheral nuclear chromatin.
- Granular or ground-glass cytoplasm.
- Motile forms are common in diarrheal stool.

Stool cysts: CDC morphologic features

- Mature cyst typically contains four nuclei.
- A central karyosome and fine peripheral chromatin are characteristic.
- Chromatoid bodies may have blunt rounded ends.
- Cyst morphology is not species-specific within the histolytica/dispar complex.

Microscopy alone has a major limitation

- *E. Histolytica*, *E. Dispar* and some other *Entamoeba* species can look identical.
- A morphologic report may not establish pathogenic species.
- Species-specific antigen or molecular testing can resolve uncertainty.

Ingested erythrocytes are not absolute proof

- Classically associated with *E. Histolytica* trophozoites.
- Rare erythrophagocytosis has also been reported in *E. Dispar*.
- Do not claim definitive species identification from RBC ingestion alone.

E. Histolytica versus E. Dispar

E. Histolytica

Recognized invasive pathogen

May cause colitis or liver abscess

Trophozoites can contain erythrocytes

Species-specific confirmation is clinically useful

E. Dispar

Generally nonpathogenic colonizer

Usually does not cause invasive disease

Morphology overlaps with E. Histolytica

Avoid misclassifying incidental carriage

Amebic versus bacterial dysentery

Amebic colitis

Protozoal tissue invasion

Flask-shaped colonic ulcers

Trophozoites may be seen

Extraintestinal hepatic spread possible

Bacterial invasive diarrhea


Invasive bacteria include Shigella and Salmonella

Exudative bloody diarrhea can occur

Bacterial studies are required for identification

Mechanisms and management differ

How vascular dissemination begins



The liver is the major extraintestinal target

Colonic tissue injury



Splanchnic vascular
invasion




Hepatic seeding



Amebic liver abscess

Liver abscess: Pathologic architecture



Abscess cavity

Central necrotic cavity

Shaggy fibrinous lining

Scant marginal inflammatory reaction

Native-shape liver and lesion schematic; Not gross pathology.

Liver abscesses can become large

- Lesions exceeding 10 cm in diameter.
- A large necrotic cavity may form in hepatic parenchyma.
- Size does not by itself establish the microbial cause of a liver lesion.


Margins need not be densely inflamed

- Scant inflammatory reaction at abscess margins.
- The lining may be shaggy and fibrinous.
- Morphologic impression must be integrated with microbiology and clinical setting.

Hepatic infection may outlast intestinal symptoms

- An amebic liver abscess may persist after acute intestinal illness.
- Current diarrhea may therefore be absent at presentation.
- A negative stool examination cannot exclude hepatic disease.


Local extension may involve thoracic organs



Bloodstream spread can reach other organs

- Extraintestinal spread beyond liver is uncommon.
- Hematogenous dissemination to kidney and brain.
- Unusual sites require specific clinical and laboratory confirmation.

Clinical spectrum: Intestinal amebiasis



Bloody diarrhea reflects destructive colitis

- Epithelial and lamina propria destruction breach the mucosal barrier.
- Bleeding and inflammatory exudate produce dysenteric stools.
- Bloody diarrhea warrants evaluation for multiple invasive pathogens.

Weight loss reflects sustained intestinal illness

- Persistent abdominal pain and diarrhea may reduce intake.
- Colonic mucosal disease can have systemic consequences.
- Clinical weight loss is a clue, not a diagnostic criterion by itself.

Fulminant necrotizing colitis is a danger

- The infection can occasionally produce acute necrotizing colitis.
- Severe tissue loss raises risk of catastrophic bowel complications.
- Rapid recognition and specialist assessment are essential.

Megacolon signals severe disease

- Megacolon is an uncommon but grave complication.
- It occurs in the setting of severe colonic inflammation and dysfunction.
- Necrotizing colitis and megacolon with significant mortality.

Three clues to extraintestinal disease

- A patient may have past, but not current, dysentery.
- A focal liver lesion can arise from prior colonic invasion.
- Consider amebiasis when epidemiology and hepatic findings align.

Diagnostic investigation is compartment specific

Suspected amebiasis

```
graph TD; A[Suspected amebiasis] --> B[Intestinal disease]; A --> C[Colonic tissue]; A --> D[Hepatic abscess]; B --> E[Stool analysis; Molecular or specific antigen tests]; C --> F[Biopsy for invasive lesion and trophozoites]; D --> G[Imaging + appropriate serology or molecular testing];
```

Intestinal disease

Stool analysis; Molecular or specific antigen tests

Colonic tissue

Biopsy for invasive lesion and trophozoites

Hepatic abscess

Imaging + appropriate serology or molecular testing

Stool sampling affects yield

- Parasites are not uniformly detected in every specimen.
- Examination of multiple stools improves microscopy sensitivity.
- Specimen handling must match microscopy, antigen or molecular method.

Species-specific antigen assays

- Selected immunoassays can detect *E. Histolytica* antigens.
- Assay specificity varies: Not every kit distinguishes *E. Dispar*.
- Follow test-specific specimen requirements, often fresh unfixed stool.

PCR can establish species identification

- Molecular assays differentiate *E. Histolytica* from related *Entamoeba* species.
- Stool and selected extraintestinal specimens may be tested.
- Specimen preservation is critical for DNA-based methods.

Antibody testing in liver abscess

- Serologic tests are especially useful in extraintestinal disease.
- Prior endemic exposure may complicate interpretation.
- Interpret antibody evidence alongside imaging and clinical history.

A biopsy is diagnostic only in context

- Trophozoites may be seen in endoscopic or surgical tissue.
- Host macrophages can be confusing on H&E.
- Organism identification and species confirmation are distinct tasks.

An integrated diagnostic sequence



Why tissue-active treatment is required

- Invasion creates organisms within colonic tissue and extraintestinal sites.
- Metronidazole for systemic amebiasis.
- Treatment must address invasive trophozoites.

Tissue disease and luminal cyst carriage differ

Invasive tissue infection

Trophozoites cause active tissue injury

Nitroimidazole therapy is tissue active

Examples: Metronidazole or tinidazole

Specific therapy determined clinically

Residual luminal infection


Residual intestinal cyst carriage may persist

Follow with a luminal agent after invasive treatment

Example: Paromomycin

Prevents persistence and onward transmission

Prevention targets the infectious cyst



Prevention follows directly from the fecal–oral life cycle.

Case 1: Invasive colitis

A returning traveler has abdominal cramping and bloody diarrhea. Colonoscopy identifies a deep ulcer; Biopsy shows macrophage-like organisms near necrotic tissue.

Which feature most strongly suggests amebic colitis?

Case 1: Pathologic resolution

- Look for flask-shaped undermining ulceration.
- Seek trophozoites within injured tissue; Assess for erythrophagocytosis.
- Confirm pathogenic *Entamoeba* where available.
- Do not mistake tissue histiocytes for parasites.

Case 2: Delayed hepatic disease

A patient has a liver abscess several weeks after a self-limited episode of diarrhea. Current stool examination is unrevealing.

Does a negative stool examination exclude amebic liver abscess?

Case 2: Pathologic resolution

- No: Hepatic disease can persist after intestinal symptoms resolve.
- Obtain appropriate hepatic imaging and targeted testing.
- Serology can be helpful in an appropriate exposure setting.
- Evaluate alternatives, including bacterial liver abscess.

Case 3: A species-identification trap

A stool report states “E. Histolytica/E. Dispar complex” based on microscopy. The patient has no symptoms.

Can this microscopy result alone prove invasive amebiasis?

Case 3: Diagnostic resolution

- No: Cysts and trophozoites of several species overlap morphologically.
- Asymptomatic colonization does not establish invasive pathology.
- Species-specific antigen or molecular testing distinguishes clinically relevant infection.

Contrast with other intestinal protozoa

Entamoeba histolytica

Invasive colonic disease

Flask-shaped colonic ulcer

Can seed hepatic abscess

Bloody diarrhea may occur

Giardia / Cryptosporidium


Predominantly noninvasive enteric injury

Giardia: Brush-border dysfunction

Cryptosporidium: Enterocyte-apical localization

Watery diarrhea / malabsorption emphasized

Mechanism-to-morphology chain



Clinicopathologic priorities

- Identify the infectious stage and the invasive trophozoite.
- Explain the narrow-neck, broad-base ulcer by lateral extension.
- Recognize vascular spread and the distinctive hepatic abscess.
- Use species-resolving diagnostics where microscopy is ambiguous.
- Address tissue infection and residual luminal organisms when treating invasive disease.

Selected teaching references

- Robbins, Cotran & Kumar: Pathologic Basis of Disease, 11th edition — gastrointestinal infection section.
- CDC DPDx: Amebiasis — life cycle, morphology and diagnostic methods.
- CDC DPDx: Stool antigen detection and molecular diagnosis.
- CDC Yellow Book / Merck Manual Professional: treatment context.