

Fibrinous Inflammation

Morphogenesis • Serosal pathology • Resolution and organization

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

Explain why fibrinogen reaches an injured tissue.

Recognize fibrinous exudate grossly and histologically.


Predict resolution versus organization and its consequences.

Why the morphologic pattern matters

The dominant exudate provides clues to injury intensity and tissue site.

Pattern recognition complements—not replaces—identification of the cause.

Fibrinous change is especially informative on serosal surfaces.




Defining the fibrinous exudate

Fibrin is formed in the extracellular compartment after plasma proteins escape.

A local procoagulant stimulus accompanies increased permeability.

Deposits can line inflamed body cavities and organ surfaces.




Fibrinous inflammation is an acute pattern

Acute inflammation involves vascular dilation, fluid leakage and leukocyte recruitment.

The mixture of exuded material varies with cause, intensity and anatomic site.

Fibrin deposition is a particular morphologic expression of this response.




Two requirements for fibrin deposition

Vascular permeability must permit large plasma proteins to escape.

An appropriate local procoagulant environment must generate fibrin.

Neither simple fluid accumulation nor protein leakage alone defines the pattern.



Principal locations: Body-cavity linings

Pericardial serosa around the heart.

Pleural serosa surrounding the lungs.

Meningeal surfaces; Distribution is context dependent.

Serosal surface in cross-section

Fibrin on surface



Serosal lining


Schematic—not a tissue section

Organ surface versus organ substance

Fibrin typically coats an inflamed serosal surface.

Adjacent tissue may also exhibit vascular dilation and recruited inflammatory cells.

The visible surface exudate should not be confused with an intravascular thrombus.




From vessel lumen to tissue compartment

Plasma fibrinogen normally circulates inside blood vessels.

Permeability changes create access to the extracellular compartment.

Extravascular fibrin then becomes a morphologic hallmark.




Early vascular reactions in acute inflammation

Vasodilation increases local blood flow.

Altered endothelial permeability enables plasma protein exudation.

Circulating leukocytes can also be recruited to the injured site.




Permeability governs exudate composition

A greater increase in permeability permits larger molecules, including fibrinogen, to escape.

This distinguishes a fibrin-rich response from a relatively fluid-rich one.

Permeability alone does not determine the final distribution of deposits.




Fibrinogen: Essential plasma precursor

Fibrinogen enters the interstitial or serosal extracellular space from plasma.

Local coagulation reactions convert fibrinogen into insoluble fibrin.

The resulting strands accumulate where leakage and coagulation coincide.




Local procoagulant activity

Robbins specifies a local procoagulant stimulus alongside protein leakage.

Its role is to favor fibrin formation at the inflamed site.

Specific triggers vary with the underlying disease; Do not infer etiology from fibrin alone.




Integrated mechanism of exudation

Inflammatory mediators act on local small vessels.

The resulting permeability changes enable efflux of plasma proteins.

With a procoagulant stimulus, fibrin forms at the site of injury.




Histamine and vascular leakage

Histamine is a principal vasoactive mediator in acute inflammation.

It contributes to increased permeability and vasodilation.

The mediator response facilitates exudation but is not uniquely specific for fibrinous inflammation.



Other permeability-associated mediators

Serotonin, C3a/C5a-associated vasoactive amine release, and cysteinyl leukotrienes participate in permeability responses.

Multiple mediators can overlap in function.


Do not assign one mediator as the unique cause of a fibrinous deposit.

Fibrin deposition is spatially localized

Leakage occurs near inflamed microvessels.

Fibrin can accumulate along adjacent serosal or injured extracellular surfaces.

The pathologic surface pattern reflects both permeability and local coagulation.




Exudate is not synonymous with effusion

Exudate refers to inflammatory extravascular material rich in plasma constituents.

An effusion denotes accumulated fluid in a body cavity.

Fibrin may coat the serosa with or without a prominent fluid component.




A fibrinous response may contain fluid

Serofibrinous exudate combines serous fluid with fibrin.

The proportions of fluid and fibrin differ between cases.

This is particularly relevant when describing pericarditis and pleuritis.



Leukocytes accompany, but do not define, the pattern


Acute inflammation commonly recruits neutrophils.

In fibrinous inflammation, the defining finding is fibrin in the exudate.

A neutrophil-rich liquefied exudate instead suggests suppurative inflammation.

Illustrative H&E appearance

Eosinophilic fibrin meshwork



Inflamed underlying serosa


Schematic—not authentic histology

Mechanistic summary: Three linked compartments

Blood supplies fibrinogen.

Leaky microvessels permit escape into extracellular space.

Locally formed fibrin becomes a surface-deposited network.



Why these steps matter diagnostically


An eosinophilic deposit should be localized relative to the affected surface.

The reaction may include edema and inflammatory infiltrates beneath the deposit.

Tissue context is needed before assigning a specific cause.

Illustrative H&E appearance

Eosinophilic fibrin meshwork



Inflamed underlying serosa


Schematic—not authentic histology

Gross appearance: An altered serosal surface

A normally smooth, glistening serosa becomes coated with fibrinous material.

Surface deposits are often rough or shaggy.

Pericarditis is the classic teaching example.




Fibrinous pericarditis: Gross recognition

Look for deposits over the pericardial surface.

The important observation is the overlying fibrin, not simply the presence of pericardial fluid.

Extent and distribution may differ with the underlying insult.



Microscopy: Eosinophilic meshwork


Fibrin stains pink/eosinophilic on H&E.

A network of strands can coat the serosa.

Identify the boundary between surface deposit and underlying tissue.

Illustrative H&E appearance

Eosinophilic fibrin meshwork



Inflamed underlying serosa

Schematic—not authentic histology

Microscopy: Amorphous fibrinous coagulum


Fibrin may also appear as a more homogeneous eosinophilic coagulum.

The same basic composition may display different microscopic textures.

Recognize both patterns without over-interpreting morphology alone.

Illustrative H&E appearance

Amorphous fibrinous coagulum



Underlying inflamed serosa

Schematic—not authentic histology

Map a fibrin deposit onto a tissue section


Start by locating the intact serosal lining or organ surface.

Identify the overlying pink strands or coagulum.

Evaluate inflammation in the subjacent tissue separately.

Illustrative H&E appearance

Eosinophilic fibrin meshwork



Inflamed underlying serosa


Schematic—not authentic histology

Surface deposits are not tissue necrosis

Fibrin accumulates on the extracellular surface.

Necrotic parenchyma is a distinct finding requiring separate assessment.

A fibrinous coating alone does not prove extensive underlying necrosis.



Fibrinous versus serous inflammation

The distinction is the dominant exudative material—not a unique disease.

Feature	Serous pattern	Fibrinous pattern
Dominant material	Relatively fluid-rich effusion	Fibrin deposited extracellularly
Typical appearance	Clearer fluid accumulation	Pink strands / surface coagulum
Mechanistic clue	Permeability or mesothelial secretion	Large protein escape + local coagulation
Teaching example	Skin blister / serous cavity	Pericarditis / pleuritis

Fibrinous versus purulent inflammation

A mixed exudate may occur; Describe the dominant morphology.


Feature	Fibrinous	Purulent / suppurative
Dominant constituent	Extracellular fibrin network	Neutrophils + liquefied necrotic debris
Common morphology	Surface strands / coagulum	Pus; may collect as an abscess
Pathophysiology	Coagulation after protein leak	Often pyogenic bacterial injury
Interpretive warning	Fibrin does not establish infection	Pus does not exclude accompanying fibrin

Do not mistake fibrin for an abscess

An abscess is a localized collection of pus within a tissue or confined space.

Its core contains liquefied debris and degenerating inflammatory cells.

Fibrinous deposits instead form strands or coagulum on a surface.




Pericardium: An anatomic substrate

The visceral and parietal layers bound a potential space.

Fibrin may coat the opposing inflamed surfaces.

Subsequent organization can thicken layers or bridge the space.




A compact gross-to-microscopic correlation

Gross: Dull, roughened, fibrin-coated surface.

Microscopic: Eosinophilic strands / amorphous coagulum.

Mechanism: Protein leakage followed by local fibrin formation.




Fibrinous and serofibrinous pericarditis

These as common forms of pericardial inflammation.

Serous fluid may be admixed with varying amounts of fibrinous exudate.

The terms describe morphology, not a single etiology.



Common settings for fibrinous pericarditis

Myocardial infarction and postinfarction Dressler syndrome.

Uremia and autoimmune or rheumatic disease.


Trauma, cardiac surgery and other insults.

Myocardial infarction as a clinical context

Acute myocardial infarction is a recognized cause.

The pericardial response can include fibrin deposition.

Document the myocardial injury separately from the serosal morphology.



Dressler syndrome: Postinfarction setting

A delayed autoimmune response after myocardial infarction.

It belongs among the causes of fibrinous or serofibrinous pericarditis.

The temporal relationship and associated findings help establish etiology.




Stages describe pathologic progression, not a fixed clinical timeline

Uremia as a pericardial trigger

Uremia is among the recognized causes of fibrinous or serofibrinous pericarditis.

The morphologic reaction remains fibrin-rich despite a noninfectious initiating condition.

A fibrinous surface should not automatically be labeled bacterial.



Immune and rheumatic associations

Systemic lupus erythematosus and rheumatic fever are cited causes.

Pericardial exudation may occur as part of wider inflammatory disease.


The patient's clinical context distinguishes these etiologies.

Postoperative and traumatic reactions

Routine cardiac surgery can be followed by a fibrinous reaction.

Trauma may likewise involve the pericardium.

The timing and procedural history are critical to interpretation.




Symptoms of fibrinous pericarditis

Pain is often sharp, pleuritic and position dependent.

Fever may be present.

Symptoms indicate pericardial inflammation but do not independently establish its morphologic subtype.




Pericardial friction rub: Structural explanation

Roughened fibrin-coated layers can produce an audible friction rub.

The rub is a notable clinical clue to pericardial inflammation.

This links a surface deposit to a bedside physical sign.




Why a friction rub can disappear

A sufficiently large fluid effusion separates the opposing pericardial layers.

Separation can reduce or abolish the friction rub despite ongoing inflammation.

Absence of the rub therefore does not exclude pericardial disease.

Fluid separates opposing pericardial layers



Less surface contact

Distinguish fibrinous from purulent pericarditis

The two reactions differ principally in their dominant exudative material.


Feature	Fibrinous / serofibrinous	Purulent
Exudate	Fluid mixed with fibrin	Cloudy fluid to thick pus
Common context	MI, uremia, SLE, trauma	Active microbial invasion
Inflammatory burden	Variable	Often severe, systemic manifestations
Diagnosis	Context + morphology	Evidence of infection important

Pleural fibrinous reaction

Serous, serofibrinous and fibrinous pleuritis form a spectrum.

The pattern is affected by reaction intensity and duration.

The underlying pulmonary or systemic disorder remains central.




Pulmonary disorders associated with pleuritis

Pneumonia, tuberculosis, pulmonary infarction and lung abscess.

Bronchiectasis may also cause an inflammatory pleural reaction.

Inflammation in the underlying lung can extend to or irritate the pleura.



Systemic causes of serofibrinous pleuritis

Rheumatoid arthritis, systemic lupus erythematosus and uremia.

Diffuse infection, pleural metastases and mediastinal or pulmonary radiotherapy.


The pattern requires etiologic correlation.

Fluid and fibrin affect pleura differently

The fluid component may be resorbed.

The fibrin component may resolve or undergo organization.

Larger effusions can compress lung tissue and cause respiratory distress.




Possible sites beyond pleura and pericardium

The general inflammation chapter also mentions meningeal surfaces.

The fibrinous designation describes material deposited at the surface.

Identification of meningeal disease etiology requires its own assessment.




Two divergent fates of the fibrinous exudate

Successful removal: Fibrinolysis with macrophage clearance.

Incomplete removal: Fibroblast and vascular ingrowth.

The balance determines resolution versus fibrous organization.




Fibrinolysis and macrophage clearance

Fibrin can be dissolved by fibrinolysis.

Macrophages help clear the resulting material.

The affected surface may return toward normal when injury is limited.




What is meant by organization?

Persistent exudate becomes invaded by fibroblasts and blood vessels.

This fibrovascular tissue can mature into scar.

Organization converts a reversible surface deposit into a structural alteration.




Pericardial organization: Morphologic result

Opaque fibrous thickening can involve the pericardium and epicardium.

Fibrous adhesions may form between the serosal layers.

More extensive fibrosis can obliterate the pericardial space.




When adhesions impair function

Thin adhesions may have little functional effect.

More substantial fibrous changes can affect cardiac movement.


Extensive constricting scar can interfere with diastolic expansion.

Thin serosa



Potential space retained

Organized fibrosis




Scar may restrict filling

Constrictive pericarditis: A downstream risk

A dense fibrous or fibrocalcific pericardial scar limits diastolic filling.

Reduced capacity to increase cardiac output may result.

Not every acute fibrinous episode progresses to constriction.



Resolution, organization and chronicity

Outcomes depend on injury, tissue context and effectiveness of clearance.

Path	Mechanism	Potential result
Resolution	Fibrin removed and edema resorbed	Recovery of serosal surface
Organization	Fibrovascular growth into persistent fibrin	Fibrous thickening / adhesions
Chronic inflammation	Persistent stimulus or failed termination	Continuing injury and repair

Pattern recognition: Three discriminating findings


Protein-rich fibrin deposition rather than free fluid alone.

Grossly visible fibrin coating or microscopically eosinophilic strands.

An inflamed serosal or extracellular surface beneath the deposit.

Illustrative H&E appearance

Eosinophilic fibrin meshwork



Inflamed underlying serosa


Schematic—not authentic histology

Clinical vignette: Chest pain after infarction

Illustrative case: A patient develops sharp positional chest pain after myocardial infarction.

An audible friction rub is detected.

Question: Which exudative pattern best matches the findings?




Case interpretation: Fibrinous pericarditis

The infarction provides an established clinical context.

Surface fibrin can roughen opposing pericardial layers, accounting for the rub.

Clinical features alone do not replace morphologic examination when needed.




Clinical vignette: Pleural reaction

Illustrative case: A patient with pneumonia develops a pleural inflammatory effusion.

A fibrin coating is documented on the pleural surface.

Question: What are the two possible fates of this deposited fibrin?




Case interpretation: Clearance or organization

Fibrinolysis and macrophages can clear the deposit.

Persistent exudate can be replaced by fibrous tissue.

Assess the disease causing the pleuritis and the extent of injury.



Microscopic spotter: Separate deposit from tissue


Locate the organ surface and the overlying eosinophilic material.

Describe strands versus amorphous coagulum.

Report accompanying cellular inflammation without mislabeling the deposit as pus.

Illustrative H&E appearance

Eosinophilic fibrin meshwork



Inflamed underlying serosa

Schematic—not authentic histology

Final synthesis: Six essential points

Fibrin requires leakage of large proteins plus local coagulation.

Serosal surfaces are characteristic sites.

On H&E, fibrin appears eosinophilic and often threadlike.

Pericarditis links morphology to a friction rub.

Clearance permits resolution; Persistence may organize.

The pattern alone never establishes the underlying cause.

Core reference and further context

Robbins, Cotran & Kumar: Pathologic Basis of Disease, 11th ed. (2026).

Chapter 3, “Morphologic Patterns of Acute Inflammation,” pp. 83–85.

Additional textbook context: “Pericardial Disease” and “Inflammatory Pleural Effusions.”