

Giardia lamblia

Protozoal enteritis: pathogenesis, morphology and diagnosis

A clinicopathologic lecture

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning outcomes

- Trace Giardia transmission and its intestinal life cycle.
- Explain epithelial injury without tissue invasion.
- Recognize biopsy morphology and relevant stool tests.
- Connect immune status with persistent or severe infection.

A flagellated protozoan of the small intestine

- Giardia is a flagellated protozoan associated with human enteric infection.
- The parasite occupies the small-intestinal lumen and villous surface.
- Clinical disease ranges from silent carriage to persistent malabsorption.

Core distinction

A noninvasive parasite can cause consequential epithelial dysfunction.

Three names for the same organism

Accepted name in sources	Interpretation
Giardia lamblia	Name used in the Robbins topic heading
Giardia duodenalis	Alternative name used by Robbins and CDC
Giardia intestinalis	Additional synonym cited by both sources

Infection is noninvasive—not harmless

Where organisms are found

- Luminal small-bowel surface
- Closely apposed to villous enterocytes
- No tissue invasion in the Robbins description

Why patients become symptomatic

- Brush-border enzyme reduction
- Microvillous damage and epithelial apoptosis
- Malabsorption and diarrhea

Two functional stages

Cyst

Environmentally stable infectious form; Ingested with contaminated material




Trophozoite

Motile luminal form; Attaches to small-intestinal surface

Recognizing the infective cyst


- The cyst is the environmentally hardy form.
- Mature cysts have four nuclei on microscopy (CDC).
- Infection follows ingestion of contaminating cysts.



Schematic mature cyst

Recognizing the trophozoite

- Trophozoites are pear shaped.
- Two similarly sized nuclei are a key morphologic clue.
- Flagella and an attachment disk are demonstrable (CDC).



Cyst and trophozoite are not interchangeable

Characteristic	Cyst	Trophozoite
Main function	Survival and transmission	Colonization of intestinal lumen
Shape	Oval/ellipsoid	Pear shaped
Nuclei	Up to four when mature	Two similarly sized
Detection	Common in formed stool	May appear in stool or duodenum

Fecal–oral transmission

Fecal material

Giardia cysts contaminate water,
food or hands



Accidental ingestion

A small inoculum can establish
infection



Intestinal infection

Trophozoites establish infection
in small bowel

Environmental exposure explains many infections

- Fecally contaminated drinking water is a principal route.
- Unfiltered public water supplies may transmit Giardia.
- Rural streams are a recognized exposure among campers.
- Contaminated food provides an additional route.

A very small inoculum can infect

- Ingestion of as few as 10 cysts may cause infection.
- A low inoculum helps explain infection after unintended ingestion.
- A low dose does not quantify risk in an individual exposure.

Interpretation

A minimal infective dose is not an individual attack probability.

Cyst survival and water disinfection

- Giardia cysts persist in the environment.
- Their chlorine resistance permits transmission in treated but unfiltered settings.
- Filtration and avoidance of contaminated water are important prevention concepts.

Two common water-related scenarios

Drinking-water exposure

- Unfiltered public supplies
- Rural streams used by campers
- Ingestion is the causal event


Recreational-water exposure

- Accidental swallowing during swimming
- Contaminated water carries cysts
- External skin contact alone is not the stated route

Food and direct fecal–oral spread

- Fecally contaminated food can transmit cysts.
- Fecal–oral transfer may occur outside a shared water source.
- The hardy cyst stage permits environmental persistence.

Environmental cycle and human infection



Excystation begins in the small bowel

Cyst ingestion

Contaminated material is
swallowed



Excystation

In the small intestine, each cyst
releases trophozoites



Surface colonization

Motile trophozoites remain in
the lumen

Trophozoite multiplication occurs in lumen

Free trophozoite

Organism resides in proximal small bowel



Longitudinal fission

Trophozoites multiply asexually



Mucosal contact

Organisms may attach to epithelial surface

Encystation permits onward transmission

Luminal movement

Parasites move distally toward colon



Encystation

Protective cyst form is produced




Stool shedding

Cysts in feces can infect a new host

Trophozoites remain on the villous surface

- Giardia trophozoites may be closely attached to enterocytes.
- They are located in the lumen, not within the lamina propria.
- This topography matters when reading duodenal biopsy sections.



Surface attachment is a pathologic interface

- Close contact occurs at the brush border of villous enterocytes.
- CDC describes the ventral adhesive disk involved in attachment.
- The tissue response can be disproportionate to visible distortion.

Noninvasive Giardia versus mucosal invasion

Giardia

- Trophozoites on luminal surface
- Brush-border dysfunction
- No invasive trophozoites in mucosa (Robbins)

Invasive amebiasis

- Entamoeba breaches colonic mucosa
- Invasion causes tissue injury and ulcers
- Can spread beyond bowel

Three epithelial effects explain disease

- Giardia causes injury and dysfunction even without invasion.
- These mechanisms converge on absorptive failure.

Brush-border enzyme reduction

Microvillous damage

Epithelial-cell apoptosis

Reduced brush-border enzymes impair digestion

- Robbins documents decreased expression of brush-border enzymes.
- Terminal carbohydrate and peptide digestion normally depends on these enzymes.
- Enzyme loss provides a mechanistic link to malabsorption.


Microvillous injury reduces surface function

- Giardia can damage enterocyte microvilli.
- Microvilli support epithelial digestion and absorption.
- A reduced functional surface can cause symptoms without marked villous atrophy.

Enterocyte apoptosis contributes to injury

- Apoptosis of small-intestinal epithelial cells is part of the described response.
- Epithelial loss can worsen brush-border dysfunction.
- Microscopic severity is not necessarily proportional to organism numbers.

How Giardia produces malabsorptive diarrhea



Connect intestinal physiology to Giardia

Absorptive phase	Relevance in Giardia
Intraluminal digestion	Not the primary lesion described by Robbins
Terminal digestion	Brush-border enzyme expression decreases
Transepithelial transport	Enterocyte injury may compromise function
Lymphatic lipid transport	Not a primary established lesion in this excerpt

Malabsorptive stool is a mechanism-based concept

- Malabsorption entails defective uptake of nutrients and water.
- Steatorrhea may accompany generalized malabsorptive disease.
- Giardia is a protozoal cause of intestinal malabsorption; Not every case has steatorrhea.


Secretory IgA contributes to clearance

- Secretory IgA as an important host defense.
- Mucosal antibody can promote immune control of intestinal Giardia.
- Impaired antibody production is linked to severe infection.

Mucosal IL-6 is another clearance signal

- IL-6 responses in the intestinal mucosa contribute to clearance.
- The source identifies a protective association—not a validated biomarker threshold.
- Clinical assessment does not depend on measuring mucosal IL-6.

Antigenic variation supports persistence



Intermittent symptoms fit chronic persistence

- Surface antigen variation permits immune evasion.
- The parasite may persist for months or years.
- The resulting illness may fluctuate rather than remain continuous.

Host factors modify infection severity

Host context	Source-supported consequence
Immunosuppression	Often severe disease
Agammaglobulinemia	Often severely affected
Malnutrition	Often severely affected
Infants / young children	Greater risk of severe illness
Travelers	Greater risk of severe illness

Agammaglobulinemia is a mechanistic clue


- Antibody-deficient patients can develop particularly severe giardiasis.
- This accords with the importance of secretory IgA in clearance.
- Chronic Giardia should remain a differential in relevant enteric illness.

Malnutrition and Giardia can coexist

- Malnourished individuals are often severely affected by Giardia.
- Giardia itself can produce malabsorption and weight loss.
- Assess both vulnerability before infection and nutritional consequences after it.


Biopsy shows luminal organisms—not tissue invasion

- Organisms may be numerous near villous tips.
- They are often closely apposed to the brush border.
- Examine the intestinal lumen and epithelial surface carefully.



Two equal nuclei and pear shape are diagnostic clues

- On duodenal biopsy, identify the characteristic pear-like trophozoite.
- Look for two similarly sized nuclei where section orientation permits.
- Organisms are small and easily overlooked in luminal material.



Small-bowel architecture may appear normal

- Large numbers of Giardia may coexist with near-normal villi.
- Absence of striking villous atrophy does not exclude giardiasis.
- Search the luminal border rather than relying only on mucosal damage.

Heavy infection may blunt villi

- Villous blunting can accompany intense Giardia infection.
- Blunting may reduce effective absorptive architecture.
- Its absence is compatible with infection.

Intraepithelial lymphocytes may increase

- Heavy infection can show increased intraepithelial lymphocytes.
- This is a host epithelial immune response, not parasite invasion.
- The finding is not specific for Giardia on its own.

The lamina propria may be inflamed

- A mixed inflammatory infiltrate in lamina propria.
- A mild or modest reaction can coexist with surface organisms.
- Do not diagnose Giardia from inflammation without identifying the parasite or a relevant test.

Morphology spans minimal to reactive mucosa

Observation	Interpretation
Nearly normal villi	Does not exclude Giardia
Luminal pear-shaped trophozoites	Direct morphologic evidence of organism
Villous blunting	May accompany heavier infection
Increased intraepithelial lymphocytes	Possible associated epithelial response
Mixed lamina propria infiltrate	Possible nonspecific mucosal reaction


Giardia and Entamoeba: A useful diagnostic contrast

Feature	Giardia	Entamoeba histolytica
Typical tissue location	Small-intestinal lumen	Colon and invaded mucosa
Invasion	Absent	Present
Characteristic clue	Pear form; two nuclei	May ingest erythrocytes
Tissue injury	Brush-border dysfunction	Ulcers and tissue destruction

Giardia versus Cryptosporidium on biopsy

Feature	Giardia	Cryptosporidium
Position	Luminal villous surface	At apical epithelial membrane
Appearance	Pear-shaped trophozoites	Tiny round blue organisms
Host interaction	Noninvasive lumen-contact	Parasite enclosed by host cell cytoplasm
Diagnosis in stool	Giardia cyst detection	Cryptosporidium oocysts

Clinical manifestations form a continuum



Giardiasis may be clinically silent

- Robbins explicitly describes subclinical infection.
- Symptom absence does not negate the possibility of intestinal carriage.
- Interpret stool findings in their clinical and exposure context.

Acute diarrhea is a common presentation

- Symptomatic Giardia can present with acute diarrhea.
- The disease mechanism emphasizes functional injury to small-bowel epithelium.
- Exposure history may make a parasitic cause more plausible.

Chronic diarrhea can be intermittent

- Persistent Giardia infection can extend for months or years.
- Variable antigen expression contributes to persistence.
- Ongoing or episodic symptoms may merit parasitic evaluation.

Malabsorption is a central consequence

- Reduced brush-border enzymes compromise terminal digestion.
- Microvillous damage undermines absorption.
- Symptoms arise from functional changes even with minimal structural abnormalities.


Weight loss integrates duration and dysfunction

- Giardia may cause clinically important weight loss.
- Chronic diarrhea and impaired nutrient absorption provide a plausible mechanism.
- Nutritional status matters especially in children and malnourished hosts.

Clinicopathologic correlation

Clinical feature	Corresponding pathologic explanation
Diarrhea	Impaired mucosal digestion and absorption
Malabsorption	Reduced enzyme expression and microvillous damage
Weight loss	Sustained impaired nutrient uptake
Intermittent illness	Long-term parasite persistence with antigen variation
Subclinical infection	Mucosal contact need not produce prominent symptoms

Select the diagnostic sample thoughtfully



Stool immunofluorescence is emphasized by Robbins

- Stool cyst immunofluorescence for documentation.
- Immunofluorescent antibodies permit visual detection of cysts.
- Positive tests must be interpreted with clinical context.

Additional laboratory methods documented by CDC

Method	What the method detects
Wet mount / concentration	Stool cysts or trophozoites
Permanent stained smear	Morphologic trophozoites
Antigen enzyme immunoassay	Giardia antigens in stool
Direct immunofluorescence	Labeled parasite cysts in stool
Duodenal sample or biopsy	Trophozoites when sampled

Intermittent detection can require repeat stool testing

- CDC notes that repeated fecal specimens may be necessary.
- A single negative sample does not invariably exclude infection.
- Test choice and repeat sampling follow the clinical scenario.

Duodenal biopsy: What the pathologist should inspect

- Find organisms between villi and along the luminal brush border.
- Confirm pear shape and paired nuclei where visible.
- Recognize that mucosa may remain nearly normal.

Distinguish the diagnostic roles of stool and biopsy

Dimension	Stool assessment	Duodenal histology
Primary finding	Cysts / trophozoites / antigen	Trophozoites near villi
Source emphasis	Immunofluorescent cyst detection	Pear shape, two nuclei
Main limitation	Detection may vary between samples	Organisms easily overlooked

Treatment is effective, but recurrence occurs

- Oral antimicrobial therapy is effective.
- The specific drug regimen is outside the supplied Giardia excerpt.
- Recurrent symptoms can occur following therapy.

Recurrence is a clinical follow-up problem

- The textbook explicitly states that recurrence is common.
- Consider ongoing exposure, persistent infection or other causes when symptoms return.
- Repeat assessment should be informed by the clinical context.

Prevention follows the transmission mechanism

- Avoid ingestion of untreated or contaminated water.
- Do not assume chlorination alone neutralizes Giardia cysts.
- Reduce fecal contamination of food, water and hands.

Case A — diarrhea after a hiking trip

Illustrative case (fictional)

An adult develops diarrhea and weight loss after repeatedly drinking untreated stream water during a camping trip.

Question

Which exposure and intestinal pathogenesis make Giardia plausible?

Case A — mechanism-based interpretation

- Stream water is a documented route of cyst exposure.
- A low infectious inoculum makes accidental ingestion relevant.
- Enzyme loss and microvillous injury explain malabsorptive symptoms.

Case B — normal villi with suspicious luminal profiles

Illustrative case (fictional)

A duodenal biopsy shows near-normal villi. Small pear-shaped structures are seen in the lumen adjacent to the brush border.

Question

What structures should the pathologist look for to support Giardia?

Case B — why normal villi can mislead

- Giardia over almost normal-appearing villi.
- Parasites can be numerous despite limited architectural distortion.
- The luminal surface deserves deliberate high-power inspection.

Case C — persistent symptoms and antibody deficiency

Illustrative case (fictional)

A person with markedly impaired humoral immunity has intermittent chronic diarrhea and weight loss. Giardia infection is being considered.

Question

Which host-defense axis and parasite strategy are relevant?


Case C — combine host and pathogen factors

- Secretory IgA is important for immune clearance.
- Agammaglobulinemic hosts may develop severe disease.
- Variant surface antigen switching favors persistence.

Avoid four common diagnostic overstatements

Overstatement	Source-grounded correction
“Normal villi exclude Giardia.”	Near-normal mucosa is possible.
“Giardia invades the mucosa.”	Robbins describes no invasion.
“Trophozoites have four nuclei.”	Trophozoites have two; mature cysts may have four (CDC).
“Chlorine makes all water safe.”	Cysts are chlorine resistant.

Synthesis: One coherent pathologic model



Reference sources

Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th edition. 2026.

CDC DPDx: Giardiasis. Last reviewed June 5, 2024. cdc.gov/dpdx/giardiasis/

Histologic figures have not been reproduced; organism and mucosal drawings are original editable schematics.