

Glaucomatous Optic Nerve Damage

Mechanisms, morphology, and clinicopathologic interpretation


Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning objectives

- Link aqueous outflow abnormalities to elevated intraocular pressure.
- Explain why pressure and optic neuropathy are not interchangeable.
- Recognize retinal and optic-nerve morphology, including age-dependent sequelae.
- Distinguish glaucomatous changes from selected optic neuropathies.



Glaucoma is defined by characteristic injury

- Distinctive optic nerve cup and visual-field changes.
- Intraocular pressure is usually elevated, but not invariably.
- The damage phenotype is more specific than a pressure reading alone.




The optic nerve is a central nervous system tract

- The optic nerve consists of neuronal axons carrying visual information.
- Meningeal coverings surround it, with cerebrospinal fluid around the nerve.
- Its pathology therefore shares important features with central nervous system injury.

Schematic • optic nerve and meningeal sheath

Dura / sheath

**Optic nerve
Axons**




Optic axons form a sensory tract invested by meninges.

Ganglion cells connect retina to optic nerve


- Retinal ganglion-cell bodies are within the inner retina.
- Their axons form the retinal nerve-fiber layer.
- Axons converge at the optic disc and populate the optic nerve.

Schematic • selective inner retinal loss



Structural loss produces functional consequences

- Loss of ganglion cells removes retinal output neurons.
- Loss of their axons reduces tissue within the nerve-fiber layer and optic nerve.
- Glaucoma is associated with characteristic visual-field changes.



Pressure is a major contributor, not a definition

Intraocular pressure


- Important exposure
- Often elevated in glaucoma
- Not sufficient alone

Neuronal injury

- Ganglion-cell / axon loss
- Cupping + field change
- Variable susceptibility


Pressure-related damage has a common endpoint

- Distinct outflow defects can elevate intraocular pressure.
- Prolonged exposure is linked with retinal ganglion-cell and optic-nerve loss.
- Different initiating mechanisms can converge on the same morphologic phenotype.




Aqueous humor is produced in the ciliary body

- Aqueous humor originates in the ciliary body.
- It first enters the posterior chamber behind the iris.
- Production continues even when drainage is obstructed.




Aqueous humor moves through the pupil

- Fluid passes from posterior chamber through the pupillary aperture.
- It subsequently occupies the anterior chamber.
- The pupillary passage can become a point of obstruction in predisposed eyes.




The trabecular route is the major outflow path

- Most aqueous humor exits through the trabecular meshwork.
- The meshwork lies at the iridocorneal angle.
- Aqueous passes toward Schlemm canal; Minor routes also exist.



Aqueous outflow obstruction elevates pressure

- Resistance within the major drainage route impairs egress.
- Ongoing aqueous formation sustains a pressure load when outflow falls.
- The obstruction site distinguishes major glaucoma mechanisms.



Open-angle and angle-closure are mechanistic classes

Open-angle

- Angle anatomically accessible
- Outflow resistance increased
- Primary / secondary


Angle-closure

- Iris obstructs angle
- Physical egress block
- Primary / secondary

Open-angle glaucoma retains angle access

- Aqueous humor can reach the trabecular meshwork.
- The major impediment is increased resistance in the open drainage angle.
- This distinguishes resistance from anatomic closure of the angle.

Schematic • iridocorneal drainage angle




Aqueous can approach the meshwork

Primary open-angle disease may look subtle anteriorly

- The anterior-chamber angle remains open.
- Few structural changes may be apparent in the anterior segment.
- The precise cause of increased outflow resistance is unknown.


Schematic • iridocorneal drainage angle



Aqueous can approach the meshwork


Genetic associations do not equal one mechanism

- Genetic factors are suspected in primary open-angle glaucoma.
- The source reports genome-wide associations involving 40 genes.
- The findings do not establish one universal mechanism for impaired aqueous egress.




Pseudoexfoliation can obstruct open-angle drainage

- Fibrillar material is deposited throughout the anterior segment.
- The condition is described as the commonest secondary open-angle glaucoma type.
- Deposits may also occur around connective-tissue vessels and in other organs.




Lens proteins may clog the trabecular meshwork

- Dissolution of lens material releases high-molecular-weight proteins.
- The resulting particulate load may impede trabecular drainage.
- This source-defined mechanism is called phacolytic glaucoma.




Senescent erythrocytes can obstruct the angle

- Following ocular trauma, senescent red cells may remain in the eye.
- These cells can block the trabecular outflow apparatus despite an open angle.
- The named entity is ghost-cell glaucoma.




Iris pigment can become outflow debris

- Pigment granules arise from the iris pigment epithelium.
- Pigment can accumulate within trabecular drainage structures.
- The resulting secondary open-angle process is pigmentary glaucoma.




Tumor debris can impair open-angle egress

- Necrotic tumor material may enter the aqueous compartment.
- This particulate load can obstruct trabecular drainage.
- The text names melanolytic glaucoma as one example.



Elevated episcleral venous pressure impedes drainage

- Increased pressure in the episcleral venous system can elevate intraocular pressure.
- The iridocorneal angle remains anatomically open.
- Examples include Sturge–Weber vascular malformations and carotid–cavernous fistula.



Secondary open-angle causes have different substrates

Trabecular obstruction

- Pseudoexfoliation material
- Lens proteins / ghost cells
- Pigment / tumor debris


Downstream pressure

- High episcleral venous pressure
- Vascular malformations
- Carotid–cavernous fistula

Angle closure physically blocks the main exit

- The peripheral iris obstructs the trabecular meshwork.
- Aqueous cannot freely access the major outflow route.
- The distinction from open-angle glaucoma is anatomical, not merely pressure magnitude.


Schematic • iridocorneal drainage angle



Shallow chambers predispose to primary angle closure


- Primary angle closure commonly develops in anatomically shallow anterior chambers.
- Hyperopia is a frequent associated ocular configuration.
- Lens–iris apposition at the pupil can initiate the obstructive sequence.

Schematic • iridocorneal drainage angle



Pupillary block prevents posterior-to-anterior flow


- Iris contact with the anterior lens surface may seal the pupillary passage.
- Aqueous accumulates behind the iris while production continues.
- Posterior-chamber pressure then increases.



Iris bombé converts block into angle closure


- Pressure behind the iris bows the iris forward.
- The bowed peripheral iris approaches and occludes the trabecular meshwork.
- A major aqueous outflow pathway is thereby closed.

Schematic • iridocorneal drainage angle




Primary closure is a sequence, not one event

- Predisposed chamber geometry permits iris–lens apposition.
- Pupillary block raises posterior-chamber pressure.
- Iris bombé occludes the angle and produces marked pressure elevation.




Sustained pressure elevation can injure the cornea

- Corneal edema may follow persistent severe intraocular pressure elevation.
- Bullous keratopathy can develop despite healthy endothelial-cell numbers.
- Corneal consequences coexist with the potential for optic nerve damage.



The lens epithelium also responds to severe closure

- The avascular lens relies on aqueous humor for nutritional support.
- Unremitting pressure elevation can damage lens epithelium.
- Minute anterior subcapsular opacities may be visible by slit lamp.



Secondary angle closure involves other barriers

Contractile membrane


- New vessels + stroma
- Myofibroblast contraction
- Peripheral anterior synechiae

Mechanical displacement

- Ciliary-body tumor
- Iris pushed forward
- Angle blocked

Retinal ischemia can initiate neovascular glaucoma


- Chronic retinal ischemia upregulates VEGF and other angiogenic signals.
- Proangiogenic factors reach the aqueous compartment.
- Thin fibrovascular membranes can develop on the iris surface.



A neovascular membrane alters the iris surface


- The neovascular membrane may be clinically transparent.
- It can smooth the usual folds and crypts of the iris.
- The important pathologic behavior is subsequent contraction, not merely its presence.

Schematic • iridocorneal drainage angle



Myofibroblast contraction seals the angle

- Myofibroblasts within the fibrovascular membrane generate contractile force.
- Contraction draws peripheral iris toward trabecular structures.
- Peripheral anterior synechiae can obstruct outflow.



Tumors can contribute to neovascular or direct closure

Neovascular route


- Tumor-associated angiogenesis
- Fibrovascular iris membrane
- Contractile closure

Direct mechanical route

- Mass near ciliary body
- Iris displaced into angle
- Outflow occlusion

Different drainage lesions converge on the same risk

- Primary open-angle: Unexplained resistance despite angle access.
- Secondary open-angle: Debris or venous-pressure effects.
- Angle-closure: Iris obstructs the meshwork directly or through synechiae.



Neuronal susceptibility modifies the pressure response

- People differ in how vulnerable their optic nerve axons are to pressure.
- The source documents injury at normal pressure and tolerance of elevated pressure.
- These contrasting observations imply a spectrum of susceptibility.

Susceptibility to pressure exposure

High
susceptibility

Intermediate

Lower
susceptibility

Same exposure ≠ same neuronal outcome

Normal-tension glaucoma preserves the injury pattern

Pressure status

- Within normal range
- Not necessarily increased
- One component of evaluation

Injury phenotype

- Characteristic cupping
- Visual-field changes
- Still glaucomatous

Elevated pressure may occur without optic neuropathy

Pressure finding


- Elevated intraocular pressure
- Prolonged exposure possible
- Risk factor

Damage evidence

- No characteristic cup change
- No characteristic field change
- Damage not demonstrated

Pressure and damage must be assessed on two axes


- A single pressure threshold cannot describe all susceptibility patterns.
- Structural cupping and retinal loss characterize established damage.
- The source supports a continuum rather than a simple pressure-only dichotomy.



Retinal ganglion-cell loss is a primary lesion

- Glaucoma characteristically produces diffuse loss of retinal ganglion cells.
- These neurons supply axons to the optic nerve.
- Their disappearance explains part of the associated nerve atrophy.

Schematic • selective inner retinal loss



Preserved retina


Glaucomatous loss

Ganglion cells are depleted; Schematic thickness only.


The retinal nerve-fiber layer becomes thin

- Ganglion-cell axons form the nerve-fiber layer.
- Loss of axons reduces the thickness of this inner retinal layer.
- This thinning as a characteristic morphologic finding.

Schematic • selective inner retinal loss



Preserved retina




Glaucomatous loss

Axons are depleted; Schematic thickness only.


Inner retinal layers show linked atrophy

- The nerve-fiber layer and ganglion-cell layer are depleted.
- The figure caption identifies the inner plexiform layer as an anatomical reference.
- Structural loss is most informative when aligned with the affected neuronal compartment.

Schematic • selective inner retinal loss



Preserved retina




Glaucomatous loss

Inner retina are depleted; Schematic thickness only.


Optical coherence tomography quantifies thinning

- The textbook explicitly identifies OCT as a method to measure nerve-fiber-layer thickness.
- An imaging measurement can track a structural consequence of neuronal loss.
- The scan is evidence about tissue structure, not a substitute for all clinical findings.

Retinal nerve-fiber-layer thickness



Intact layer (schematic)




Thinned layer (schematic)

OCT measures thickness; This diagram is not an OCT scan.


Optic-disc cupping reflects tissue loss

- The surface cup becomes enlarged or deepened as glaucoma progresses.
- The source links cupping in part to retinal ganglion-cell loss.
- Cupping describes a structural change at the optic nerve head.



Advanced glaucoma combines cupping with atrophy

- The optic nerve is atrophic in advanced chronic disease.
- The surface cup may become markedly widened and deepened.
- The combination of cupping and atrophy is particularly characteristic in this context.




A wide subdural space may signal severe nerve loss

- The optic nerve is surrounded by meningeal sheaths including dura.
- Marked axonal atrophy reduces the caliber of nerve tissue.
- In the source figure, this produces a conspicuously wide subdural space.

Schematic • optic nerve and meningeal sheath

Dura / sheath


**Optic nerve
Axons**

A schematic diagram illustrating the layers of the meningeal sheath surrounding the optic nerve. It consists of three concentric rounded rectangles. The innermost rectangle is dark blue and labeled 'Optic nerve Axons'. The middle rectangle is light blue and represents the 'Dura / sheath'. The outermost rectangle is white and represents the 'Dura / sheath'. The space between the inner and middle rectangles is labeled 'Optic nerve Axons', and the space between the middle and outer rectangles is labeled 'Dura / sheath'.

With advanced atrophy the nerve shrinks inside its meningeal sheath.


Three anatomical compartments tell one injury story

- Retina: Ganglion cells and nerve-fiber layer are depleted.
- Optic nerve head: Surface cupping becomes conspicuous.
- Retrobulbar nerve: Advanced atrophy leaves expanded surrounding space.



Pediatric pressure elevation changes eye size

- In infants and children, elevated pressure can enlarge the globe.
- Developing ocular coats deform differently from adult tissue.
- This age dependence is an important morphologic distinction.




Developing eye

Mature eye

Buphthalmos denotes diffuse globe enlargement

- Prolonged pressure elevation in early life may distend the entire eye.
- The textbook terms this appearance buphthalmos.
- It is a consequence of a developing eye subjected to pressure load.




Developing eye


Mature eye

Megalocornea is an associated pediatric change

- Corneal enlargement can occur with elevated childhood intraocular pressure.
- The source names megalocornea alongside buphthalmos.
- An enlarged cornea is a pressure-related structural sign, not a retinal lesion.




Developing eye



Mature eye

The adult globe may develop focal staphyloma

- After eye growth is complete, pressure can cause localized scleral thinning.
- Uveal tissue may line the resulting ectatic sclera.
- This configuration is called staphyloma.



Developing eye

Mature eye

Age changes the morphology of chronic pressure

Infant / child


- Buphthalmos
- Megalocornea
- Diffuse enlargement

Adult

- Focal scleral thinning
- Uveal-lined ectasia
- Staphyloma

Structure and function are complementary clues

- Visual-field abnormality establishes functional impairment characteristic of glaucoma.
- OCT thinning and disc cupping characterize different structural levels.
- Pressure measurement informs risk but does not replace structure–function correlation.



Papilledema must not be confused with cupping

Glaucoma

- Optic nerve cup deepens
- Chronic nerve atrophy
- Retinal ganglion-cell loss

Papilledema

- Disc swelling / hyperemia
- Venous stasis / transport block
- CSF-pressure or compression link

AION has a different injury mechanism

Glaucomatous neuropathy

- Often pressure-associated
- Chronic cupping / atrophy
- Retinal ganglion-cell loss

Anterior ischemic neuropathy

- Vascular supply interruption
- Acute pale disc swelling
- Infarction may be permanent

Inherited and toxic neuropathies broaden the differential

Glaucomatous pattern


- Characteristic cup change
- Retinal nerve-fiber loss
- Pressure often implicated

Other optic neuropathy

- Inherited / nutritional / toxic
- May damage macular fibers
- Alternative pathogenesis

Leber hereditary optic neuropathy is mitochondrial

- The source attributes Leber optic neuropathy to inherited mitochondrial gene mutations.
- It follows a maternal pattern and commonly affects young adults.
- This example illustrates optic nerve injury without a glaucoma-type outflow lesion.



Case: Normal pressure, abnormal nerve morphology

- An eye has characteristic cup enlargement and nerve-fiber thinning.
- Measured intraocular pressure is within the usual range.
- Interpretation: Normal-tension glaucoma remains possible; Pressure alone does not rule it out.

Teaching case • interpretation is illustrative, not patient data

Case: Sustained pressure, no documented injury

- An individual has long-observed elevated intraocular pressure.
- There is no characteristic field change or progressive cupping documented.
- Interpretation: High pressure alone is not proof of glaucomatous neuronal loss.

Teaching case • interpretation is illustrative, not patient data

Case: Retinal ischemia and iris membrane

- Chronic retinal ischemia is accompanied by an abnormal iris-surface membrane.
- The membrane contracts and the angle becomes closed.
- Interpretation: The source mechanism is secondary neovascular angle-closure glaucoma.

Teaching case • interpretation is illustrative, not patient data


Case: Trauma followed by an open-angle obstruction

- Senescent erythrocytes accumulate after ocular trauma.
- The angle remains anatomically open but aqueous outflow is impaired.
- Interpretation: Ghost-cell secondary open-angle glaucoma.

Teaching case • interpretation is illustrative, not patient data


Integrated morphologic diagnosis

- Look for the characteristic retinal ganglion-cell and axonal loss.
- Relate an enlarged optic nerve cup to the loss of tissue.
- Interpret pressure and drainage mechanisms as etiologic context, not replacements for injury evidence.



Final synthesis

- Aqueous outflow abnormalities commonly elevate intraocular pressure.
- Ganglion-cell depletion, nerve-fiber thinning and cupping define the chronic damage phenotype.
- Normal-tension injury and pressure tolerance demonstrate variable susceptibility.
- Age determines additional globe and corneal structural consequences.



Primary reference and study boundaries

- Robbins, Cotran & Kumar: Pathologic Basis of Disease, 11th edition (2026).
- Chapter 29, “The Eye”: glaucoma pp. 1195–1197; optic nerve pp. 1207–1209.
- Morphologic schematics in this lecture are original instructional diagrams, not specimen photographs.