

Grading and Staging of Tumors

Morphologic aggressiveness, anatomic extent, and clinical interpretation

A morphology-to-clinical-extent framework

Dr Abu Suraih Sakhri


MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Explain why tumor grade and stage answer different clinical questions.
- Identify the morphologic features used to assess differentiation.
- Interpret the general T, N, and M axes without inventing site-specific cutoffs.
- Recognize limitations of morphologic grading, sampling, and purely anatomic staging.

Why assign grade and stage?



Two complementary axes

GRADE — tumor phenotype

- How closely neoplastic cells resemble normal tissue
- Architectural, cytologic, and sometimes proliferative findings
- A morphologic estimate of aggressiveness

STAGE — disease extent

- Primary lesion and local spread
- Regional lymph-node involvement
- Distant metastatic dissemination

Biologic appearance is not anatomic spread

Histologic appearance ↓

Less extensive disease

More extensive disease

More differentiated

Localized, differentiated tumor

Disseminated, differentiated
tumor

Less differentiated

Localized, anaplastic tumor

Disseminated, anaplastic
tumor

Anatomic spread →

Two different risks

Localized, well-differentiated thyroid cancer

- Cells resemble thyroid tissue
- Anatomic extent remains confined
- Differentiation and spread both matter


Anaplastic thyroid cancer invading adjacent tissues

- Poor morphologic differentiation
- Extension into surrounding structures
- Clinical course cannot be inferred from a single feature


Grade describes tumor-cell differentiation

- Differentiation is resemblance to normal parenchymal cells.
- Resemblance is judged morphologically and sometimes functionally.
- Grading converts selected tissue features into a prognostic descriptor.
- The exact features and category boundaries depend on tumor type.

Differentiation forms a continuum




Well-differentiated tumors retain structure



Simplified schematic — not histologic evidence

- Relatively uniform cells preserve a recognizable tissue pattern.
- Some cancers form structures resembling their site of origin.
- Morphologic resemblance alone does not exclude malignancy.


Moderate differentiation preserves partial pattern



Simplified schematic — not histologic evidence

- Tumor still forms some recognizable structures.
- Architecture is more irregular than its normal counterpart.
- Invasion, not differentiation alone, establishes malignant behavior.

Poor differentiation obscures lineage morphology



Simplified schematic — not histologic evidence

- Normal architectural organization is largely lost.
- Cellular and nuclear atypia may be conspicuous.
- Precise tumor classification may require ancillary studies.

Anaplasia is loss of differentiation

DIFFERENTIATED

- Organized tissue-like structure
- Some specialized functions may remain
- Normal counterparts are recognizable


ANAPLASTIC

- Disordered appearance
- Numerous atypical cell features
- Tissue of origin can be difficult to recognize

Architecture can contribute to grade

- Assess gland formation or other tissue-appropriate structures.
- Ask whether the neoplasm respects normal cellular relationships.
- Architecture is one component—not a universal standalone grade.
- Each organ-specific grading system defines its own relevant patterns.

Pleomorphism reflects morphologic diversity



Simplified schematic — not histologic evidence

- Cells vary in size and shape within the same lesion.
- Tumor giant cells may be present in markedly anaplastic neoplasms.
- Pleomorphism contributes to atypia but is not itself a staging variable.

Abnormal nuclear features support anaplasia

- Nuclei may become disproportionately large.
- Nuclear outlines and chromatin distribution may be irregular.
- Hyperchromasia and conspicuous nucleoli may be present.
- Interpret nuclear features in tissue and tumor-type context.

Mitotic activity informs selected grading systems

- Frequent mitoses can reflect active tumor proliferation.
- Atypical mitoses strengthen evidence of malignant cytology.
- Mitoses also occur in normal regenerating or hyperplastic tissue.
- No universal mitotic-count threshold applies across all cancers.

Mitosis and malignancy are not synonyms


A mitotic figure

- Observed during normal tissue turnover
- May accompany repair or hyperplasia
- Requires clinical and histologic context

An atypical mitosis

- May show abnormal spindle geometry
- Can be seen in highly anaplastic tumors
- Supports atypia but does not define TNM stage

Loss of polarity indicates architectural disorder



Disrupted alignment and polarity


Basement membrane

Stromal compartment

- Normal epithelial cells have organized orientation.
- Anaplastic cells may lose coordinated alignment.
- The architectural disruption is separate from distant spread.

Simplified schematic — not histologic evidence

Ischemic necrosis can accompany rapid growth



Specialized function may track differentiation

Preserved function

- Some thyroid tumors form follicles
- Differentiated squamous cancers may form keratin
- Endocrine tumors may still secrete hormones

Lost or altered function

- Poorly differentiated tumors may lose mature functions
- Some cancers acquire unexpected secretory properties
- Function alone does not establish stage

Grade is not a universal numerical scale

Grading pattern	Source-supported principle	Interpretation
Two-tier	Low grade / high grade	Broad qualitative separation
More categories	Up to four categories described	Finer morphologic stratification
Across tumor types	Criteria differ by organ and neoplasm	Do not transfer thresholds

The same appearance can have different meaning

- Grade definitions are tailored to the biology of each malignancy.
- Gland architecture may matter in one tumor, mitoses in another.
- Specific grading criteria cannot be extrapolated from unrelated organs.
- The generic Robbins framework does not supply organ-specific thresholds.

A descriptive diagnosis can be more defensible


Morphologic description

- “Well differentiated” or “poorly differentiated”
- Record tissue and tumor type
- State conspicuous diagnostic features

Formal grade

- Use only the applicable validated grading framework
- Avoid spurious precision where no scheme is supplied
- Do not fabricate a numerical category

Grade predicts behavior imperfectly



Differentiation does not prove benignity


Well-differentiated carcinoma

- May closely mimic normal cells
- Can still invade adjacent tissue
- May metastasize

Benign neoplasm

- Usually well differentiated
- Lacks malignant invasive/metastatic capacity
- Grade cannot substitute for diagnosis

Invasion is a distinct morphologic question



Simplified schematic — not histologic evidence

- A tumor may look highly differentiated yet invade stroma.
- A preserved basement membrane separates in situ from invasive epithelium.
- Invasion must not be inferred merely from cytologic atypia.

Dysplasia is not tumor grade


Dysplasia

- Disordered epithelial growth
- May be a precursor to malignancy
- Progression is not inevitable

Cancer grading

- Characterizes an established malignancy
- Describes morphologic differentiation
- Applied through tumor-specific systems

Carcinoma in situ is noninvasive



No tumor cells beneath boundary


Basement membrane

Stromal compartment

- Severe epithelial atypia may involve the full thickness.
- Basement membrane remains intact.
- An in situ lesion is not equivalent to stromal invasion.

Simplified schematic — not histologic evidence

Biopsy may sample only part of the tumor



Tumor-center necrosis reduces useful sampling

Tumor center

- May contain ischemic necrosis
- Can yield nonviable material
- May obscure architecture

Viable representative tissue

- Preserves cellular morphology
- Improves architectural evaluation
- Requires suitable handling


The specimen determines what can be evaluated

Material	Strength	Interpretive limitation
Resection	Broad architecture and extent	Availability depends on treatment
Core/incisional biopsy	Tissue architecture in sampled region	May underrepresent heterogeneity
Cytology/FNA	Cell-level atypia	Cannot reliably demonstrate tissue invasion

Cytology does not directly establish invasion

- Cytologic preparations primarily show individual cells or clusters.
- Cellular atypia can suggest malignancy and altered differentiation.
- The tissue boundary needed to demonstrate invasion may be absent.
- Distinguish cytologic impression from full histologic assessment.

Morphologic grading requires technical quality



What grade does—and does not—say

Grade informs

Differentiation and selected atypical features

Sometimes proliferation and architecture

An estimate of tumor aggressiveness

Grade alone cannot establish

Regional node status

Presence of distant metastasis

Precise clinical outcome for an individual

Stage summarizes anatomic disease extent

- Staging examines the primary lesion, regional lymph nodes, and metastases.
- It aims to quantify the apparent spread at assessment.
- The result supports prognosis and comparison of treatment outcomes.
- For solid cancers, TNM provides a structured language.

TNM describes three separate dimensions

T

Primary lesion — size
and/or local extent


N

Regional lymph-node
involvement

M

Distant metastatic
disease

T refers to the primary lesion



T categories are cancer-specific

- T1–T4 as categories of increasing primary-tumor burden.
- Size and local extension must be interpreted by cancer site.
- An identical dimension need not imply an identical T category in two organs.
- Do not assign formal T categories without the site-specific staging manual.

Tumor size is not its entire biological story


Smaller primary tumor

- May still possess aggressive cytology
- Metastasis can sometimes occur
- Size alone does not determine grade

Larger primary tumor

- May show relatively differentiated cells
- Local invasion can affect resectability
- Size alone does not prove metastasis

Local invasion complicates complete excision



Simplified schematic — not histologic evidence

- Malignant growth may cross normal tissue boundaries.
- Tumor margins can be irregular or microscopically infiltrative.
- An apparently circumscribed cancer may still invade.

Anatomic depth differs from cytologic grade


Depth / extent

- Where the tumor has reached
- Whether adjacent structures are infiltrated
- A question for local tumor assessment

Cytologic grade

- How tumor cells and architecture look
- Degree of differentiation
- A biologic morphology descriptor


N refers to regional lymph nodes



Lymphatic spread commonly begins regionally

- Carcinomas commonly disseminate first through lymphatics.
- Routes often follow the ordinary regional drainage pathways.
- Regional spread is not synonymous with hematogenous metastasis.
- Unusual drainage and skip metastases can occur.

A sentinel node samples first-echelon drainage



An enlarged node is not necessarily malignant

Reactive enlargement

- Tumor antigens may provoke nodal immune responses
- Size can increase without tumor deposits
- Clinical impression is not tissue proof

Metastatic involvement

- Tumor cells present in nodal tissue
- Histologic examination can establish involvement
- N interpretation needs relevant nodal evidence

N categories encode regional involvement

- N0 indicates no regional nodal involvement in the source framework.
- N1–N3 represent increasing nodal disease by applicable system.
- The involved number, range, or stations are cancer-dependent.
- A generic N label is not a substitute for organ-specific rules.

M records distant metastatic disease

No identified distant metastasis

- No distant deposits established in the assessment
- Notation depends on formal staging rules
- Do not equate absence of evidence with a guaranteed absence

Distant metastasis identified

- Cancer found at a physically discontinuous site
- Defines systemic spread for many solid cancers
- Major clinical and prognostic consequence


Blood-borne metastases extend beyond the primary region



Metastatic pattern follows anatomy and biology

- Venous drainage influences where circulating tumor cells arrest.
- Liver and lungs are common sites of hematogenous deposits.
- Organ-specific tumor biology influences metastatic distribution.
- The presence of metastasis is not predicted perfectly by grade.

Serosal seeding is another route of spread



Invasion and metastasis are separable

Locally invasive, rarely metastatic

- Some malignant tumors infiltrate despite few distant deposits
- Local progression still matters clinically
- Do not equate invasion with inevitable M1

Distant metastatic tumor

- Cells have spread to discontinuous sites
- Can occur despite an initially small primary
- M status must be assessed directly

T, N, and M should not be collapsed early

T

Record the primary-site
extent

N

Record regional nodal
evidence

M

Record distant spread
evidence

TNM components require site-specific synthesis

Descriptor	What it describes	What it is not
T	Extent of the primary cancer	A universal size cutoff
N	Regional nodal spread	Any lymph-node enlargement
M	Distant metastatic spread	A surrogate for histologic grade

Grading and TNM answer different questions

Histologic appearance ↓

Limited anatomic extent

Extensive anatomic spread

More differentiated

Low morphologic atypia;
Limited spread

Low morphologic atypia;
Extensive spread

Less differentiated

High morphologic atypia;
Limited spread

High morphologic atypia;
Extensive spread

Anatomic spread →

Stage is generally more clinically useful

- Stage over grade in broad prognostic value.
- Anatomic spread strongly influences the possibility of definitive control.
- The same grade may occur with different extent of disease.
- Grade still contributes useful independent morphologic information.

Clinical assessment and tissue findings complement

Imaging / clinical evaluation

- Estimates extent and distant disease
- Helps identify involved structures
- Cannot always determine microscopic spread

Surgical / pathologic assessment

- Can directly examine sampled tumor and nodes
- Documents tissue infiltration
- Limited by what has been removed or sampled

Unexamined tissue creates classification limits

Sampling is incomplete

Unobserved anatomic compartments



Extent is uncertain

Evidence does not justify precision



Report limitations clearly

Avoid unsupported TNM claims


Anatomic staging needs diagnostic context

- Confirm the nature of the primary malignancy.
- Interpret local spread in the correct organ context.
- Distinguish regional from distant sites using the relevant definitions.
- Use the appropriate site-specific staging rules for a formal assignment.

Grade and stage must appear separately in reports

Diagnostic dimension	Typical report content	Potential error
Tumor diagnosis	Type and site of malignancy	Replacing diagnosis with grade
Grade	Histologic differentiation / system	Using generic grading thresholds
Extent / TNM	Anatomic primary, nodal, metastatic data	Inferring missing components from grade

Morphology and anatomy do not explain all risk



Molecular tumor features can add prognosis

- Molecular information as complementary to anatomic staging.
- Molecular prognostic information can be independent of apparent spread.
- The textbook uses selected early breast cancers as an example.
- No specific assay or biomarker thresholds are prescribed in this excerpt.

Early breast cancer illustrates multimodal assessment

Anatomic description

T1–2, N0, M0 is the textbook example

Molecular characterization

May refine prognostic evaluation

Interpretation

Combined information is preferable to one axis alone

A prognostic biomarker is not a TNM category

Molecular or biologic feature

- Can help stratify likely outcome
- May add information beyond stage
- Requires validated disease-specific interpretation

Anatomic descriptor

- Records primary, regional, and distant extent
- Must follow staging definitions
- Does not by itself reveal every biological feature

Do not silently modernize staging criteria

- TNM manuals and disease-specific categories can change.
- This lecture teaches the general framework in the supplied Robbins text.
- Exact numeric size cutoffs and stage groups are intentionally omitted.
- Formal contemporary case staging requires current site-specific authority.

Case exercise: Differentiated tumor with spread

CLINICAL STEM

A biopsy shows recognizable gland formation. Imaging identifies deposits at a discontinuous distant site.

DISCUSSION PROMPTS

- Which evidence informs grade?
- Which evidence informs metastatic status?
- Can an apparently differentiated carcinoma be metastatic?

Case interpretation: Appearance does not cancel metastasis

Grade evidence

Gland formation suggests retained differentiation

Stage evidence

Distant deposits establish dissemination

Core principle

Morphologic differentiation does not preclude metastasis

Case exercise: Marked atypia, apparently localized

CLINICAL STEM

A resected tumor is highly pleomorphic, with abnormal mitoses. Available assessment finds no nodal or distant tumor.

DISCUSSION PROMPTS

- What findings describe its phenotype?
- What data describe extent?
- Should a numeric stage group be assigned here?

Case interpretation: Phenotype is not extent

Grade evidence

Pleomorphism and atypical mitoses

Stage evidence

Extent depends on primary, nodes, metastases

Limitation

No formal TNM category without site-specific data

Case exercise: Palpable regional lymph node

CLINICAL STEM

An individual with carcinoma has an enlarged regional draining lymph node. The node has not been sampled.

DISCUSSION PROMPTS

- Does enlargement itself establish nodal metastasis?
- What other explanation can account for enlargement?
- What finding would improve N assessment?

Case interpretation: Nodal enlargement is ambiguous

Observation

The node is enlarged

Competing explanation

Reactive immune
hyperplasia

Needed evidence

Examination for tumor
deposits, as clinically
indicated

Five distinctions that prevent staging errors

Do not confuse

With

Differentiation

Anatomic spread

Atypical mitoses

Evidence of metastasis

In situ growth

Stromal invasion

Enlarged regional node

Proven nodal metastasis

Molecular prognosis

A universal TNM substitute

Reporting checklist for a solid malignancy

- Identify the tumor type and origin.
- Use a defensible tumor-specific histologic grade when applicable.
- Document primary extent and available regional nodal findings.
- Record distant spread evidence and unresolved assessment limits.
- Keep molecular prognostic information distinguishable from anatomy.

Clinical synthesis: Three distinct information layers

What is it?

Diagnosis / lineage /
differentiation

Where is it?

Primary extent, nodes,
distant sites

How might it behave?

Integrated morphology,
anatomy, molecular
features

Consolidation questions

- Why can a well-differentiated carcinoma still have a poor prognosis?
- How do mitoses contribute to some—but not all—grading systems?
- What additional information is needed beyond tumor size?
- Why is a single regional node enlargement not automatically N-positive?

Source and interpretation notes

Robbins, Cotran & Kumar: Pathologic Basis of Disease. 11th ed. (2026). Chapter 7: Neoplasia.

Principal section: “Grading and Staging of Tumors,” pp. 295–296.

Adjacent context: “Differentiation and Anaplasia,” pp. 239–241; invasion/metastasis, pp. 241–244.

Associated clinical/laboratory context: “Histologic and Cytologic Methods,” pp. 296–297.

Formal site-specific TNM criteria are not reproduced; consult the relevant current staging manual before clinical use.