

Histone Organization

Chromatin architecture, epigenetic regulation and related gene-control mechanisms

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Describe the nucleosome, histone octamer and linker DNA.
- Relate euchromatin and heterochromatin to transcriptional accessibility.
- Explain chromatin remodeling and the functions of writers, readers and erasers.
- Connect reversible epigenetic regulation with disease and treatment.
- Distinguish chromatin control from miRNA, lncRNA and CRISPR mechanisms.

Three molecular levels of gene control

DNA sequence



Chromatin accessibility




RNA-level control



Protein output

Identical genome, different cell identities



Different phenotypes arise from selective gene expression—not a different genome.

Epigenetic factors: Regulation above DNA sequence

Core concept

- Differentiated cells share nearly the same genetic composition.
- Cell-type identity depends on lineage-specific gene-expression programs.
- Epigenetic mechanisms alter the use of DNA without requiring sequence change.

Why it matters

Chromatin state
contributes to selective
transcription and cellular
phenotype.

Transcription is not dictated by DNA sequence alone

- Promoters and enhancers encode regulatory potential.
- Nucleosome position determines whether regulatory sites can be accessed.
- Histone modifications and chromatin organizers add another regulatory layer.


Distinction

DNA sequence defines
possible interactions;
Chromatin affects
physical access.

The physical DNA-packaging problem

~ 1.8 m

DNA in a single human cell



Packaging makes the genome physically compatible with nuclear dimensions.

Why DNA binds histone proteins

DNA backbone
Negative charge (–)




Histone proteins
Positive charge (+)

Electrostatic association

Nucleosome: The repeating unit of chromatin


Histone octamer



DNA wound around core

Schematic; Core DNA: 147 base pairs.

The histone core is an octamer



H2A × 2

H2B × 2

H3 × 2

H4 × 2

Two copies each of H2A, H2B, H3 and H4 form the octamer.

The core wraps 147 base pairs of DNA

Nucleosomal DNA

- A nucleosome core contains approximately 147 bp of DNA.
- DNA winds around a positively charged histone octamer.
- This repeating structure compacts DNA while permitting regulation.


Remember

147 bp describes the DNA wrapped on the core—not the linker.

Linker DNA and histone H1


Core DNA: 147 bp

Linker DNA: ~20–80 nt



H1 is a linker histone, distinct from the H2A/H2B/H3/H4 core octamer.

Chromatin as a beads-on-a-string fiber



Repeated nucleosomes are linked by short DNA segments—the “beads on a string” model.

Compaction and transcriptional access are coupled


Packaging

DNA–histone association allows a very long genome to fit within the nucleus.


Regulation

Local remodeling and histone marks modulate access to regulatory DNA.

Chromatin is not uniformly condensed



Euchromatin



Heterochromatin

Left: Dispersed chromatin; Right: Condensed, intensely stained chromatin. Schematic only.

Euchromatin: Dispersed, transcriptionally active

- Relatively open chromatin in the light-microscopic nucleus.
- Genes in these regions are generally more available for transcription.
- Histone marks supporting access to RNA polymerase are common.

Microscopy

Less densely stained
nuclear regions
correspond broadly to
euchromatin.

Heterochromatin: Condensed, transcriptionally inactive

- Relatively dense chromatin identifiable at light microscopy.
- Compaction reduces access to regulatory sequences and transcription machinery.
- Chromatin-associated marks help maintain gene silencing.

Microscopy

More intense nuclear staining reflects a compact chromatin state.

Euchromatin versus heterochromatin

Feature	Euchromatin	Heterochromatin
Organization	More dispersed	More condensed
Typical transcription	More active	More inactive
Regulatory access	More accessible	More restricted
Light microscopy	Less dense regions	Dense chromatin regions

What chromatin pattern reveals in a nucleus

- A light-microscopic image shows spatial organization of chromatin.
- The euchromatin–heterochromatin distinction is a useful structural approximation.
- Transcriptional status cannot be fully inferred from a single nuclear image.

Interpretation

Morphology suggests chromatin state; It does not identify the exact gene program.


Access to DNA determines transcriptional opportunity

Accessible region



Regulatory DNA exposed

Compact region



Regulatory DNA obscured

A simplified illustration of relative nucleosome density and DNA access.

Regulatory DNA must be physically accessible

- Promoters are docking regions for the transcription apparatus.
- Enhancers can influence transcription when appropriately positioned.
- Nucleosomes can expose or conceal these regulatory elements.

Mechanism

Chromatin remodeling
changes
promoter/enhancer
accessibility without
changing the DNA letters.


Chromatin-remodeling complexes move nucleosomes

- Remodeling complexes reposition nucleosomes along DNA.
- Relocation can expose a formerly obstructed regulatory element.
- Relocation can also hide an element and suppress transcription.

Key distinction

A remodeling complex primarily changes nucleosome position, not DNA sequence.

Nucleosome repositioning changes promoter exposure



Initially hidden promoter

Remodeling complex

Promoter exposed

Nucleosome repositioning can obscure or reveal cis-regulatory DNA.

Writers, readers and erasers of histone marks

Writers

Place modifications

Readers

Bind selected marks

Erasers

Remove modifications

Regulatory outcome depends on which residue is modified and which protein recognizes it.

Histone marks are dynamic regulatory signals

- More than 70 types of histone modification.
- Different marks are placed on particular histone amino-acid residues.
- The meaning of a mark depends on its position and protein context.

Avoid oversimplification

A “histone mark” does not invariably mean activation or repression.

Three major classes of histone modification

Acetylation

Lysine

Methylation

Lysine / arginine

Phosphorylation

Serine

Modification identity alone does not specify the direction of gene expression.


Histone acetylation favors accessible chromatin

- Histone acetyltransferases (HATs) acetylate lysine residues.
- Acetylation tends to promote an open chromatin configuration.
- Such regions are generally more permissive to transcription.

Typical effect

Acetylation → relaxed chromatin tendency → more transcriptional access.

Histone acetyltransferases: Adding acetyl groups



Typical tendency: Increased transcription


Histone deacetylases reverse lysine acetylation

- Histone deacetylases (HDACs) remove acetyl modifications.
- Deacetylation generally favors chromatin condensation.
- Acetylation state is consequently a potentially reversible regulatory feature.

Typical effect

HDAC activity → lower acetylation → more compact chromatin tendency.

Acetylation and deacetylation are reversible



These are tendencies, not universal on/off switches for every gene.

Histone methylation has context-dependent effects

- Specific enzymes methylate histone lysine or arginine residues.
- Histone lysine methylation can be associated with activation or silencing.
- The particular marked residue determines much of the interpretation.

Exam pitfall

Do not equate every methyl group on a histone with gene repression.

Methylation is not a universal off switch

One histone residue

Methylation may contribute to a transcriptionally active chromatin state.

Another histone residue

Methylation may instead promote condensed, silent chromatin.

The source emphasizes residue specificity; It does not provide a universal list of named sites.

Histone phosphorylation can support different states

- Phosphate groups may be added to selected serine residues.
- The associated chromatin may become more accessible or more condensed.
- Functional significance depends on the residue and context.


Concept

“Phosphorylation” specifies a chemical mark—not a fixed transcriptional outcome.

Compare histone modifications

Modification	Target residue	Effect
Acetylation	Lysine	Usually more accessible chromatin
Methylation	Lysine or arginine	Activation or repression, site-dependent
Phosphorylation	Serine	Access or condensation, site-dependent

DNA methylation is a separate regulatory layer



Contrast: Histone methylation may either activate or repress transcription.

DNA methylation has its own regulators

- DNA methyltransferases place methyl marks on DNA.
- Demethylating enzymes help reverse such modifications.
- Methylated-DNA-binding proteins recognize selected methylated sites.

Usual result

High methylation of regulatory elements commonly accompanies transcriptional silencing.

Histone methylation is not DNA methylation

Feature	Histone methylation	DNA methylation
Modified substrate	Histone amino-acid residues	DNA
Common outcome	Activation or repression	Regulatory-element methylation usually silences
Regulators	Histone writer/reader/eraser systems	DNA methyltransferases and demethylating systems

Chromatin looping brings distant elements together

3D looping juxtaposes distant DNA elements



Enhancer

Promoter

Conceptual only: Spatial regulation depends on chromatin-organizing proteins.


Chromatin organizers modulate 3D DNA relationships

- Chromatin-organizing factors bind noncoding genomic regions.
- DNA looping can bring enhancers into functional proximity with promoters.
- Spatial genome organization adds regulation beyond local histone marks.

Limits


The Robbins chapter explicitly notes that these organizing factors remain incompletely characterized.

Cell-type gene expression emerges from several levels



No single chromatin modification can explain all cell-specific gene-expression patterns.

Epigenetic dysregulation and malignancy



Epigenome dysregulation as a central feature in malignant disease.

Epigenetic changes may be inherited or acquired

- Altered epigenetic regulation can occur without changing protein-coding DNA sequence.
- Epigenetic abnormalities to malignant disease.
- Inherited and acquired epigenetic changes are also implicated in other diseases.

Distinction

Epigenetic mechanisms are separate from—but interact with—genetic mutations.

Reversibility provides a therapeutic rationale

- Histone acetylation and DNA methylation are modifiable chemical states.
- Some epigenetic alterations are therefore amenable to pharmacologic intervention.
- DNA methylation inhibitors are used in cancer treatment, as.

Clinical principle

Targeting a reversible chromatin regulator may restore or alter gene expression.

Histone organization: Mechanism recap

Physical substrate

DNA is wrapped around a histone octamer; H1 associates with linker DNA.

Regulatory state

Accessible euchromatin contrasts with condensed heterochromatin.

Dynamic control

Remodelers and histone-mark writers/readers/erasers govern DNA access.

Clinical link

Epigenome disruption is implicated in malignancy and is potentially druggable.

Application: Interpreting nuclear chromatin

Two differentiated cells have the same genomic DNA sequence. One shows more dispersed nuclear chromatin and expresses a lineage-specific gene panel.

Consider

- What structural feature could help explain the expression difference?
- Why would different promoter sequences not be necessary?

Application: Blocking an epigenetic enzyme

A compound decreases the activity of a histone deacetylase in a model cell line.

Consider

- What happens to the acetylation–deacetylation balance?
- What general change in chromatin access would be expected?
- Why does that not guarantee activation of every gene?

Noncoding RNAs extend gene regulation beyond chromatin

MicroRNAs (miRNAs)

Short noncoding RNAs that guide post-transcriptional silencing of target mRNAs.

Long noncoding RNAs (lncRNAs)

Long transcripts with diverse regulatory effects, including chromatin and transcriptional control.


MicroRNAs modulate protein expression

- MiRNAs are small RNAs, averaging about 22 nucleotides.
- They do not encode proteins.
- By pairing with target mRNAs, they can reduce translation or favor mRNA cleavage.

Regulatory stage

MiRNAs act primarily after transcription, whereas histone marks affect access for transcription.

Biogenesis of an active microRNA




Dicer helps produce mature regulatory RNAs

- MiRNA genes are transcribed into a primary precursor (pri-miRNA).
- Progressive processing includes cleavage by the enzyme Dicer.
- Mature miRNA strands are about 21–30 nucleotides and associate with RISC.

Important term

RISC = RNA-induced silencing complex.

RISC targets messenger RNA



MiRNA–mRNA base pairing guides post-transcriptional gene silencing.

One microRNA can affect many protein-coding genes

- A single miRNA can recognize multiple mRNA targets.
- Consequently, one miRNA can coregulate a broad expression program.
- This contributes to developmental control and disease states, including cancer.

Systems effect

Target multiplicity links short RNAs to network-scale changes in protein expression.

Synthetic siRNAs provide experimental gene knockdown

- Small interfering RNAs (siRNAs) can be introduced into cells.
- The source describes their use of Dicer/RISC-related machinery.
- Designed siRNAs permit reduction of expression from selected mRNA targets.

Research application

Knockdown changes transcript abundance or translation without editing the DNA sequence.

SiRNA technology also has therapeutic applications

- Synthetic siRNAs can target excessive or inappropriate gene function.
- Approved therapeutic applications.
- Selected forms of amyloidosis are given as an example.

Boundary

RNA interference is not
equivalent to CRISPR
DNA editing.


Long noncoding RNAs can regulate chromatin

- LncRNAs are non-protein-coding transcripts longer than 200 nucleotides.
- Some lncRNAs associate with chromatin and limit RNA polymerase access.
- Other lncRNAs can enhance gene transcription through distinct mechanisms.

Difference from miRNAs

LncRNAs may operate at the chromatin or transcription level, rather than chiefly on mature mRNA.

XIST mediates X-chromosome inactivation



XIST lncRNA acts in cis to establish a repressive environment on its chromosome.

XIST provides a physiological silencing example

- XIST is transcribed from the X chromosome and escapes its inactivation.
- Its transcript spreads as a repressive “cloak” over the source X chromosome.
- The resulting chromatin environment supports chromosome-scale gene silencing.

Principle

A noncoding RNA can alter expression across an entire chromosomal domain.

Enhancer-associated lncRNAs may increase transcription

- Some enhancer loci produce long noncoding RNA transcripts.
- These transcripts can participate in positive regulation of gene promoters.
- The source emphasizes diversity rather than one universal mechanism.

Contrast

LncRNA activity can be activating or repressive depending on its context.

MiRNA and lncRNA: Different regulatory strategies

Property	MicroRNA	Long noncoding RNA
General size	~22 nt on average	>200 nt
Main examples here	MRNA targeting through RISC	Chromatin silencing; Enhancer effects
Functional scope	Post-transcriptional repression	Variable transcriptional regulation


CRISPR–Cas originated in microbial defense

- CRISPR sequences and Cas genes function in prokaryotic defense.
- Bacteria retain sequence information from invading genetic elements.
- Guide RNAs then help Cas proteins recognize matching targets.

Origin


Genome editing adapts a naturally occurring sequence-recognition and cleavage system.

The natural CRISPR defense concept



Simplified conceptual sequence from the Robbins section.

Engineered CRISPR–Cas9 editing



Simplified CRISPR–Cas9 editing mechanism.

Guide RNA defines the target sequence

- An engineered guide RNA is complementary to a selected DNA sequence.
- An approximately 20-base guide targeting segment.
- Guide RNA directs Cas9 to the corresponding genomic site.

Editing step

Target recognition
precedes DNA cleavage
by Cas9.

DNA repair determines the editing outcome

Cas9 DNA cleavage

```
graph TD; A[Cas9 DNA cleavage] --> B[Nonhomologous end joining]; A --> C[Homology-directed repair]; B --> D[Often disruptive indels]; C --> E[Precise sequence change with template];
```

Nonhomologous end joining

Often disruptive indels

Homology-directed repair

Precise sequence change with template

Random disruptive mutations versus precise introduction of genetic material.

Repair mechanisms after Cas9 cleavage

Pathway	Mechanism	Typical editing result
Nonhomologous end joining	Rejoin broken DNA ends	Potentially disruptive mutations
Homology-directed repair	Use a homologous donor sequence	Precisely introduced DNA change

Gene editing is a tool for disease modelling

- Targeted mutations can be introduced in cells or tissues.
- Edited cells can model cancer and other disease mechanisms.
- Transgenic animal models can be generated from edited stem cells.

Research value

The approach probes causal consequences of sequence changes.

CRISPR has entered treatment of inherited disease

- Therapeutic application in inherited disorders.
- Sickle cell disease is an example cited in the chapter.
- Clinical deployment must be distinguished from experimental uses.

Clinical boundary

A gene-editing intervention changes DNA; Epigenetic therapies alter reversible regulatory states.

Genome editing also raises ethical questions

- Sequence modification can be used for laboratory research and clinical intervention.
- Potential use to alter non-disease-associated traits raises ethical concerns.
- Robbins explicitly identifies a vigorous debate about these applications.

Scope

Evaluate intended use, affected cells and consequences—rather than treating all editing as equivalent.

Compare levels of intervention in gene regulation

Chromatin



DNA access for transcription

MiRNA / lncRNA



RNA-mediated gene regulation

CRISPR-Cas9



Intentional alteration of DNA sequence

Case: Identify the regulatory mechanism

A cell maintains the same DNA sequence but changes transcription after nucleosomes are repositioned away from a promoter. A second experiment depletes a target mRNA using an introduced short RNA.

Consider

- Which experiment changes chromatin accessibility?
- Which intervention acts at the RNA level?
- Does either require changing the genomic sequence?

Case: Explain a targeted DNA modification

An investigator delivers Cas9 and an engineered guide RNA. A double-strand break is produced at the targeted genomic site.

Consider

- What is the guide RNA's function?
- Which repair route commonly yields disruptive mutations?
- Which route can install donor-encoded sequence?

Synthesis: Principles to retain

Nucleosomes

147 bp of DNA around a core octamer, with H1 associated with linker DNA.

Chromatin state

Euchromatin is generally accessible; Heterochromatin is condensed and silent.

Epigenetic marks

Acetylation generally opens; Methylation and phosphorylation are site-dependent.

Other regulators

MiRNAs and lncRNAs influence gene expression; Cas9 can directly alter DNA.

References and source boundaries

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. (2026). Chapter 1, “The Cell as a Unit of Health and Disease,” pp. 3–5.
- Primary sections: Histone Organization; MicroRNA and Long Noncoding RNA; MicroRNA; Long Noncoding RNA; Gene Editing.
- All molecular schematics in this presentation are original, editable teaching diagrams; they are not reproduced textbook figures.