

Indoor Air Pollution

Exposure, mechanisms, and clinicopathologic consequences

Dr Abu Suraih Sakhri


MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Classify major indoor airborne pollutants by source and biologic action.
- Explain combustion-related inflammation, hypoxic injury, and carcinogenesis.
- Distinguish infectious, allergic, and toxic indoor-air presentations.
- Apply exposure-oriented reasoning to representative clinical scenarios.

Indoor air is an exposure compartment



The indoor environment can intensify exposure when emissions persist and ventilation is inadequate.

Four principal indoor hazard families

Combustion

Smoke, particles, CO, nitrogen oxides

Biologic

Microbes and allergens

Geologic

Radon and its decay products

Materials


Formaldehyde and asbestos

Category describes source; Clinical consequences depend on the agent and dose.

Ventilation changes the exposure context

- Increasingly sealed buildings may retain internally generated pollutants.
- The same indoor space can contain gaseous, particulate, and biologic hazards.
- Poor air exchange is a potential contributor—not a universal explanation—for symptoms.
- Risk assessment must link a source, a plausible route, and an outcome.

From pollutant source to clinical effect



Different agents may produce irritation, hypoxia, allergy, infection, or malignancy.

Why individuals respond differently

- Host susceptibility varies with age, genetic predisposition, and tissue sensitivity.
- Asthma and chronic pulmonary disease can increase the impact of irritant exposure.
- Exposure intensity, duration, and mixed pollutants complicate simple risk thresholds.
- A safe level inferred for a population need not predict every individual response.

Indoor versus outdoor air: Overlapping hazards

Shared hazards

- Carbon monoxide and nitrogen dioxide occur in both settings.
- Particulate exposure may arise from combustion sources.
- The respiratory system is a principal entry site.

Indoor-specific emphasis

- Confinement can concentrate locally emitted pollutants.
- Building materials introduce additional volatile compounds.
- Indoor animals, dust and moisture influence bioaerosols.

Organ responses depend on pollutant class

Agent class	Principal pathway	Representative effect
Combustion gases	Irritation or oxygen transport	Asthma symptoms; systemic hypoxia
Particles	Airway/alveolar inflammation	Respiratory infection susceptibility
Bioaerosols	Infection or sensitization	Pneumonia; rhinitis/asthma
Radon	Ionizing radiation	Lung carcinoma

Domestic combustion: Source diversity

Solid biomass

Wood, dung, charcoal

Other fuels

Kerosene, lignite, peat

Coal combustion

Smoke and aromatic compounds

Cooking oil


Volatile aromatic compounds

Fuel and cooking practices influence the mixture encountered indoors.

Household fuel burning is a global exposure

- Approximately one-third of the world's people were estimated to burn carbon-containing fuel in their homes.
- Cooking, heating, and lighting are relevant activities.
- The specified fuels include wood, dung, charcoal, kerosene, lignite, and peat.
- Household burning represents everyday domestic exposure, not only an industrial hazard.

Combustion creates mixed airborne hazards




Combustion is not a single-pollutant phenomenon.

Smoke acts first as a respiratory irritant

- Organic combustion smoke contains various oxides of nitrogen and carbon particulates.
- Inhaled irritants contact respiratory mucosa and can provoke local injury.
- Combustion smoke exposure is associated with increased susceptibility to lung infections.
- The exact disease depends on the mixture and the exposed host.

Smoke and respiratory infection susceptibility




Smoke-related infection susceptibility does not establish which pathogen causes an illness.

Particles can reach the distal lung

- Fine particles are more likely than larger particles to reach the distal respiratory tract.
- Particles reaching alveoli may be engulfed by macrophages and neutrophils.
- Activated phagocytes release inflammatory mediators.
- Particle deposition and innate immune activation explain a pulmonary inflammatory response.

Alveolar particulate response



Schematic: Biologic response to respirable particles; No tissue photomicrograph.

Local irritation versus systemic consequence

Airway-dominant injury

- Smoke and nitrogen oxides irritate mucosal surfaces.
- Inflammatory responses alter airway function.
- Preexisting lung disease increases vulnerability.

Systemic hypoxic injury

- CO does not need to irritate the airway.
- CO binds hemoglobin and compromises oxygen carriage.
- Neurologic injury may be prominent.

Polycyclic aromatic hydrocarbons are a separate hazard

- Volatile compounds containing PAHs may arise from cooking oils and coal burning.
- These emissions are important indoor hazards in lower-resource settings, notably parts of Asia.
- Some organic combustion smoke may contain carcinogenic PAHs.
- PAH exposure links combustion environments to potential long-term cancer risk.

A carcinogen can require metabolic activation



The adjacent toxicology discussion uses benzo[a]pyrene as an example of activation.

A crucial distinction: Presence versus cancer causation

- Finding a potential carcinogen in smoke identifies a biologic hazard.
- Dose, duration, absorption, and host factors modify eventual risk.
- Cancer risk cannot be calculated from simple detection of PAHs in indoor air.
- Mechanistic plausibility does not determine an individual's probability of developing malignancy.


Tobacco smoke is a major indoor pollutant

- Tobacco smoke is a major and frequently encountered indoor pollutant.
- Environmental or secondhand inhalation exposes people other than active smokers.
- Passive tobacco exposure is associated with lung cancer in nonsmokers.
- Indoor tobacco exposure should be separated from deliberate active smoking.

Important constituents of tobacco smoke

Constituent	Principal biologic effect
Carbon monoxide	Impaired oxygen transport/utilization
Nitrogen oxides	Ciliary toxicity; mucosal irritation
Formaldehyde	Ciliary toxicity; mucosal irritation
Polycyclic aromatic hydrocarbons	Carcinogenesis
Benzo[a]pyrene	Carcinogenesis

Secondhand tobacco smoke: Causal reasoning



Passive inhalation can increase lung-cancer risk in nonsmokers.

Smoke constituents have different biologic actions

- Carcinogens contribute to mutation-related risk.
- Irritants affect mucosal and ciliary function.
- Carbon monoxide reduces blood oxygen carriage.
- A single indoor emission can therefore generate multiple pathologic outcomes.

Carbon monoxide: An indoor combustion emergency

- Incomplete oxidation of hydrocarbons generates carbon monoxide (CO).
- Potential sources include furnaces, engines, cigarettes, and gasoline-powered generators.
- Poorly ventilated or enclosed settings are especially concerning.
- CO exposure produces systemic toxicity rather than a simple irritant syndrome.

Common environments with CO generation

Heating

Faulty or poorly vented furnace

Power

Gasoline-powered generator

Transport

Running engine in an enclosure

Smoking


Cigarette combustion

All listed scenarios involve fuel combustion and potential enclosed-space exposure.

Carbon monoxide cannot be detected by smell

- CO is colorless, tasteless, odorless, and nonirritating.
- Lack of obvious warning sensations makes exposure difficult to recognize.
- CNS depression may evolve insidiously.
- A plausible source and multiple affected occupants can be more informative than odor.

CO causes functional anemia without loss of hemoglobin




Hemoglobin binds carbon monoxide with approximately 200 times the affinity for oxygen.

The carboxyhemoglobin mechanism

- CO occupies oxygen-binding sites on hemoglobin.
- The resulting carboxyhemoglobin cannot transport oxygen normally.
- Systemic hypoxia follows as functional oxygen-carrying capacity declines.
- Neurologic manifestations reflect tissue vulnerability to inadequate oxygen delivery.

Why the brain is vulnerable in CO poisoning



Hypoxic injury is especially prominent in the basal ganglia and lenticular nuclei.

Acute CO poisoning can be deceptive

- Acute exposure may cause CNS depression, unconsciousness, or death.
- Cherry-red skin or mucosal coloration can occur in some light-skinned individuals.
- That coloration is not a reliable exclusion or screening feature.
- Rapid deaths may occur before visible morphologic changes develop.


CO poisoning has nonspecific tissue morphology

- With longer survival the brain may be mildly edematous.
- Punctate hemorrhages and hypoxia-related neuronal changes may occur.
- The changes reflect systemic hypoxia rather than a CO-specific histologic lesion.
- Absence of characteristic morphology cannot exclude severe exposure.

Persistent exposure creates cumulative neurologic risk

- Stable carboxyhemoglobin can accumulate during prolonged exposure.
- Slowly developing hypoxia may cause widespread CNS ischemic injury.
- Persistent deficits may affect memory, vision, hearing, and speech.
- Chronic toxicity may be less dramatic than acute collapse.

CO exposure: Recognition pathway



Blood carboxyhemoglobin measurement directly supports assessment of CO exposure.

CO toxicity versus airway irritancy


Feature	Carbon monoxide	Combustion smoke/NO ₂
Sensory warning	Odorless; nonirritating	May irritate airways
Principal injury	Impaired oxygen carriage	Respiratory irritation/inflammation
Clinical emphasis	CNS hypoxia	Airway symptoms / asthma
Useful clue	Source + COHb testing	Exposure + respiratory response

Clinical case: Sudden symptoms in a closed room

Hypothetical teaching vignette

- An occupant is found confused after a gasoline generator has run in an enclosed space.
- There is no unusual smell and little cough.
- Question: Which indoor agent and pathogenic mechanism best explain the presentation?

Case resolution: Generator-associated CO injury



Test the exposure hypothesis using blood carboxyhemoglobin and the clinical context.

Nitrogen dioxide is also an indoor pollutant

- Nitrogen dioxide occurs in both indoor and outdoor polluted air.
- Combustion can introduce nitrogen oxides into indoor air.
- Nitrogen dioxide is associated with increased airway reactivity and reduced lung function.
- Children and people with asthma are particularly relevant susceptible groups.

Nitrogen dioxide: Respiratory susceptibility

Population	Associated effect
Healthy adults	Increased airway reactivity
People with asthma	Decreased lung function
Children	Increased respiratory infections

Asbestos is an additional indoor-air concern

- Asbestos is among the additional agents that can contaminate indoor air.
- Inhaled mineral dusts are associated with pneumoconioses.
- Asbestos-related fibrosis and cancers have a distinct mechanism from CO or radon.
- Do not confuse radon, CO, and asbestos: Their mechanisms are distinct.

Bioaerosols comprise two clinical groups

Infectious agents

Airborne microbes

Allergens


Pet dander, mites, fungi, molds

Aerosol exposure may cause disease through infection or through allergic responses.

Infectious bioaerosols: Textbook examples

- Legionnaires' disease is a representative illness in the infectious bioaerosol group.
- It also includes viral pneumonia and the common cold.
- The presence of an airborne organism creates an infectious exposure pathway.
- Agent-specific confirmation is not established by a generic indoor-exposure history.

Infectious indoor aerosol pathway




Pathway illustrates exposure; Actual acquisition depends on organism and circumstances.

Indoor allergens are not synonymous with infection

- Pet dander and dust mites represent allergenic exposures.
- Fungi and molds also contribute allergens in indoor environments.
- Allergic responses can be distressing without a microbial infection.
- An allergen-associated illness should not automatically be treated as pneumonia.

Allergen-associated respiratory illness



Eye irritation may accompany allergen-related indoor-air presentations.

Distinguish microbial from allergenic aerosols

Comparison	Infectious bioaerosol	Allergenic bioaerosol
Representative source	Microorganisms	Dander, dust mites, molds
Primary process	Infection	Allergic response
Examples	Pneumonia; common cold	Rhinitis; asthma
Reasoning	Consider pathogen	Consider allergen exposure

Clinical case: Symptoms in a damp indoor space

Hypothetical teaching vignette

- An individual develops recurrent rhinitis and wheeze when spending time in a room with visible mold.
- Question: Is the strongest hypothesis an infectious illness or allergen-associated disease?
- What additional clinical history would help distinguish the two?


Case interpretation: Exposure is not a diagnosis

- The symptom pattern supports considering allergen-associated rhinitis or asthma.
- Molds can be relevant allergens, but exposure alone does not prove etiology.
- Infectious disease requires compatible clinical evidence and appropriate assessment.
- Ask about symptom timing, other triggers, and response outside the building.

Radon is a soil-derived radioactive gas

- Radon arises naturally from uranium present in soil.
- It can enter homes and contribute to indoor exposure.
- The principal long-term pathologic concern is lung carcinoma.
- Radon differs mechanistically from smoke irritation and CO poisoning.

Radon hazard originates below the building




Original conceptual environmental pathway, not a building measurement.

Radon daughters deliver the damaging radiation

- Polonium-214 and polonium-218 are alpha-emitting radon decay products.
- These daughter products emit alpha particles.
- Inhaled daughter products may deposit in the lung.
- Chronic exposure is associated with pulmonary carcinoma in uranium miners.

Radon carcinogenic mechanism



The diagram depicts a carcinogenic pathway, not a quantified dose-response.

Evidence is strongest for occupational radon exposure

- Sustained radon exposure can cause lung carcinoma in uranium miners.
- Household exposures are also considered a lung-cancer hazard.
- Estimating risk from low-level domestic exposure is less straightforward.
- Evidence for a hazard is not a case-specific diagnosis.

Radon and tobacco have interacting risk contexts

- Coexisting tobacco exposure is particularly concerning in radon-exposed people.
- Tobacco smoke is independently carcinogenic in the respiratory tract.
- Co-exposure is relevant when taking an environmental risk history.
- An individual joint-risk estimate needs appropriate exposure and epidemiologic data.

Textbook radon burden estimates need context

- A textbook-cited US EPA estimate is roughly 21,000 radon-related lung-cancer deaths each year.
- It reports roughly 2,900 among people who have never smoked.
- These numbers are reproduced as textbook-attributed estimates, not current-year surveillance.
- Radon is identified as a major lung-cancer cause in nonsmokers.

Radon versus carbon monoxide: Incompatible mechanisms

Radon

- Radioactive gas associated with uranium-bearing soil.
- Alpha-emitting decay products affect the lung.
- Primary described long-term endpoint: Lung carcinoma.


Carbon monoxide

- Gas released by incomplete hydrocarbon combustion.
- Binds hemoglobin and decreases oxygen delivery.
- Acute or chronic hypoxic neurologic injury.

Formaldehyde can originate in indoor materials

- Manufactured building materials are a recognized formaldehyde source.
- Cabinetry, furniture, and adhesives are potential emission sources.
- Inadequately ventilated housing may permit airborne accumulation.
- This is a source-to-exposure issue, not necessarily a visible material defect.

Materials, ventilation, and mucosal exposure



Qualitative schematic: Pollutant concentrations cannot be inferred from this sequence.

Formaldehyde causes mucosal irritation

- Formaldehyde exposure can cause breathing difficulty and burning of the eyes and throat.
- Irritant effects were described at concentrations of 0.1 ppm or higher.
- It also identifies asthma attacks as a possible response.
- This concentration should not be treated as a universal individual-response threshold.

Formaldehyde has acute and chronic hazard dimensions

Immediate manifestations

- Eye and throat burning.
- Breathing difficulty.
- Triggering of asthma attacks.

Long-term classification

- Formaldehyde is classified as carcinogenic to humans and animals.
- Carcinogenicity does not predict that every exposed person develops cancer.
- Avoid inferring individual risk from a symptom alone.

Comparison of indoor chemical hazards

Agent	Dominant acute issue	Long-term concern
Carbon monoxide	Systemic hypoxia	Neurologic sequelae
Formaldehyde	Mucosal irritation	Carcinogenic classification
Radon	Usually not an irritant syndrome	Lung cancer
Smoke/PAHs	Airway irritation	Carcinogen exposure


Sick building syndrome is a symptom pattern

- The syndrome has been described among office workers who spend long periods indoors.
- Headache and respiratory complaints are characteristic.
- It is a descriptive syndrome rather than proof of one causative chemical.
- A careful exposure assessment must avoid premature causal attribution.

Its underlying cause remains uncertain

- No single underlying causal agent has been established.
- One or more indoor pollutants may contribute.
- Poor ventilation is proposed as a possible contributing condition.
- Do not label all nonspecific building symptoms as a confirmed toxic exposure.

Sick building syndrome: Hypothesis map



This is a hypothesis map only; The causal explanation remains unresolved.

Symptom patterns orient indoor exposure assessment

Presentation	Candidate pathway	Discriminating clue
Confusion / collapse	CO-related hypoxia	Fuel-burning source
Wheeze / rhinitis	Allergenic aerosol	Exposure-linked symptoms
Burning eyes / throat	Irritant gas / formaldehyde	Material or combustion source
Long-latency cancer	Radon / smoke carcinogens	Chronic exposure history

Exposure history should be source-specific

- Ask which fuels, heating appliances, and generators operate indoors.
- Review tobacco exposure and visible or documented building materials.
- Consider dampness, pets, dust, ventilation, and occupancy patterns.
- Separate agent-specific evidence from general symptom associations.

Three levels of causal evidence

Exposure clue

Source plausibly present

Health pattern


Manifestations fit agent

Confirmation

Agent-specific tests when available

An exposure history alone is insufficient to establish an individual diagnosis.

Prevention targets the exposure pathway



Conceptual exposure control; Specific interventions require appropriate environmental and clinical advice.

Integrated case: Unexplained office complaints

Hypothetical teaching vignette

- Several office workers describe headache and respiratory symptoms during long indoor workdays.
- The building has limited fresh-air exchange; No individual agent has been measured.
- Question: What can be concluded, and what remains uncertain?

Case resolution: Avoid overdiagnosis

- The cluster is compatible with the symptom pattern called sick building syndrome.
- Poor ventilation is a plausible factor, not a confirmed mechanism.
- Evaluate occupancy-linked symptoms and possible pollutant sources.
- Do not claim a specific toxin without supporting environmental or medical findings.

Clinical synthesis: Four mechanisms, different endpoints

- Irritants and particles predominantly affect airway and alveolar function.
- Carbon monoxide impairs systemic oxygen transport.
- Bioaerosols can cause infection or allergic manifestations.
- Radon and certain organic chemical exposures raise carcinogenic concerns.

High-yield mechanism comparison

Exposure	Mechanism	Selected clinicopathologic consequence
Smoke particles	Airway/alveolar inflammation	Infection susceptibility
CO	Carboxyhemoglobin formation	CNS hypoxia
Bioaerosols	Microbial infection / allergy	Pneumonia / rhinitis / asthma
Radon daughters	Alpha radiation	Lung carcinoma
Formaldehyde	Mucosal irritation	Burning eyes/throat, asthma attacks

Key interpretive principles

- Multiple pollutants can coexist in a home or workplace.
- Different manifestations suggest different mechanisms; They are not interchangeable.
- Established toxic effects must be distinguished from uncertain exposure associations.
- Exposure assessment requires agent-specific evidence rather than symptom pattern alone.

Primary reading

- Robbins, Cotran & Kumar: Pathologic Basis of Disease, 11th edition.
- Environmental Pollution: Indoor Air Pollution and related carbon monoxide discussion.
- Effects of Tobacco; Cancer Risks From Exposures to Radiation.