

Infectious Rhinitis

Viral etiology · mucosal pathology · clinicopathologic differential

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Identify the named viral causes and the source-defined acute nasal morphology.
- Explain how edema, enlarged turbinates, and catarrh form a coherent clinical syndrome.
- Distinguish rapid uncomplicated recovery from local spread and bacterial superinfection.
- Compare acute infection with allergy, chronic rhinitis, sinusitis, and COVID-19.

Infectious rhinitis is the common cold

- An acute viral inflammatory disorder centered in the nasal mucosa.
- Profuse catarrhal nasal discharge is a defining manifestation.
- Most episodes resolve rapidly rather than evolving into persistent disease.



The nose lies within a continuous upper airway


- The nose, pharynx, and larynx within the upper airway.
- Nasal inflammation may extend locally to pharyngeal and tonsillar tissues.
- The anatomic location explains the distinction between rhinitis and pharyngotonsillitis.



Contiguous upper-airway compartments (schematic)

Nasal inflammatory diseases are common

- The common cold is among the most frequent disorders of the nose.
- Inflammation also affects the accessory air sinuses.
- Viral disease is usual; Superimposed bacterial infection may modify its course.



Infectious rhinitis belongs to a wider differential

- Acute infectious rhinitis represents one nasal inflammatory pattern.
- Allergic rhinitis, chronic rhinitis, and nasal polyps are nearby but distinct entities.
- These conditions can share swelling or discharge without sharing a cause.

Infectious

Allergic

Chronic

Polypoid

More than one virus produces the same syndrome

- Multiple viral causes, not a single cold pathogen.
- Adenoviruses, echoviruses, and rhinoviruses are named major offenders.
- A common mucosal phenotype across these agents.

Adenoviruses

Echoviruses

Rhinoviruses

Named etiologic agents are examples—not a unique test

- The viral list establishes etiologic heterogeneity.
- The section does not assign pathogen-specific nasal gross lesions.
- Therefore, the observed mucosal appearance alone does not identify which named virus is present.

Multiple viral causes


Shared inflammatory pattern

No organism-specific gross pattern

No source-defined diagnostic cutoff


Viral disease and secondary bacteria are different stages

- The initial common-cold process is viral.
- Bacteria may subsequently superinfect inflamed mucosa.
- Secondary infection intensifies inflammation and changes the exudate.



Viral infection initiates the catarrhal reaction


- The named viruses evoke prominent nasal secretion.
- The catarrhal reaction occurs with acute mucosal inflammation.
- Secretion and tissue swelling develop together in the acute stage.



Mucosal erythema is a prominent early finding

- The nasal mucosa becomes visibly red during the initial acute stage.
- Redness is part of the same inflammatory pattern as edema and thickening.
- Erythema documents active mucosal change, not an etiologic species.

**Nasal airway
(schematic)**




Reddened mucosa

Schematic — not histology

Edema makes the nasal mucosa thicker

- The mucosa as edematous and thickened.
- Swelling occupies space within the nasal cavity.
- Reduced luminal space connects tissue pathology with obstructive symptoms.

**Nasal airway
(schematic)**




**Thickened / edematous
lining**

Schematic — not histology

Turbinate enlargement adds to the obstruction

- Turbinates become enlarged in the acute stage.
- Their enlargement occurs alongside diffuse mucosal swelling.
- Together these alterations narrow the nasal airway.

**Nasal airway
(schematic)**




Swollen turbinates

Schematic — not histology

Nasal cavities become narrower

- Narrowed cavities are an explicit gross finding in acute infectious rhinitis.
- Both swollen lining and enlarged turbinates contribute anatomically.
- This is the morphologic basis for impaired nasal airflow.

**Nasal airway
(schematic)**




Reduced airway space

Schematic — not histology

Catarrhal discharge is an exudative phenotype

- The initial discharge is profuse and catarrhal.
- Catarrhal denotes the mucus-rich secretion accompanying mucosal inflammation.
- The early discharge does not by itself establish bacterial superinfection.



Inflammation alters tissue and surface secretions

- Tissue compartment: Thickening, edema, erythema.
- Airway compartment: Narrowing and turbinate enlargement.
- Surface compartment: Profuse catarrhal discharge.

Mucosa


Lumen

Turbinates

Secretions


The nasal phenotype links morphology to symptoms

- Catarrhal secretion is clinically expressed as nasal discharge.
- Mucosal and turbinate swelling provide a basis for congestion.
- These manifestations arise from tissue responses, not from an identifying histologic lesion.



A causal map of early viral rhinitis

- Acute viral infection precedes the mucosal inflammatory reaction.
- Redness and edema describe the mucosal component.
- Turbinates enlarge, cavities narrow, and secretions increase.



Acute morphology is a constellation—not one sign

- No single gross finding described is unique to the common cold.
- Interpret simultaneous redness, edema, narrowed cavities, and discharge.
- Neighboring allergic rhinitis can also produce edema, erythema, and mucus.

Acute infectious: Viral origin


Catarrh and edema

Allergic: Hypersensitivity origin

Edema, erythema, mucus

The earliest stage is described morphologically

- Initial acute disease: Mucosa red, thickened, and edematous.
- Turbinates enlarge and nasal lumina narrow.
- Profuse catarrhal secretions accompany these local changes.



Local extension may involve the throat

- Rhinitic changes can spread to produce pharyngotonsillitis.
- This represents contiguous upper-airway involvement.
- Pharyngeal extension is not identical to bacterial superinfection.



Contiguous upper-airway compartments (schematic)

Rhinitis and pharyngotonsillitis describe sites

- Rhinitis identifies nasal mucosal inflammation.
- Pharyngotonsillitis identifies inflammation involving pharynx and tonsils.
- One acute upper-airway process can involve both locations.

Rhinitis

Nasal mucosa

Catarrhal discharge


Pharyngotonsillitis

Pharynx + tonsils

Local extension

Bacterial superinfection intensifies inflammation

- Bacteria as a secondary event.
- When superinfection occurs, the inflammatory reaction is enhanced.
- The character of the exudate may evolve beyond simple catarrh.



Mucopurulent discharge signals a changed exudate

- Secondary bacterial infection can produce mucopurulent exudate.
- This is contrasted with the initial profuse catarrhal secretion.
- The source does not attribute this exudate to any particular bacterium.

Initial illness


Catarrhal discharge

With superinfection

Mucopurulent discharge

Suppuration is a possible—not obligatory—change

- A suppurative exudate can sometimes occur with bacterial superinfection.
- It is presented as an alternative to mucopurulent exudate.
- Do not treat suppuration as a routine feature of uncomplicated viral rhinitis.



Exudate descriptors have different meanings

- Catarrhal: Predominant pattern in the primary viral illness.
- Mucopurulent: Described after secondary bacterial infection.
- Suppurative: An occasional form of enhanced inflammatory exudate.

Context	Described exudate
Primary viral episode	Profuse catarrhal
Secondary bacterial event	Mucopurulent
Occasional enhanced reaction	Suppurative

Avoid a false shortcut from appearance to species

- The text associates bacterial superinfection with altered exudate.
- It does not supply an organism-specific appearance or diagnostic accuracy.
- Thus morphology suggests a process; It does not microbiologically type the cause.

What the source supports


Change in exudate

What it does not establish

Species identification


Uncomplicated and complicated courses diverge

- Many acute viral rhinitis episodes clear quickly.
- Some cases extend locally to the pharynx or tonsils.
- Bacterial superinfection creates a second inflammatory pathway.



Spontaneous clearance is typical

- That infections usually clear rapidly.
- Its seven-day aphorism communicates the self-limited course.
- The passage does not provide individual prognostic thresholds or treatment comparisons.



The textbook aphorism is qualitative

- The statement “a week if treated, 7 days if ignored” is deliberately humorous.
- Its point is that an uncomplicated common cold is often short-lived.
- It is not a clinical trial or a precise duration prediction for each patient.

The common cold has no described specific biopsy signature

- Acute gross mucosal changes rather than an acute microscopic grading scheme.
- The excerpt does not describe diagnostic immunohistochemistry or molecular pathology.
- Native schematics in this lecture illustrate anatomy—not observed histology.

**Nasal airway
(schematic)**




Uninflamed illustration

Schematic — not histology

Interpret red, edematous nasal mucosa as inflammation

- Reddening and edema are direct morphological observations.
- Neither observation alone distinguishes infectious from allergic rhinitis.
- Etiologic interpretation requires the wider clinical and source-defined context.

**Nasal airway
(schematic)**




Reddened mucosa

Schematic — not histology

The swollen turbinate is an anatomic clue

- Turbinate enlargement is specifically described during acute infectious rhinitis.
- Its position within the airway makes the narrowed cavity visible conceptually.
- A schematic is preferable to inventing a histologic image.

**Nasal airway
(schematic)**




Swollen turbinates

Schematic — not histology

Airway narrowing explains clinical impact

- A widened mucosa occupies some of the nasal passage.
- Enlarged turbinates leave less patent space.
- The combination explains why a common viral process can be troublesome without being destructive.

**Nasal airway
(schematic)**




Reduced airway space

Schematic — not histology

Superinfection is recognized as a process change

- The initial disease is described as viral and catarrhal.
- Subsequent bacterial involvement enhances the inflammatory response.
- Mucopurulent or suppurative exudate fits this source-defined evolution.



A localized cold can broaden its anatomic territory

- Primary location: Nasal cavity.
- Contiguous spread: Pharynx and tonsils.
- Separately, rhinitis may precede sinusitis in adjacent anatomical spaces.



Contiguous upper-airway compartments (schematic)

Acute rhinitis is not synonymous with chronic rhinitis

- Acute infectious rhinitis is described as usually rapidly resolving.
- Chronic rhinitis follows repeated microbial or allergic episodes.
- The latter may develop superimposed bacterial infection and persistent tissue injury.

Acute infectious

Typical rapid clearance

Acute edema / catarrh

Chronic

Repeated episodes

Surface injury / mixed cells


Chronic rhinitis introduces epithelial damage

- In chronic rhinitis, superficial epithelial desquamation or ulceration is common.
- The infiltrate may include neutrophils, lymphocytes, and plasma cells.
- These findings belong to the chronic section, not automatically to every cold.

Chronic lesion	Source-described feature
Epithelial surface	Desquamation or ulceration
Inflammatory infiltrate	Neutrophils, lymphocytes, plasma cells

Impaired drainage favors persistent infection

- A deviated nasal septum can impair secretory drainage.
- Nasal polyps can also obstruct drainage pathways.
- These alterations contribute to microbial invasion in chronic rhinitis.




Acute and chronic rhinitis differ in evidence

- Acute: Viral causes, catarrh, swollen red mucosa, rapid resolution.
- Chronic: Repeated inflammation, bacterial infection, surface desquamation/ulceration.
- The distinction is etiologic, temporal, and morphologic.

Dimension	Acute infectious	Chronic
Pattern	Usually brief	Repeated episodes
Source emphasis	Edema + catarrh	Surface injury
Secondary bacteria	May occur	Often superimposed


Allergic rhinitis begins without a viral trigger

- Allergen hypersensitivity as the initiating event.
- Examples include pollens, fungi, animal allergens, and dust mites.
- The causal category differs from the virus-initiated common cold.



IgE mechanisms distinguish allergic rhinitis

- Allergic rhinitis is an IgE-mediated immediate hypersensitivity disorder.
- Early and late phases of the allergic response.
- The cellular infiltrate is eosinophil-rich.



Clinical morphologies overlap despite different causes

- Both acute infection and allergy may cause mucosal edema and erythema.
- Both can increase mucus production.
- Shared morphology should not erase the distinction between viral infection and IgE-mediated hypersensitivity.

Infectious rhinitis

Multiple viruses

Profuse catarrh

Allergic rhinitis

IgE hypersensitivity

Eosinophil-rich infiltrate


Eosinophil-rich inflammation is an allergic clue

- Robbins explicitly describes eosinophils in allergic rhinitis.
- The infectious-rhinitis paragraph does not give a corresponding acute cellular profile.
- Do not invent a microscopy-based cold diagnostic rule from this comparison.

Entity	Supported morphologic detail
Allergic rhinitis	Eosinophil-rich infiltrate
Infectious rhinitis	Acute gross edema, erythema, catarrh

Nasal polyps can follow recurrent rhinitis

- Recurrent rhinitis can produce focal mucosal protrusions.
- These lesions may project into the nasal airway.
- Their presence represents a structural consequence rather than the initial viral cold lesion.




Polyp pathology is not the same as acute catarrh

- Polyps show edematous loose stroma and altered mucus glands.
- Inflammatory cells may include eosinophils, neutrophils, and plasma cells.
- This is the separately described polyp pattern, not a diagnostic micrograph of acute rhinitis.

Polyp component	Morphology
Stroma	Loose, edematous
Glands	Hyperplastic or cystic
Inflammatory cells	Eosinophils, neutrophils, plasma cells


Polyps can compromise both airway and sinus drainage

- Large or multiple polyps may obstruct nasal airflow.
- Polyp-related obstruction can impair sinus drainage.
- This connects repeated nasal disease to later anatomical complications.




Most acute sinusitis is preceded by rhinitis

- Robbins explicitly links many episodes of acute sinusitis to prior rhinitis.
- Inflammatory swelling can impair sinus outflow.
- Sinus inflammation belongs to an adjacent compartment and should be identified separately.




Impaired drainage can amplify sinus disease

- Edema reduces the effectiveness of sinus drainage.
- The source describes possible empyema of the sinus.
- The relationship is a complication pathway—not an inevitable result of a cold.



Sinusitis has uncommon but serious extensions

- The sinusitis section describes potential extension into the orbit.
- Bone involvement can lead to osteomyelitis.
- Further spread may involve the cranial vault and dural venous sinuses.



COVID-19 belongs in the differential

- SARS-CoV-2 infection may produce upper respiratory symptoms overlapping a common cold.
- Taste and smell disturbances may accompany COVID-19.
- Cough from lower respiratory involvement and fever are additional described features.

Common cold

Upper-airway symptoms

Catarrhal rhinitis

COVID-19

Overlapping UR symptoms

Taste/smell change, cough, fever

Symptom overlap prevents categorical discrimination

- A common cold and COVID-19 can both have upper respiratory manifestations.
- The additional manifestations are not described as absolute criteria.
- The source excerpt does not provide a definitive diagnostic testing algorithm.

A four-way conceptual diagnostic comparison

- Infectious: Acute viral disease with catarrhal discharge.
- Allergic: IgE-mediated disorder with eosinophil-rich response.
- Chronic: Repeated episodes, altered drainage, epithelial injury.
- COVID-19: Overlapping upper respiratory symptoms with additional possible manifestations.

Process	Discriminating emphasis
Acute infectious	Viruses; catarrh; rapid clearance
Allergic	Allergen / IgE / eosinophils
Chronic	Repeated disease; injury
COVID-19	Taste/smell change; cough; fever may occur

Case 1: An uncomplicated acute presentation

- A patient has recent nasal discharge with swollen, red nasal mucosa.
- The turbinates are enlarged and nasal passages narrowed.
- Teaching question: Which morphologic features support acute infectious rhinitis?

Discuss the reasoning before revealing the answer in speaker notes.

Case 2: The exudate changes

- A viral rhinitis episode initially produces profuse catarrh.
- Subsequently the inflammatory reaction intensifies and the exudate becomes mucopurulent.
- Teaching question: What additional process should be considered?

Discuss the reasoning before revealing the answer in speaker notes.

Case 3: Recurrent rather than acute disease

- Episodes recur and secretions drain poorly in the setting of nasal polyps.
- Surface epithelial desquamation and mixed inflammatory cells are described.
- Teaching question: Which adjacent Robbins category better fits this pattern?

Discuss the reasoning before revealing the answer in speaker notes.

Case 4: Shared edema, different mechanism

- A rhinitis episode follows exposure to an allergen.
- The inflammatory tissue infiltrate is eosinophil-rich.
- Teaching question: Which mechanism differentiates this from a cold?

Discuss the reasoning before revealing the answer in speaker notes.

Case 5: Upper respiratory overlap

- A patient has cold-like upper airway symptoms with taste disturbance, cough, and fever.
- These additional findings in COVID-19.
- Teaching question: What differential diagnosis must remain under consideration?

Discuss the reasoning before revealing the answer in speaker notes.


Case 6: Rhinitis precedes a sinus complication

- Rhinitis is followed by impaired sinus drainage.
- Mucosal edema narrows the outflow pathway and sinus infection develops.
- Teaching question: What mechanism connects the two disorders?

Discuss the reasoning before revealing the answer in speaker notes.

Integrating the disease in one causal schema

- Initiation: Several respiratory viruses trigger acute nasal inflammation.
- Local response: Redness, thickening, edema, turbinate enlargement, catarrh.
- Outcome: Rapid clearance is usual; Pharyngeal spread and bacterial superinfection are possible.



Distinguish mechanisms from consequences

- Viral cause is the initiating condition in the common cold.
- Edema and catarrh are inflammatory manifestations, not independent etiologies.
- Pharyngotonsillitis, bacterial superinfection, and sinus involvement are potential extensions.


Level	Example
Cause	Respiratory viruses
Tissue response	Edema, erythema, catarrh
Possible extension	Pharyngotonsillitis / superinfection

Questions to carry into clinical practice

- What feature indicates an acute viral mucosal reaction rather than a chronic epithelial injury pattern?
- What exudate change suggests superimposed bacterial infection?
- Why does similar nasal edema not establish a shared cause between infection and allergy?

Synthesis: What Robbins establishes

- The common cold is a typically rapidly clearing, virus-mediated nasal inflammatory disorder.
- Acute gross changes are red, swollen mucosa, enlarged turbinates, narrowed cavities, and catarrh.
- Recognize secondary bacteria, local extension, and overlap with allergic disease or COVID-19.



Source and interpretation boundaries

- Primary source: Robbins, Cotran & Kumar, Pathologic Basis of Disease, 11th ed. (2026), Chapter 16, Head and Neck.
- Core infectious-rhinitis passage: page 672; comparisons draw on adjacent nasal inflammatory sections (pages 672–673).
- Diagrams are original editable schematics; the supplied source does not include reusable acute-rhinitis photomicrographs.
- Clinical testing thresholds, pathogen assays, and treatment recommendations are outside the source excerpt.