

Malnutrition in High-Resource Countries

Secondary nutritional depletion, inflammation and cachexia

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Distinguish inadequate nutrient supply from inflammation-related malnutrition.
- Recognize characteristic loss of subcutaneous fat and skeletal muscle in adults.
- Connect catabolic mechanisms to adverse clinical outcomes and cachexia.
- Interpret common hospital and long-term-care scenarios using these mechanisms.

Malnutrition persists in resource-rich settings

- High national food availability does not ensure adequate nutrition for an individual patient.
- Secondary malnutrition is described particularly in chronically ill, older and bedridden patients.
- The central clinical question is why tissue stores are being depleted.

HIGH-RESOURCE SETTING



Vulnerable patient




Secondary malnutrition

What makes this primarily secondary malnutrition?

Primary malnutrition	Secondary malnutrition
Deficiency in the diet itself	Inadequate intake, digestion, absorption or use
Food availability may be limiting	Disease can create a mismatch despite available food
Nutrients supplied are insufficient	Supply, losses or requirements may be altered


The pathologic endpoint is tissue loss




- Robbins characterizes malnutrition by loss of fat and muscle mass, weight loss and generalized weakness.

Nutrition has three distinct biological functions

- Carbohydrates, fats and proteins provide energy for daily metabolism.
- Amino acids and fatty acids furnish building blocks for new proteins and lipids.
- Vitamins and minerals support metabolic pathways and structural functions.




A normal diet may still be biologically inadequate



- Secondary malnutrition is not synonymous with starvation or absent food access.


Four ways disease can disrupt nutritional balance

- Malabsorption prevents normal nutrient delivery from the gut.
- Impaired utilization or storage reduces the usefulness of absorbed nutrients.
- Excess losses diminish retained nutrient stores.
- Increased needs can exceed the available dietary supply.



Three high-risk clinical settings

- Chronic disease can create sustained metabolic demand and inflammation.
- Advanced age is an important context for secondary nutritional compromise.
- Prolonged bed rest or hospitalization makes wasting clinically consequential.




Nursing-home residents: Textbook estimate

- Malnutrition in more than half of older nursing-home residents in the United States.
- This illustrates the magnitude of vulnerability in institutional care.
- Treat the figure as a textbook-reported estimate, not a newly verified current prevalence rate.

Older community adults are not exempt

- That approximately one-quarter of older US adults living in the community are malnourished or at risk.
- The reported category includes both established malnutrition and risk of malnutrition.
- A community dwelling alone does not exclude nutritional vulnerability.

Illness changes nutrient requirements




- Failure to recognize illness-related requirements can delay recovery.

A mechanistic differential is more useful than location

Energy / nutrient deficiency	Inflammation-related malnutrition
Inadequate intake or absorption	Disease-associated inflammatory mediators
Example: post-stroke dysphagia	Example: chronic inflammatory disease
Effective nutrient supply falls	Catabolic drive and expenditure rise

Pathway I: Insufficient intake



- Dysphagia following stroke is the explicit clinical example in the chapter.

Post-stroke dysphagia: Clinical logic

Clinical vignette

An older patient has difficulty swallowing following a cerebrovascular event. Progressive weight loss and loss of muscle bulk are observed.


Interpretive question

Which pathway is directly supported?

Interpretation: Energy and nutrient deficiency due to inadequate intake.

- The neurologic disorder limits delivery of food rather than primarily increasing catabolism.
- Concurrent inflammation cannot be judged without additional clinical information.

Pathway I can begin with malabsorption



- Eating does not establish that the intestine can absorb the required nutrients.

Short bowel syndrome: The key distinction

Clinical vignette

A patient with markedly reduced functional small-bowel length continues to eat but loses body mass.

Interpretive question


Why is oral intake an incomplete assessment?

Interpretation: The limiting step is nutrient absorption, not necessarily food consumption.

- Short bowel syndrome as a direct example of energy and nutrient deficiency.


Broader sources of impaired effective supply

- Gastrointestinal disease or malabsorption syndromes can reduce nutrient uptake.
- Some drugs can interfere with nutrient uptake or utilization.
- Abnormal nutrient storage and excessive nutrient loss are additional mechanisms.




High-resource malnutrition is often multifactorial

- Reduced intake, poor absorption, metabolic demand and chronic disease can occur in the same patient.
- A single visible physical sign does not identify the dominant mechanism.
- Etiologic assessment should distinguish supply failure from inflammatory catabolism.




Pathway II: Inflammation-related malnutrition



- This pathway is more common in adults than isolated energy/nutrient deficiency.


Which chronic diseases are implicated?

- Cancer is a major setting for inflammation-related malnutrition.
- Chronic obstructive pulmonary disease is explicitly included.
- Inflammatory bowel disease and renal failure are also listed.



Cancer: Sustained catabolic pressure

- A neoplasm may coexist with an inflammatory, hypercatabolic host response.
- Weight loss need not be explained solely by decreased food intake.
- The related syndrome of cancer cachexia is discussed separately Chapter 7.



COPD: Chronic disease as nutritional stress

Clinical vignette

An older adult with chronic obstructive pulmonary disease loses muscle mass despite reporting regular meals.

Interpretive question


What mechanism should be considered?

Interpretation: Inflammation-related malnutrition, even if intake appears adequate.

- The disease appears' list of chronic inflammatory settings.
- A dietary history alone cannot rule out metabolic tissue loss.


Inflammatory bowel disease: Dual pathways

- Inflammatory bowel disease among chronic disorders associated with inflammation-related malnutrition.
- Gastrointestinal disease may also impair effective nutrient uptake in the broader dietary-insufficiency framework.
- More than one causal pathway may therefore be plausible in an individual case.




Renal failure and inflammatory malnutrition

- Renal failure is another chronic disorder' inflammation-related malnutrition group.
- The provided text links this group to increased metabolic expenditure and skeletal muscle breakdown.
- Do not assume that deficient intake alone explains wasting in this setting.




Inflammatory mediators act upstream




- The direction is disease-associated inflammatory signaling toward heightened metabolic demand.

Resting energy expenditure is important

- Energy is required even without intentional physical exertion.
- Inflammatory mediators can increase this baseline expenditure.
- Sustained demand promotes use of endogenous tissue substrates.




Skeletal muscle becomes a metabolic substrate




- The benefit of substrate supply is purchased at the cost of muscle mass.

Why amino acids are diverted to glucose

- Muscle protein degradation provides amino acid substrates.
- These amino acids to increased glucose production during catabolic stress.
- The result is progressive depletion of functional skeletal muscle.



Adequate protein intake may not prevent wasting




- The chapter explicitly describes skeletal-muscle catabolism even when protein intake is adequate.

Deficient supply versus accelerated breakdown

Supply-limited pathway	Catabolism-driven pathway
Inadequate intake / absorption	Cytokine and prostaglandin signaling
Lower available nutrients	Higher resting expenditure
Depletion due to shortage	Depletion despite apparently adequate supply
Stroke dysphagia / short bowel	Cancer / COPD / IBD / renal failure


One patient can have both mechanisms



- This combined-pathway illustration is a clinical synthesis of the adjacent Robbins sections.

Clinical recognition begins with body composition

- Loss of subcutaneous fat is a conspicuous visible feature.
- Wasting of large skeletal-muscle groups is another major sign.
- Dependent edema may coexist and complicate visual interpretation.



Subcutaneous fat: Look beyond body weight

- Robbins specifically describes fat depletion in the arms and chest wall.
- The shoulder and metacarpal regions are additional sites of visible tissue loss.
- Fat loss is an anatomic finding, not proof of a single underlying cause.

Arms / chest wall



Shoulders



Metacarpal region

Loss of fat in the upper limb

- Inspect the contours of the arms for diminished subcutaneous tissue.
- Reduced soft-tissue covering may make bony and tendinous contours more obvious.
- Interpret these changes together with muscle bulk and clinical context.

Examination sites — schematic


Arms

Shoulders

Metacarpal region


Quadriceps wasting: Lower-extremity clue

- Quadriceps wasting is an explicitly named sign of secondary malnutrition.
- The large muscle group is a useful site for observing loss of skeletal muscle mass.
- Atrophy is consistent with prolonged negative protein balance or catabolic drive.




Deltoid wasting: Shoulder-girdle clue

- Deltoid atrophy provides an additional clinically visible sign.
- Concurrent loss in multiple regions supports systemic tissue depletion.
- The sign should trigger assessment of nutrient supply and inflammatory disease.



Edema can coexist with tissue depletion

- Ankle edema and sacral edema are described among conspicuous clinical findings.
- Fluid accumulation and tissue wasting are different dimensions of the examination.
- Edema should not be used as evidence that nutritional stores are preserved.



Where to look for edema

- Ankle edema is a useful clue in an ambulatory examination.
- Sacral edema may be particularly relevant in a recumbent patient.
- Neither distribution identifies the etiology on its own.

Examination sites — schematic


Ankles

Sacrum

Three physical domains should be integrated

Fat	Muscle	Fluid
Arms, chest, shoulders, hand regions	Quadriceps and deltoids	Ankle or sacral edema
Subcutaneous reserve	Skeletal protein reserve	Edema phenotype
Depletion finding	Wasting finding	Requires separate interpretation


A structured bedside assessment



- This teaching sequence organizes the examination findings listed by Robbins; It is not a validated diagnostic instrument.


Why malnutrition matters in hospital

- The chapter identifies higher risks of infection and sepsis in malnourished hospitalized or bedridden patients.
- Impaired wound healing is another reported consequence.
- Death after surgery is more likely in the malnourished group.




Malnutrition and infection risk

- Malnourished hospitalized patients have increased infection risk.
- Underlying chronic illness and tissue loss can occur simultaneously.
- The source establishes a clinical association; It does not quantify a patient-specific risk.




Sepsis is a serious associated outcome



- Increased sepsis risk; This is not a claim that every infection progresses to sepsis.


Impaired wound healing has practical consequences

- The source associates malnutrition with delayed or impaired wound healing.
- This matters for wounds and for the perioperative patient.
- Assessment of nutrition should be part of understanding poor recovery.



Surgery amplifies clinical importance

- Death after surgery is an increased risk among malnourished hospitalized or bedridden patients.
- The source does not supply an absolute risk or an effect size for the individual patient.
- Nutritional compromise is clinically relevant before and after operative care.




Outcomes are not a diagnostic definition

Clinical signs	Clinical associations
Fat depletion / muscle wasting	Infection and sepsis risk
Ankle or sacral edema	Impaired wound healing
Support suspicion of malnutrition	Explain why recognition matters

Cachexia: An important related syndrome

- Cachexia as secondary malnutrition in patients with advanced cancer or AIDS.
- Its detailed discussion is located in the neoplasia chapter.
- Cachexia particularly emphasizes systemic catabolism and loss of muscle.




Cancer cachexia is not just poor intake


Ordinary food restriction	Cancer cachexia
Primary shortfall in intake	Hypercatabolic state
Intake deficit accounts for tissue loss	Muscle loss not explained by reduced intake alone
Food-supply mechanism central	Inflammatory and tumor-host mechanisms prominent

Phenotype of cancer cachexia

- Profound weight loss, fatigue and muscle atrophy.
- Anorexia, anemia and edema may accompany the wasting.
- Muscle loss can occur with or without adipose loss.




Inflammatory cytokines in cancer cachexia




- These mediators as important contributors, not a complete single-cause explanation.

Protein breakdown: Ubiquitin–proteasome system



- Robbins specifically gives myosin heavy chain as an example of a degraded skeletal-muscle protein.

GDF-15 introduces an appetite pathway



- This pathway complements—not replaces—the inflammatory catabolism model.

Cachexia is mechanistically complex

- Interventions directed at a single cytokine such as TNF have not reversed cancer cachexia in clinical experience.
- GDF-15 / GFRAL antagonism has shown activity in experimental and early clinical settings.
- The textbook explicitly emphasizes incomplete mechanistic understanding.

Multiple contributors



Single-pathway limitation



Active uncertainty

Case: Pneumonia after prolonged bed rest

Clinical vignette

A bedridden older patient has obvious quadriceps wasting and minimal arm subcutaneous fat. The patient is admitted with an infection.

Interpretive question

What additional clinical concern is supported?

Interpretation: Increased risk of infection-related complications and sepsis in malnourished inpatients.

- The physical findings support suspicion of secondary malnutrition.
- The example does not establish which nutritional pathway predominates.

Case: Postoperative wound recovery

Clinical vignette

Following abdominal surgery, a chronically ill patient has poorly healing wounds, loss of deltoid bulk and sacral edema.

Interpretive question

How should these observations be connected?

Interpretation: Malnutrition may coexist with impaired healing and adverse postoperative risk.

- The associations support nutritional assessment, not a claim that malnutrition is the only cause.

Case: Cancer with adequate intake

Clinical vignette

A patient with advanced cancer reports substantial food intake yet continues to lose skeletal muscle mass.

Interpretive question

Why is intake alone insufficient to explain the phenotype?


Interpretation: Inflammation-related hypercatabolism / cancer cachexia can cause ongoing muscle loss.

- Robbins explicitly allows skeletal catabolism even with adequate protein intake.

Case: Distinguish the two pathways

Clinical situation	Dominant mechanism suggested
Dysphagia after stroke	Insufficient intake
Short bowel syndrome	Malabsorption
COPD with wasting despite meals	Inflammation-related catabolism
Advanced cancer with muscle loss	Inflammation / cachexia may contribute

A practical etiologic reasoning sequence




- This is an educational reasoning framework based on the cited textbook mechanisms, not a formal diagnostic guideline.

Common interpretation errors

- Do not equate food availability with adequate tissue nutrition.
- Do not infer normal skeletal protein balance from reported adequate intake.
- Do not interpret edema as an absence of wasting.
- Do not transfer diagnostic thresholds for childhood wasting to older adults.

Integrated clinicopathologic model



- Different upstream pathways converge on clinically important tissue depletion.

Five conclusions for clinical pathology

- Secondary malnutrition is important in older, chronically ill and bedridden patients.
- Differentiate reduced effective supply from inflammation-driven expenditure and catabolism.
- Examine subcutaneous fat, quadriceps, deltoids and dependent edema.
- Complications include infection, sepsis, poor healing and higher postoperative mortality risk.
- Cancer cachexia demonstrates that muscle loss cannot always be corrected by intake alone.

References and source boundaries

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th edition (2026). Chapter 9: Environmental and Nutritional Diseases; dietary insufficiency, adult malnutrition and cachexia sections.
- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th edition (2026). Chapter 7: Neoplasia; Cancer Cachexia.
- Clinical vignettes, pathway figures and comparison tables in this lecture are original teaching reconstructions—not published patient data or diagnostic criteria.