

Mycobacterial Arthritis

Tuberculous synovitis · granulomatous injury · chronic joint destruction

Infection → granuloma → pannus → ankylosis

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraisakhri.com

Learning objectives

- Trace hematogenous and contiguous routes of mycobacterial joint infection.
- Explain how tuberculosis immunopathogenesis produces caseating granulomas and articular destruction.
- Connect synovial pannus, marginal erosion and fibrous ankylosis to chronic symptoms.
- Distinguish this process from selected alternative arthritides and interpret TB tests appropriately.


Tuberculous arthritis is a chronic joint infection

- Mycobacterial arthritis in the Robbins section refers to joint infection caused by *Mycobacterium tuberculosis*.
- The disorder is chronic and progressive rather than characteristically abrupt.
- Synovial granulomatous inflammation drives cartilage and bone injury.
- Late disease can obliterate the joint space and produce fibrous ankylosis.

Its place among infectious arthritides

Dimension	Tuberculous arthritis	Acute suppurative arthritis
Typical tempo	Insidious, progressive	Abrupt, acutely painful
Principal lesion	Caseating granulomas; pannus	Acute purulent infection
Long-term hazard	Erosion, fibrosis, ankylosis	Rapid destructive joint injury
Clinical priority	Recognize gradual progression	Urgent identification and therapy

Why an infected joint can suffer permanent damage



1. Articular cartilage has less regenerative capacity than bone.
2. The tempo differs by pathogen, but irreversible structural damage is a shared concern.

Two source pathways reach the joint


Hematogenous spread

1. A visceral source provides circulating bacilli.
2. Pulmonary tuberculosis is the usual visceral source noted by Robbins.
3. Organisms seed the joint.

Contiguous spread

1. Tuberculous osteomyelitis is present nearby.
2. Infection extends from an osseous focus.
3. Joint structures become involved.

Hematogenous dissemination: Source to synovium




1. A pulmonary focus is usually implicated when a visceral source is identifiable.
2. A joint manifestation may appear after systemic dissemination.

Why a lung focus is relevant

- Tuberculosis is acquired predominantly by inhalation from people with active pulmonary disease.
- Organisms can disseminate hematogenously from primary or later disease.
- An articular presentation therefore prompts consideration of a visceral source.
- Pulmonary symptoms are not a required feature of the articular definition.


Early bacteremia creates a route to distant sites



1. Early dissemination during the initial immune response.
2. Joint infection is one potential skeletal consequence of dissemination.

Mechanistic context from the broader tuberculosis chapter.

Contiguous extension begins in infected bone




1. The parent section specifically identifies adjacent osteomyelitis as a route.
2. The anatomic relationship explains why joint and bone findings may coexist.

Bone tuberculosis supplies the neighboring focus

- Mycobacterial osteomyelitis typically arises from bloodborne seeding of bone.
- Osseous disease may remain unrecognized for a prolonged period.
- Caseating granulomas are a shared morphologic lesion in bone and joint infection.
- The bone section explicitly lists tuberculous arthritis as a possible complication.

Adjacent bone and synovium: Spatial relationship




Simplified section of an articular interface (schematic; Not a diagnostic image).

Host factors may shape dissemination

- Tuberculosis is more likely to progress when cellular immune defenses are weakened.
- HIV infection, malnutrition and immunosuppression among important TB risk contexts.
- The osteomyelitis section describes more extensive infection in immunocompromised patients.
- These are tuberculosis background factors—not joint-specific incidence estimates.

Macrophage uptake initiates mycobacterial infection



1. Uptake through receptors including mannose-binding lectin and CR3.
2. Intracellular survival contributes to persistence before effective T-cell immunity develops.

Phagosome maturation is blocked


Normal antimicrobial pathway

1. Phagosome matures.
2. Fusion with lysosome supports killing.
3. Microbicidal conditions destroy ingested microbes.

Tuberculosis evasion


1. M. Tuberculosis interferes with maturation.
2. Phagolysosome formation is inhibited.
3. The bacillus survives in its intracellular compartment.

Antigen presentation recruits adaptive immunity




1. Mycobacterial antigens are presented in draining lymphoid sites.
2. Cell-mediated immunity is central to the resulting granulomatous response.

IL-12 and IL-18 favor a Th1 response



1. This is the broader Robbins TB mechanism, not a joint-specific cytokine measurement.
2. The pathway provides the immunologic explanation for chronic granuloma formation.

IFN- γ activates infected macrophages



1. Macrophages use reactive species and antimicrobial pathways to control infection.
2. The same inflammatory environment can contribute to tissue destruction.

Protective immunity and tissue injury coexist


Protective consequence

1. Activated macrophages contain bacilli.
2. Granulomas restrict spread.
3. Cell-mediated responses are central to control.

Pathologic consequence

1. Persistent inflammation injures tissue.
2. Caseous necrosis can develop.
3. Destruction around a joint compromises function.

Granuloma formation: A tissue-level sequence




1. Some activated macrophages fuse into multinucleated giant cells.
2. A granuloma expresses containment and a persistent inflammatory focus.

Caseation is the necrotic center of the lesion


- Robbins specifically describes confluent caseating granulomas after seeding of the joint.
- Caseous necrosis accompanies the tissue-damaging component of the TB immune response.
- Epithelioid macrophages and surrounding inflammatory cells form the granulomatous architecture.
- Histologic identification must be interpreted with organism-focused evidence.

Anatomy of a caseating granuloma




Original conceptual granuloma schematic — not histology or evidence of a specific patient.

Confluence expands the area of synovial injury



1. Confluent caseating granulomas are explicitly identified in the joint section.
2. The diagram is a conceptual lesion sequence, not a measured timeline.

TNF and chemokines sustain cellular recruitment



1. This feedback is described in general tuberculosis pathogenesis.
2. It helps explain how an initially localized infection can sustain chronic lesions.


Immunosuppression can alter containment

- Macrophage activation and Th1 responses help restrain M. Tuberculosis.
- Loss of effective immune control can permit persistent or disseminated infection.
- Reactivation risk during TNF-antagonist treatment.
- Do not equate immunosuppression with a specific articular morphology or frequency.

The synovium becomes an agent of destruction

- Granulomatous inflammation involves the synovium.
- Inflammatory synovial tissue can extend over the articular surface.
- That advancing tissue as a pannus.
- Pannus creates the anatomical route to cartilage and marginal bone injury.

Pannus extends across articular cartilage




Simplified section of an articular interface (schematic; Not a diagnostic image).

Cartilage damage follows direct surface encroachment

- Pannus advances over the articular cartilage.
- Chronic inflammation and direct encroachment damage the cartilage surface.
- Loss of cartilage compromises smooth articulation and joint integrity.
- Because cartilage repairs poorly, structural loss may persist.

Marginal erosion follows pannus invasion




Simplified section of an articular interface (schematic; Not a diagnostic image).

Caseation and erosion are distinct observations

Feature	Primary meaning	Location / relationship
Caseating granuloma	TB-associated granulomatous lesion	Inflamed synovium
Pannus	Extending inflammatory synovial tissue	Over cartilage
Marginal erosion	Articular bone loss	Joint edge invaded by pannus
Ankylosis	Late loss of mobility	Fibrous bridging of articular structures

Joint-space obliteration is a late structural outcome



1. The passage identifies obliteration as a consequence of chronic disease.
2. The diagram deliberately avoids assigning unsupported imaging measurements.

Fibrous ankylosis limits motion

- Fibrous ankylosis is the late fusion process described in the Robbins section.
- Fibrous tissue restricts normal separation and movement of articular surfaces.
- Ankylosis is the endpoint of sustained structural damage—not an early diagnostic requirement.
- It should not be confused with a claim of obligatory bony fusion.


The lesion evolves through recognizable stages

Mycobacterial seeding

Granulomatous synovitis


Pannus / marginal erosion

Fibrous ankylosis



Sequence shows pathologic relationships; It does not assign fixed stages or timings.

Pathogenesis links infection to disability



1. These are the causal anchor points explicitly identified in the joint passage.
2. The clinical correlate is a progressive loss of normal joint function.

Weight-bearing joints predominate

Hip



Knee



Ankle

Emphasized first

Second in order

Third in order

Source gives rank order but no site-specific percentages.

Hip involvement: A weight-bearing articulation

- The hip is listed first among commonly affected joints.
- Hip involvement exemplifies the vulnerability of a major weight-bearing articulation.
- Insidious pain should be interpreted alongside the possibility of a chronic destructive joint process.
- Site alone is not diagnostic of tuberculous infection.

Knee involvement: Another frequent site

- The knee follows the hip in the order reported by Robbins.
- Inflammatory synovial tissue may affect a large articular surface.
- Progressive cartilage and marginal bone injury can threaten function.
- The joint-site distribution supports suspicion but cannot establish etiology.


Ankle involvement: The third listed site

- The ankle is the third weight-bearing joint highlighted.
- The same granulomatous synovial mechanism applies regardless of the joint involved.
- Chronic pain plus structural damage warrants an infectious differential.
- Avoid interpreting the ordered list as an exhaustive distribution of disease.

Insidious onset shapes the clinical presentation

- The disease does not characteristically begin with a sudden explosive episode.
- Gradually increasing pain is the central symptom specified in the joint section.
- Its chronic nature can delay recognition when joint pain has multiple possible causes.
- Absence of abrupt symptoms does not exclude an infectious process.

Pain progression mirrors tissue progression



1. Clinical and morphologic events are linked conceptually.
2. Robbins explicitly identifies increasing pain, pannus, erosion and late ankylosis.

This is a mechanism-based synthesis, not a validated time-to-symptom curve.

Systemic clues come from skeletal TB context

- The mycobacterial osteomyelitis discussion notes localized pain and possible low-grade fever.
- Chills or weight loss can occur in bone tuberculosis.
- These features provide contextual clues for osseous or systemic disease.
- The dedicated arthritis section does not specify their frequency in joint infection.

Clinical and pathologic clues reinforce one another

Clinical or structural observation	Pathologic explanation
Slowly worsening pain	Persistent granulomatous synovitis
Adjacent osseous disease	Potential contiguous source
Articular marginal injury	Pannus-associated erosion
Severe stiffness in chronic disease	Fibrous ankylosis / loss of joint space

Early and advanced disease must be distinguished

Earlier process

1. Mycobacterial seeding.
2. Granulomatous synovitis.
3. Insidious pain.

Advanced process

1. Pannus and marginal erosion.
2. Obliteration of joint space.
3. Fibrous ankylosis.

Suppurative arthritis differs in its tempo

Finding	Mycobacterial arthritis	Suppurative arthritis
Onset	Insidious	Sudden
Pain course	Gradually increasing	Acute, intensely painful joint
Primary inflammatory picture	Chronic caseating granulomas	Purulent fluid / acute infection
Consequences	Erosion and ankylosis	Potential rapid joint destruction

Rheumatoid arthritis also produces pannus

Feature	Tuberculous arthritis	Rheumatoid arthritis
Underlying process	M. tuberculosis infection	Immune-mediated inflammatory arthritis
Synovial lesion	Caseating granulomas may occur	Chronic inflammatory synovitis
Cartilage / bone injury	Pannus and marginal erosion	Pannus with erosive changes
Key distinction	Mycobacterial evidence	Joint pattern and immune context

Osteoarthritis causes cartilage injury differently

Feature	Tuberculous arthritis	Osteoarthritis
Driver	Infectious granulomatous inflammation	Mechanical / degenerative cartilage injury
Synovial pathology	Granulomas and pannus	Usually limited secondary inflammation
Bone changes	Inflammatory marginal erosion	Osteophytes; subchondral changes
Typical interpretation	Consider infection	Noninfectious degenerative pattern


Viral arthritis has a different time course

- Viral arthritis is generally acute to subacute.
- Joint symptoms can reflect direct viral infection or triggered autoimmunity.
- This differs from the chronic progressive caseating granulomatous process of tuberculosis.
- Tempo helps prioritize causes but is not a stand-alone microbiologic test.

Lyme arthritis is a separate infectious pattern

- Lyme arthritis is caused by *Borrelia burgdorferi* rather than *M. Tuberculosis*.
- It often involves large joints, particularly the knee.
- The Robbins section describes attacks that can persist weeks to months and may move to new sites.
- Shared large-joint involvement does not establish the same mechanism or diagnostic test.

Pattern recognition narrows—not closes—the differential



Diagnostic evidence needs both tissue and organism context

- The characteristic synovial lesion is a caseating granuloma.
- Granulomatous morphology strongly frames the differential but does not by itself establish the species.
- Robbins recommends biopsy and microbial culture for many osteomyelitis diagnoses.
- An articular workup should seek appropriate microbiologic confirmation; The joint-specific specimen pathway is a clinical extrapolation.

Acid-fast staining examines the bacterial phenotype

- The lipid-rich mycobacterial cell wall underlies acid-fast staining.
- Staining demonstrates acid-fast organisms if present at detectable abundance.
- Acid-fast morphology alone does not uniquely identify M. Tuberculosis.
- A negative smear does not settle the cause of a granulomatous joint lesion.

Detection caveats are general diagnostic reasoning, not sensitivity claims for synovial specimens.

Culture identifies organisms and supports susceptibility testing

- Robbins calls culture the gold standard for active tuberculosis diagnosis.
- The tuberculosis chapter describes slow growth on solid culture media.
- Culture supports comprehensive antimicrobial susceptibility testing.
- Reported pulmonary culture timings should not be presented as joint-fluid performance data.

Nucleic-acid amplification provides earlier TB evidence

- PCR detection of M. Tuberculosis DNA as more rapid than culture.
- Some assays also evaluate resistance-associated markers.
- Performance varies with specimen and bacterial burden.
- The chapter's pulmonary-test sensitivity statements must not be transferred to joints without evidence.


IGRA and tuberculin tests do not prove active arthritis

Test group	What it measures	Critical interpretive limit
IGRA	Antigen-specific T-cell IFN- γ response	Does not distinguish latent from active TB
Tuberculin skin test	Cell-mediated skin response to PPD	Positive result does not localize active disease
Culture / NAAT	Direct organism-oriented evidence	Test interpretation depends on specimen and context

Diagnostic interpretation must avoid four errors

- Do not diagnose active joint tuberculosis from an IGRAs or skin test alone.
- Do not identify M. Tuberculosis from an acid-fast morphology pattern alone.
- Do not assume every erosive pannus is rheumatoid arthritis.
- Do not assume a slow clinical course excludes infection.

Integrate clinical, histologic and microbiologic information



1. Interpret results in the context of disease location and possible visceral / bone source.
2. This is an educational framework, not a replacement for local clinical protocols.

Framework synthesized from source sections; Not a formal Robbins algorithm.

Long-standing lesions threaten joint function

- Pannus expansion damages cartilage and marginal bone.
- Chronic infection may obliterate the joint space.
- Fibrous ankylosis restricts mobility and can leave lasting functional impairment.
- The joint section does not provide a numerical complication rate.

Clinical exercise: Gradually painful hip

An adult has slowly progressive hip pain. Tissue from a destructive synovial lesion shows confluent caseating granulomas.

Which pathogen class and morphologic sequence should be prioritized?

Clinical clue 1: Insidious pain

Clinical clue 2: Weight-bearing joint

Clinical clue 3: Caseating granulomatous synovitis

Case analysis: Hip lesion

- Prioritize tuberculous arthritis due to the chronic course and caseating granulomas.
- Link joint seeding to granulomatous synovitis and advancing pannus.
- Anticipate marginal bone erosion and, if longstanding, fibrous ankylosis.
- Microbiologic evidence is needed to identify the causative mycobacterium.

Hypothetical educational case, not a reported patient.

Clinical exercise: Inflammatory pannus

Synovial tissue forms pannus and has eroded joint margins. A discussion proposes rheumatoid arthritis solely because pannus is present.

What important etiologic alternative and tissue finding must be considered?

Clinical clue 1: Pannus is not disease-specific

Clinical clue 2: Marginal erosions occur in several disorders


Clinical clue 3: Morphology and microbiology are discriminators

Case analysis: Pannus is not a diagnosis

- Pannus occurs in tuberculous arthritis as well as rheumatoid arthritis.
- Confluent caseating granulomas strongly raise tuberculosis in this specific context.
- Clinical time course and source of infection help structure the differential.
- Investigate for mycobacteria rather than assigning an autoimmune cause from pannus alone.

Hypothetical educational case, not a reported patient.

The core morphologic chain



1. Organism, immune response, tissue response and structural consequence form one causal sequence.
2. Each component addresses a distinct diagnostic or teaching question.

Distinguishing observations to retain

- Primarily adult disease with insidious and progressive pain.
- Often arises through bloodborne dissemination or adjacent osteomyelitis.
- Caseating granulomas, synovial pannus and marginal erosions are key lesions.
- Hip, knee and ankle are the emphasized weight-bearing sites; Chronic disease can culminate in fibrous ankylosis.

Primary reference and companion sections

Robbins, Cotran & Kumar: Pathologic Basis of Disease — Mycobacterial Arthritis.

Companion material: Mycobacterial Osteomyelitis; Infectious and Suppurative Arthritis.

Background framework: Tuberculosis—pathogenesis and diagnostic testing.

Contrasts: Rheumatoid Arthritis, Osteoarthritis, Viral Arthritis and Lyme Arthritis.