

Nonimmune Hydrops

Etiology · pathogenesis · morphology · clinicopathologic reasoning

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives


- Distinguish nonimmune hydrops from immune-mediated fetal hemolysis.
- Explain how cardiac, hematologic and lymphatic abnormalities produce fetal fluid accumulation.
- Identify major etiologic groups and their characteristic pathology.
- Integrate morphology with the likely underlying fetal disorder.

Fetal hydrops: The central concept

Abnormal fluid accumulation during intrauterine growth

- The distribution and severity of edema vary widely.
- A shared morphologic endpoint may arise from different diseases.
- Hydrops is a manifestation—not a single etiologic diagnosis.

Severity forms a spectrum



The most severe generalized form may be lethal; Lesser forms can be compatible with life.

Fluid may accumulate in distinct sites

Soft tissues

Generalized
subcutaneous edema;
Prominent neck swelling
may occur


Pleural / peritoneal spaces

Isolated effusions can
represent lesser degrees
of fluid accumulation

Postnuchal region

Cystic hygroma reflects
localized fluid
accumulation, often
lymphatic

Why nonimmune hydrops predominates



Relative frequency changed after effective prevention of Rh alloimmunization.

Two pathogenetic categories

Immune hydrops

- Maternal red-cell alloantibodies cross the placenta
- Fetal erythrocyte destruction is central

Nonimmune hydrops

- No requirement for maternal anti-red-cell antibodies
- Multiple cardiovascular, genetic, hematologic or placental causes

Nonimmune hydrops is a syndrome

Different initiating lesions converge on fetal edema


- A structural cardiac defect and a severe fetal anemia can share the same gross appearance.
- The mechanism must be traced backward from fluid accumulation to its cause.
- Morphology can provide clues but does not establish etiology alone.

The balance governing fetal edema




Hydrostatic pressure, oncotic pressure and lymphatic return influence the final distribution of fluid.

Elevated venous pressure promotes edema




Cardiac failure to raised hydrostatic pressure in fetal hydrops.

Reduced oncotic pressure promotes edema




A contributor especially when anemia causes hepatic injury.

Anemia also stresses the fetal heart



Anemia is a systemic oxygen-delivery problem, not solely a blood-count abnormality.

Lymphatic dysfunction: A distinct route



Particularly relevant to 45,X / Turner syndrome.

Different mechanisms may coexist

Mechanism	Illustrative disorder	Immediate consequence
Hydrostatic pressure ↑	Cardiac malformation / arrhythmia	Congestion
Oncotic pressure ↓	Severe anemia with liver injury	Reduced plasma-protein support
Lymphatic drainage ↓	Turner syndrome	Postnuchal fluid accumulation

Three major etiologic groups

Cardiovascular

Congenital structural lesions and functional rhythm disturbances

Chromosomal

45,X and trisomies 21/18, often with cardiac anomalies

Fetal anemia

Severe anemia unrelated to Rh or ABO maternal antibodies

Cardiovascular defects: Two routes


Structural

- Congenital cardiac malformations
- Abnormal intracardiac anatomy compromises fetal circulation

Functional


- Fetal arrhythmias
- An abnormal rhythm can impair effective cardiac output

Structural cardiac malformation



Congenital cardiac malformations as major structural causes.

Arrhythmia can cause hydrops



Functional cardiac defects do not require an anatomic malformation.

Why a normal-looking heart is insufficient

- Hydrops may result from functional cardiac disease, including arrhythmia.
- Structural assessment alone cannot exclude all cardiac mechanisms.
- The etiologic reasoning must consider both morphology and function.

The cardiovascular final pathway

Heart failure → venous congestion → fetal edema

- Both structural lesions and arrhythmias can initiate this pathway.
- Concurrent anemia can aggravate myocardial oxygen deficiency.
- The visible hydrops is downstream of the cardiovascular disorder.

Chromosomal causes: Key examples

45,X

Turner syndrome; Cardiac
and lymphatic
abnormalities

Trisomy 21

Hydrops associated with
accompanying cardiac
abnormalities

Trisomy 18

Hydrops associated with
accompanying cardiac
abnormalities

45,X: Two mechanisms in one fetus


Cardiovascular component

- Associated structural cardiac anomalies
- Potential intrauterine cardiac failure

Lymphatic component

- Abnormal drainage from the fetal neck
- Postnuchal cystic hygroma

Turner syndrome and cystic hygroma




A useful etiologic clue; Not exclusive to 45,X.

Cystic hygroma is not pathognomonic

- Prominent posterior neck fluid may accompany a 45,X karyotype.
- That cystic hygromas also occur outside Turner syndrome.
- Interpret the finding with other malformations and the overall phenotype.

Cardiac anomalies link aneuploidy to hydrops



Structural cardiac defects as the link.

Dysmorphism directs etiologic reasoning

- Dysmorphic features in a hydropic fetus suggest possible chromosomal disease.
- Morphologic suspicion is a clue, not a definitive chromosome diagnosis.
- Postmortem study may disclose previously unrecognized cardiac abnormalities.

Chromosomal disease: Multiple phenotypes


Finding	Pathogenetic interpretation
Cardiac malformation	Potential circulatory failure
Posterior neck fluid	Possible lymphatic-drainage disturbance
Dysmorphic features	Consider underlying constitutional chromosome anomaly

Fetal anemia: A nonimmune mechanism

Marked oxygen-delivery failure can culminate in hydrops

- The anemia is not caused by maternal Rh or ABO antibodies.
- Severe anemia can damage heart and liver through hypoxia.
- Hepatic protein synthesis and cardiovascular function may both deteriorate.

Alpha-globin gene dosage




Each normal alpha-globin gene contributes to chain synthesis.

Alpha-thalassemia severity tracks deletions


α -gene deletions	Syndrome	Relation to hydrops
1	Silent carrier	Not the severe fetal phenotype
2	α -thalassemia trait	Mild, often microcytosis
3	HbH disease	Significant anemia
4	α -thalassemia major	Severe fetal anemia / hydrops

Four deleted α genes: α -thalassemia major




Alpha-thalassemia major as a major nonimmune hydrops cause in parts of Southeast Asia.

How hemoglobin Bart's forms



Hb Bart's is a gamma-chain homotetramer.

Hb Bart's binds oxygen too tightly



The central defect is impaired unloading of oxygen, not merely hemoglobin concentration.

Fetal oxygen transport collapses

Oxygen carriage does not guarantee effective oxygen delivery

- Hemoglobin Bart's retains oxygen with unusually high affinity.
- Severe tissue hypoxia damages the heart and other fetal organs.
- Fetal edema and organ enlargement may follow.

Clinical morphology in α -thalassemia major

Severe pallor

Reflects profound fetal anemia

Generalized edema

Severe nonimmune hydrops phenotype

Massive hepatosplenomegaly

Erythropoietic demand and cardiovascular consequences

α -Thalassemia major: Natural history

Before fetal interventions

- Severe tissue anoxia often caused intrauterine or early neonatal death


With intrauterine transfusion

- Some affected fetuses can survive;
Ongoing transfusion needs may follow

Long-term consequence after survival

- Survivors of α -thalassemia major often remain transfusion dependent.
- Repeated transfusions introduce clinically important iron-overload risk.
- Hematopoietic stem-cell transplantation as potentially curative.

Parvovirus B19 enters transplacentally




Erythroid tropism is central

Parvovirus B19 preferentially infects erythroid precursors

- The virus replicates in normoblasts.
- Infected red-cell progenitors undergo apoptosis.
- The resulting red-cell aplasia can produce severe fetal anemia.

Parvovirus B19: Marrow sequence



Mechanism specified by Robbins; Not generalized marrow hyperplasia.

Parvoviral inclusions are a tissue clue

- Intranuclear inclusions in circulating or marrow erythroid precursors.
- Inclusion-bearing erythroblasts directly support infection of the red-cell lineage.
- This differs from the usual compensatory erythroid expansion in other anemic causes.

The crucial marrow exception

Most severe fetal anemias

- Compensatory erythroid hyperplasia
- Extramedullary hematopoiesis may expand

Parvovirus B19 fetal anemia

- Direct progenitor destruction
- Red-cell aplasia despite anemia

Parvovirus pregnancy outcomes vary

Most infected pregnancies

- The source notes that the great majority have a normal outcome

Potential adverse outcomes

- Fetal anemia, hydrops, spontaneous abortion or stillbirth

Timing of transplacental infection matters

- Transplacental infections can occur during pregnancy.
- Gestational timing affects clinical manifestations.
- Parvovirus-associated adverse outcomes include second-trimester pregnancy loss.

Two fetal anemia pathways compared

Feature	α -thalassemia major	Parvovirus B19
Primary lesion	Four α -gene deletions	Erythroid progenitor infection
Mechanism	Hb Bart's / severe tissue hypoxia	Red-cell precursor apoptosis
Marrow response	Compensatory erythropoiesis	Aplastic suppression is a key exception

Twin hydrops: Start with the placenta

Shared placental vascular channels can redistribute fetal blood

- Monochorionic twin placentas contain vascular anastomoses.
- Some shunts preferentially move blood between circulations.
- Severe imbalanced perfusion may threaten one or both twins.

Twin placentation determines risk


Dichorionic placentation

- Two chorionic units; May occur in dizygotic or monozygotic twins


Monochorionic placentation

- Implies monozygotic twins; Vessels can connect circulations


Placental vascular anastomoses



Twin-to-twin transfusion: Donor side



Twin-to-twin transfusion: Recipient side



Twin-to-twin transfusion is not symmetric

Donor twin

- Relatively underperfused
- Lower circulatory blood delivery
- At risk from perfusion insufficiency

Recipient twin

- Relatively volume overloaded
- Greater circulatory load
- At risk from volume-related cardiac strain

Twin-to-twin transfusion and fetal loss

- Severe intertwin shunting may cause death of one or both fetuses.
- Approximately 10% of nonimmune hydrops cases are linked to monozygotic twin pregnancies account.
- Twin transfusion represents a placental vascular—not red-cell alloimmune—mechanism.

An unexplained subset remains


The cause is not identifiable in every hydropic fetus

- Up to 20% of cases without an established cause.
- Unexplained hydrops should not be assigned a specific molecular diagnosis by appearance alone.
- This is a limit of etiologic classification, not an additional disease category.

Gross pathology: External fluid


- The degree of intrauterine edema varies from localized to generalized.
- Generalized anasarca produces a markedly edematous fetus.
- Posterior neck fluid is especially prominent in some lymphatic disorders.

Gross pathology: Pallor of fetus and placenta




Anemia-associated hydrops shows characteristic fetal and placental pallor.

Hepatosplenomegaly in anemic hydrops



Visceral enlargement in most cases partly to cardiac failure/congestion.

Bone marrow: Compensatory response



This response is expected except with direct suppression such as parvovirus.

Extramedullary hematopoiesis

Liver

Erythroid cell islands
among hepatocytes


Spleen / nodes

Additional sites of
hematopoietic expansion

Other tissues

Kidneys, lungs and even
myocardium may be
involved

Why extramedullary hematopoiesis develops



Compensatory marrow and extramedullary erythropoiesis.

Erythroblastosis fetalis is a blood finding

- Numerous immature red cells can enter fetal peripheral circulation.
- These cells include reticulocytes and erythroblasts.
- The finding is a response to marked anemia; It is not limited to immune hydrops.

Anemia-induced hydrops: Integrated morphology

Compartment	Typical pathologic finding	Interpretation
Fetus / placenta	Pallor	Severe anemia
Liver and spleen	Enlargement	Congestion and hematopoiesis
Bone marrow	Erythroid hyperplasia	Compensation for red-cell deficit
Peripheral blood	Reticulocytes / erythroblasts	Accelerated erythropoiesis

Distinguish marrow hyperplasia from aplasia

Compensatory phenotype

- Expanded erythroid precursors
- Often extranodal / extramedullary hematopoiesis

Parvovirus phenotype

- Infected erythroid precursors undergo apoptosis
- Production failure may dominate

Outcome depends on severity and cause

Severe phenotype

Stillbirth or death shortly
after birth may occur

Variable burden

Some infants have lesser
or localized collections

Potential recovery

Some infants recover
completely


Immune versus nonimmune: Causal contrast

Feature	Immune hydrops	Nonimmune hydrops
Primary process	Maternal anti-fetal red-cell antibody	Diverse non-antibody disorders
Prominent drivers	Rh/ABO incompatibility	Cardiac, chromosomal, fetal anemia
Potential specific findings	Hemolysis; Neonatal hyperbilirubinemia	Cystic hygroma, arrhythmia, genetic anemia, aplasia

Kernicterus: Maintain causal precision

- Bilirubin-mediated CNS injury is a major threat in immune hemolytic disease.
- The mechanism requires severe unconjugated hyperbilirubinemia.
- Hydrops itself does not imply that kernicterus is present.

Interpretation begins with the mechanism



Case: Fetal arrhythmia with edema

Clinical vignette

A fetus has generalized edema and a sustained rhythm abnormality. No structural malformation is demonstrated.

Which direct pathway explains fluid accumulation?

Case: Posterior neck fluid and 45,X

Clinical vignette

A fetus with a 45,X karyotype has marked postnuchal fluid accumulation.

What mechanism most directly explains the neck finding?

Case: Four α -globin gene deletions

Clinical vignette

A hydropic fetus has severe pallor, massive hepatosplenomegaly, and deletion of all four α -globin genes.

Why is fetal oxygen delivery profoundly impaired?

Case: Anemia without erythroid hyperplasia

Clinical vignette

A hydropic fetus is severely anemic. Erythroid precursors contain intranuclear inclusions and erythropoiesis is suppressed.

Which infectious mechanism fits the pattern?

Case: Discordant monochorionic twins

Clinical vignette

Two monochorionic twins have markedly discordant perfusion; One is underperfused and the other volume overloaded.

What placental lesion provides the mechanism?

Causal comparison across etiologies

Etiology	Primary disruption	Shared endpoint
Arrhythmia / malformation	Cardiac failure	Hydrops
45,X / aneuploidy	Cardiac defect $\pm$ lymphatic failure	Hydrops / cystic hygroma
α -Thalassemia major	Oxygen-delivery failure	Hydrops
Parvovirus B19	Erythroid aplasia	Severe anemia $\rightarrow$ hydrops
Twin transfusion	Unequal blood distribution	Overload / circulatory dysfunction

Pathology principles to retain

Mechanism

Hydrops is a final common consequence of several distinct disturbances

Morphology

Pallor, organ enlargement, erythroid response and edema reveal key physiology

Etiology

Cardiac, chromosomal, anemic, viral and twin-related causes demand distinction

Primary textbook sources

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. (2026). Chapter 10: Fetal Hydrops, pp. 419–422.
- Chapter 14: α -Thalassemia / α -Thalassemia Major.
- Chapter 22: Twin Placentas; twin-to-twin transfusion syndrome.
- Chapter 5: Turner syndrome (supplementary genetic context).