
Other Orbital Inflammatory Conditions

Anatomic pathways · inflammatory morphology · differential diagnosis

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning objectives

- Explain how orbital anatomy permits spread of inflammation from adjacent paranasal sinuses.
- Distinguish infectious, systemic, idiopathic, and neoplastic causes of orbital enlargement.
- Interpret the major histologic findings of idiopathic orbital inflammation and their diagnostic implications.
- Recognize when granulomatosis with polyangiitis or IgG4-related disease must be considered.

Orbital inflammation is an anatomic and etiologic problem


Orbital setting	Interpretive consequence
Paranasal sinuses next to orbit	Adjacent infection may extend across shared bony boundaries
Muscles, lacrimal gland, fat, fascial tissue	Different compartments can be selectively involved
Eye in a limited bony cavity	Increased orbital contents displace the globe
Systemic immune-mediated disease	Orbital disease may be the first recognized manifestation

Orbital walls explain routes of sinus extension



Original anatomic schematic — not a clinical scan

The anatomic route to infectious orbital inflammation



Sinus disease is an anatomic source—not evidence that an orbital lesion is idiopathic.


Proptosis is a manifestation—not a diagnosis

- Expansion of orbital contents displaces the globe forward.
- An orbital mass, inflammatory swelling, or tissue accumulation can produce the same outward sign.
- The direction of displacement may help localize the underlying lesion.
- Proptosis can threaten the corneal surface through incomplete lid coverage.

Axial versus non-axial proptosis


Feature	Axial	Non-axial
Globe position	Displaced directly forward	Displaced forward with directional deviation
Anatomic implication	Often centered in the muscle cone	Suggests a more eccentric orbital process
Robbins example	Optic nerve glioma or meningioma	Superotemporal lacrimal gland enlargement

The muscle cone explains axial proptosis




Original anatomic schematic — not a clinical scan

Lacrimal gland enlargement shifts the eye inferomedially



Original anatomic schematic — not a clinical scan

How proptosis leads to corneal injury



These consequences arise from exposure and do not identify the underlying etiology.

Thyroid-associated orbitopathy provides a key comparator

- Graves-associated orbitopathy is an important cause of axial proptosis.
- Rectus muscles accumulate extracellular matrix proteins and develop variable fibrosis.
- Orbital manifestations may not parallel the contemporaneous thyroid function status.
- The tendon-sparing pattern is emphasized in the Robbins gross example.

Thyroid orbitopathy versus inflammatory pseudotumor

Question	Thyroid-associated orbitopathy	Idiopathic orbital inflammation
Primary tissue change	Rectus muscle matrix expansion / fibrosis	Inflammation and variable fibrosis
Distribution in source	Rectus muscles; Tendon sparing in figure	Diffuse or confined to selected compartments
Classification	Associated with Graves disease	No identified local or systemic cause
Pathology implication	Muscular remodeling	Mixed chronic inflammatory infiltrate may be found

Four major etiologic routes to orbital inflammation

Adjacent infection

Systemic
inflammation

IgG4-associated
fibroinflammation

Idiopathic orbital
inflammation

Do not assign the idiopathic label until identifiable local and systemic causes are excluded.

Bacterial and fungal infections can spread from the sinuses

- The ethmoidal sinuses lie medial to the orbit, separated by the lamina papyracea.
- The maxillary sinus lies directly beneath the orbital floor.
- Uncontrolled bacterial or fungal sinus infection may involve orbital structures.
- The adjacent anatomic relationship explains a contiguous rather than primary orbital origin.


Host factors increase concern for infectious extension

- Immunosuppression and diabetes as common contexts.
- Occasional cases arise without a recognized predisposing condition.
- Risk factors alter the clinical level of suspicion; Their absence does not exclude spread.
- An orbital inflammatory mass should not automatically be treated as noninfectious.

Do not equate orbital inflammation with idiopathic disease

Finding or context	Why it changes the differential
Active bacterial or fungal sinus disease	Favors infection as a potential source
Immunosuppression or diabetes	Raises concern for infectious spread
Sinus-based and orbital abnormalities	Suggests contiguous extension
No recognized risk factors	Does not by itself eliminate infection

Sinus-based disease: Causal interpretation



Mechanism schematic: This sequence does not specify microbiologic species or clinical tests.

Applied case: An orbital swelling beside sinus disease

A patient with diabetes develops orbital swelling while an adjacent paranasal sinus infection is active.

Diagnostic question

Which etiologic category must be considered before calling this an idiopathic orbital process?

Contiguous anatomic boundary · Established infection · Predisposing host context

Case interpretation: Infection is the priority exclusion

Contiguous infection extending from an adjacent sinus is a plausible and important explanation.

- Sinus anatomy provides the potential route into the orbit.
- Diabetes increases clinical suspicion but is not required for orbital spread.
- The supplied information does not establish idiopathic orbital inflammation.

Granulomatosis with polyangiitis can first appear in the orbit

- Systemic granulomatosis with polyangiitis (GPA) may initially manifest as orbital disease.
- Orbital involvement can remain apparently localized for prolonged periods.
- Alternatively, GPA may reach the orbit secondarily from affected paranasal sinuses.
- Early orbital localization does not establish that the process is idiopathic.

Two pathways by which GPA involves the orbit

Primary recognized orbital presentation

GPA involving sinonasal tissues



Secondary extension into the orbit

An orbit-dominant presentation should still prompt consideration of systemic vasculitic disease.

The systemic GPA pattern explains the orbital warning

- GPA is a necrotizing granulomatous vasculitis.
- Its characteristic systemic pattern involves the upper or lower respiratory tract and small-to medium-sized vessels.
- Focal necrotizing glomerulonephritis is part of the classic triad.
- The orbital lesion may represent one site of a broader vascular and granulomatous disorder.

Orbital histology: Which features prompt GPA?

Finding	Interpretive weight
Chronic lymphocytes / plasma cells alone	Compatible with several inflammatory processes
Vasculitis	Raises possibility of systemic disease
Necrosis	Additional warning against an unqualified idiopathic diagnosis
Degenerating collagen with vasculitis	Robbins specifically highlights concern for GPA

Necrosis is not a routine defining feature of idiopathic inflammation

- Idiopathic orbital inflammation primarily as chronic inflammation with variable fibrosis.
- Necrosis, degenerating collagen, and vasculitis should prompt consideration of GPA.
- These findings require clinicopathologic integration; They do not independently prove GPA.
- The diagnostic question shifts from pattern recognition to evaluation for an underlying cause.

Applied case: Vasculitis in an orbital biopsy

An orbital biopsy shows chronic inflammation, vascular wall injury, and areas of necrosis.

Diagnostic question

Which systemic inflammatory diagnosis deserves particular attention?

Vasculitis · Necrotic tissue injury · Orbital inflammatory presentation

Case interpretation: Investigate GPA

Granulomatosis with polyangiitis is an important possibility.

- Necrosis and vasculitis are red flags.
- The orbit may be the initial site or be involved through sinonasal disease.
- Morphology creates suspicion; Systemic clinical correlation remains necessary.

Idiopathic orbital inflammation is a diagnosis of exclusion

- It is also called orbital inflammatory pseudotumor or orbital pseudotumor.
- The process is inflammatory without an identifiable local or systemic cause.
- Both focal and diffuse orbital patterns occur.
- IgG4-related disease must be excluded before labeling the process idiopathic.

The word “pseudotumor” describes appearance, not neoplasia

Term	Teaching implication
Orbital inflammatory pseudotumor	A mass-forming inflammatory condition can mimic a tumor
Idiopathic orbital inflammation	No local or systemic etiologic explanation identified
Sclerosing dacryoadenitis	Predominant involvement of the lacrimal gland
Orbital myositis	Predominant involvement of extraocular muscles

Unilateral or bilateral disease does not define the cause

- Idiopathic orbital inflammation may be unilateral or bilateral.
- The laterality alone cannot confirm an idiopathic inflammatory process.
- The compartment distribution and histology contribute additional information.
- Systemic conditions and neoplasia remain in the differential regardless of side.

Idiopathic inflammation can involve several orbital compartments

Diffuse orbital involvement

Multiple tissue elements

Sclerosing dacryoadenitis

Lacrimal gland

Orbital myositis

Extraocular muscles

Posterior scleritis

Tenon capsule / globe-adjacent fascia

Sclerosing dacryoadenitis localizes disease to lacrimal tissue

- Inflammation can be confined predominantly to the lacrimal gland.
- The lacrimal gland sits superotemporally in the orbit.
- Lacrimal enlargement can therefore produce inferomedial globe displacement.
- Lacrimal location does not by itself distinguish idiopathic inflammation from systemic disease or neoplasm.

Orbital myositis localizes inflammation to extraocular muscles

- The extraocular muscles are a recognized site of localized idiopathic orbital inflammation.
- Muscle involvement can contribute to orbital volume expansion and proptosis.
- Thyroid-associated orbitopathy is another important muscle-centered process.
- The underlying mechanism and tissue distribution must be interpreted together.


Tenon capsule involvement links orbit to the globe surface

- Tenon capsule is the fascial layer that wraps around the eye.
- Inflammatory involvement of this region is discussed under posterior scleritis in the source.
- A disease confined to the globe-adjacent fascial layer differs anatomically from diffuse orbital involvement.
- The term “orbital pseudotumor” therefore encompasses multiple patterns of distribution.

Localized versus diffuse idiopathic orbital inflammation

Distribution	Tissue emphasized	Useful question
Diffuse orbital	Multiple orbital tissue elements	Is the process truly orbit-wide?
Sclerosing dacryoadenitis	Lacrimal gland	What else enlarges the lacrimal gland?
Orbital myositis	Extraocular muscle	Could thyroid orbitopathy explain this?
Posterior scleritis / Tenon region	Fascial tissues about the eye	Is inflammation predominantly globe-adjacent?

The central histologic pattern is chronic inflammation with fibrosis




**Mixed
lymphoplasma
cytic
inflammation**

Schematic morphology — native vector shapes; Not an H&E specimen

Lymphocytes and plasma cells are typical components

- The inflammatory infiltrate commonly contains lymphocytes and plasma cells.
- These cells form a mixed chronic inflammatory population rather than a single named neoplastic cell type.
- Histologic distribution and the background stromal response should be evaluated together.
- A mixed infiltrate alone cannot establish the etiology of orbital inflammation.

Eosinophils may accompany the chronic inflammatory infiltrate




**Eosinophils
may
accompany
chronic
inflammation**

Schematic morphology — native vector shapes; Not an H&E specimen

Reactive germinal centers may be present

- Germinal centers can occur within orbital inflammatory infiltrates.
- Their presence suggests reactive lymphoid hyperplasia in the Robbins description.
- This interpretation requires the broader tissue pattern, not the presence of a single follicle.
- Orbital lymphoma remains an important competing diagnostic category.

Fibrosis can replace orbital fat



**Dense collagen
can replace
orbital fat**

Schematic morphology — native vector shapes; Not an H&E specimen


Fibrosis is a spectrum rather than a binary finding

- The source emphasizes variable degrees of fibrosis in idiopathic orbital inflammation.
- Some biopsies are inflammation-rich; Others may be markedly sclerotic.
- The mixture of chronic inflammatory cells and stromal collagen is an important morphologic pairing.
- Severe sclerosis should reinforce the need to consider systemic fibroinflammatory disease.

Reading an orbital inflammatory biopsy systematically

Sequence	Question for the pathologist
1. Tissue context	Is the sample orbital fat, lacrimal tissue, muscle, or fascial tissue?
2. Inflammatory cells	Is the infiltrate mixed, lymphoplasmacytic, or otherwise unusual?
3. Stromal response	How much fibrosis is present, and what tissue is replaced?
4. Warning features	Are vasculitis, necrosis, or degenerating collagen present?

Vasculitis changes the diagnostic interpretation



Robbins specifically calls out GPA when necrosis and degenerating collagen accompany vasculitis.

Orbital fibrosis with other organ lesions requires explanation

- Inflammation centered in the orbit is usually localized in the idiopathic description.
- Concomitant sclerosing lesions may involve retroperitoneum, mediastinum, or thyroid.
- That distribution raises the possibility of a systemic fibroinflammatory disorder.
- Robbins particularly highlights IgG4-related disease.

IgG4-related disease must be excluded before “idiopathic”

- IgG4-related disease can involve periorbital tissues and the lacrimal gland.
- It may cause tumor-like fibroinflammatory lesions resembling orbital pseudotumor.
- Orbital disease can coexist with fibrosis in other organs.
- The presence of orbital inflammation alone does not distinguish these diagnoses.

Systemic IgG4-related disease unifies distant lesions

**Periorbital / lacrimal
fibroinflammation**

**Systemic immune-mediated
fibroinflammatory disease**



**Possible retroperitoneal, thyroid,
or mediastinal sclerosis**

Shared tissue pattern and distribution support investigation of an underlying systemic process.

Robbins characterizes the IgG4-related tissue pattern

- Infiltrates often contain abundant IgG4-producing plasma cells and lymphocytes.
- Fibrosis and obliterative phlebitis form part of the characteristic description.
- Serum IgG4 is usually elevated, but tissue pattern and clinical context remain important.
- A quantitative diagnostic threshold is not supplied in this source excerpt.

IgG4 positivity does not prove an IgG4-driven mechanism

- The source describes increased IgG4-producing plasma cells as a hallmark of disease.
- It does not establish that the IgG4 antibody is itself the cause of fibrosis.
- The pathogenesis remains incompletely understood.
- The distinction between histologic association and causal mechanism is important.

Idiopathic inflammation versus IgG4-related disease

Dimension	Idiopathic orbital inflammation	IgG4-related disease
Etiologic framing	No identifiable local / systemic cause	Systemic fibroinflammatory disease
Histologic overlap	Lymphocytes, plasma cells, fibrosis	Lymphoplasmacytic infiltrate and fibrosis
Particularly relevant clues	Variable patterns; Diagnose by exclusion	IgG4-rich infiltrates, obliterative phlebitis
Beyond orbit	Usually confined to orbit	May affect many other organs

Orbital lymphoma is a clinically important mimic

- Non-Hodgkin lymphoma can involve the entire orbit or individual compartments.
- The lacrimal gland may be affected by lymphoma as well as inflammation.
- A lymphoid infiltrate is not automatically synonymous with a benign inflammatory process.
- The disease must be classified as a hematolymphoid neoplasm if lymphoma is established.

Three competing explanations for lacrimal enlargement

Process	Why it belongs in the differential
Sclerosing dacryoadenitis	Localized idiopathic orbital inflammation
Systemic sarcoidosis	Can enlarge the lacrimal gland and cause non-axial proptosis
Lacrimal gland lymphoma	A neoplastic process can localize to the same compartment
Lacrimal epithelial tumor	Pleomorphic adenoma or adenoid cystic carcinoma are examples

Encapsulation may favor selected orbital masses

- Relatively few orbital masses are encapsulated.
- Examples include pleomorphic adenoma of the lacrimal gland, dermoid cyst, and neurilemmoma.
- Recognition of encapsulation on imaging may help anticipate the underlying pathology.
- Encapsulation is a feature of lesion architecture, not a general rule for inflammation.

Metastatic tumors can also resemble inflammation

- Metastatic prostatic carcinoma can present clinically like idiopathic orbital inflammation.
- Metastatic neuroblastoma and Wilms tumor may be associated with periorbital ecchymoses.
- Sinonasal neoplasms can extend into the orbit.
- A mass-like presentation therefore requires more than an inflammatory appearance to establish etiology.

A practical orbital differential matrix

Orbital process	Useful source-derived clue
Sinus-origin infection	Bacterial/fungal sinus disease contiguous with orbital walls
Granulomatosis with polyangiitis	Sinus association; Vasculitis ± necrosis
IgG4-related disease	Fibroinflammation; Possible systemic sclerosis
Idiopathic orbital inflammation	Mixed chronic inflammation / variable fibrosis after exclusions
Neoplasm	Lymphoma, lacrimal tumors, or metastasis may mimic an inflammatory lesion

A source-grounded diagnostic reasoning sequence

Orbital mass or inflammatory symptoms



Evaluate anatomic distribution and adjacent sinus disease



Assess systemic disease, IgG4-related disease, and neoplasm



Integrate biopsy morphology with clinical information



Reserve “idiopathic” for unexplained inflammation

Applied case: Lymphoid follicles in an orbital lesion

A lacrimal-region biopsy contains chronic inflammatory cells and several germinal centers.

Diagnostic question

What interpretation is supported—and which important alternative remains?

Germinal centers · Lacrimal localization · Lymphoid tissue

Case interpretation: Reactive pattern, but not a final label

Reactive lymphoid hyperplasia is suggested, but lymphoma and systemic processes remain considerations.

- Germinal centers as suggesting a reactive process.
- Lymphoma may involve the same lacrimal compartment.
- Assessment of the complete lesion and context is required.

Applied case: Orbital fibrosis plus retroperitoneal sclerosis

An orbital fibroinflammatory lesion is identified in a patient who also has retroperitoneal fibrosis.

Diagnostic question

Which systemic diagnosis particularly merits investigation?

Orbital chronic inflammation · Marked fibrosis · Fibrosis at a distant organ site

Case interpretation: Consider IgG4-related disease

IgG4-related disease is an important unifying possibility.

- Orbital and retroperitoneal fibroinflammation within its organ spectrum.
- Lymphoplasmacytic infiltrates and fibrosis may overlap with apparent idiopathic inflammation.
- The coexistence of lesions argues against an unqualified idiopathic designation.

Summary: Pattern, significance, differential

Observation	Diagnostic implication
Sinus disease with orbital swelling	Exclude contiguous bacterial/fungal infection
Vasculitis with necrosis or degenerating collagen	Consider granulomatosis with polyangiitis
Lymphoplasmacytic inflammation and variable fibrosis	Typical morphology but not etiologically specific
Fibrosis across several organs	Investigate IgG4-related disease
Lymphoid mass or atypical orbital expansion	Keep lymphoma / other neoplasms in the differential

Selected source references

Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. 2026. Chapter 29: The Eye, pp. 1188–1189.

Robbins, Cotran & Kumar. 11th ed. 2026. Chapter 6: IgG4-Related Disease, pp. 207–208.

Robbins, Cotran & Kumar. 11th ed. 2026. Chapter 11: Granulomatosis With Polyangiitis, pp. 469–470.

AAO EyeWiki: Nonspecific Orbital Inflammation. Supplemental terminology and diagnostic context; consulted October 2026.