

Outdoor Air Pollution

Environmental pathology and clinicopathologic mechanisms

Ambient pollutants • mechanisms of injury • clinicopathologic consequences

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives


- Classify major regulated ambient air pollutants and identify important sources.
- Explain how oxidant gases and particles injure the respiratory tract and affect systemic health.
- Recognize clinical vulnerability patterns and the mechanisms, manifestations, and morphology of carbon monoxide poisoning.

Outdoor air pollution is a mixed exposure

- Ambient air contains gaseous agents and particulate matter rather than a single toxic chemical.
- Pollutant mixtures are especially prominent in cities and near heavy industry.
- Simultaneous exposure complicates assignment of an individual symptom to one pollutant.

Why lungs are the primary target


- The respiratory tract is directly exposed to ambient gaseous and particulate contaminants.
- Epithelial injury and inflammation explain much of the immediate pulmonary disease burden.



Mechanistic schematic

Effects extend beyond the lungs

- Adverse effects in multiple organs, possibly mediated by systemic inflammation.
- Particulate exposure is particularly associated with secondary cardiovascular effects.



Mechanistic schematic

Exposure is geographically uneven

- Major urban centers experience combinations of traffic, industrial, and other combustion emissions.
- Concentrations may be especially high near major industrial or transport sources.
- Residence, occupation, and time outdoors modify the circumstances of exposure.

A useful first classification

Gaseous pollutants

- Ozone (O₃), nitrogen dioxide (NO₂), sulfur dioxide (SO₂)
- Carbon monoxide (CO): Systemic asphyxiant
- Different gases produce different tissue-level effects

Particulate pollutants

- Solid or liquid material suspended in air
- Categorized by aerodynamic size (PM₁₀; PM_{2.5})
- Fine particles can reach distal airspaces

Six pollutants covered by ambient standards


Pollutant	Symbol / category	Core clinical emphasis
Sulfur dioxide	SO ₂	Respiratory irritation and asthma
Carbon monoxide	CO	Carboxyhemoglobin; systemic hypoxia
Ground-level ozone	O ₃	Oxidative epithelial injury
Nitrogen dioxide	NO ₂	Airway reactivity; infection risk
Lead	Pb	Regulated ambient pollutant; detailed toxicity elsewhere
Particulate matter	PM ₁₀ / PM _{2.5}	Lung inflammation; cardiac effects

Particulate size is part of the definition

- PM₁₀ designates particles smaller than 10 µm in diameter.
- PM_{2.5} designates fine particles smaller than 2.5 µm in diameter.
- These are particle-size categories, not specific chemical compositions.

Smog reflects mixed atmospheric contaminants

- The term combines smoke and fog and describes a visibly polluted urban atmosphere.
- Ozone may coexist with sulfur dioxide and particles, generating overlapping exposures.



Mechanistic schematic

Air pollution long predates modern industry

- Early concerns about coal smoke and respiratory disease in London.
- John Evelyn described health consequences of thick, contaminated air in 1661.
- The distinguishing modern features are pollutant sources, composition, and emission controls.

Ambient standards are population-level controls

- In the United States, the EPA regulates six criteria pollutants through National Ambient Air Quality Standards.
- Standards are intended to limit harmful environmental exposures.
- They do not guarantee equal biological susceptibility among exposed people.

Host factors change vulnerability

More vulnerable


- Preexisting asthma or chronic obstructive lung disease
- Established cardiac or chronic pulmonary disease
- Children and persons exercising outdoors (pollutant dependent)

Physiologic considerations

- Baseline reserve affects clinical response
- Airway reactivity can amplify irritant effects
- Exposure duration and intensity matter

Inhaled xenobiotics can act locally or systemically

- The toxicology parent section distinguishes injury at the entry site from injury after systemic transport.
- Outdoor pollutants exemplify both patterns: Airway irritants versus CO-mediated tissue hypoxia.



Mechanistic schematic

Ozone has opposite roles by altitude

Stratospheric ozone

- Situated high above the earth
- Absorbs harmful solar ultraviolet radiation
- Depletion increases environmental vulnerability

Ground-level ozone

- Formed in sunlight from emitted precursors
- Reactive ambient respiratory pollutant
- Can provoke oxidant injury and inflammation

Altitude determines ozone's role


- Protective ozone in the stratosphere and harmful ozone near the ground are chemically the same molecule.
- Health impact depends on atmospheric location and exposure route.

STRATOSPHERE • O₃ absorbs ultraviolet radiation

LOWER ATMOSPHERE • O₃ injures airway epithelium

Stratospheric ozone intercepts ultraviolet radiation


- The stratospheric layer provides a protective atmospheric function.
- Robbins locates it roughly 10–30 miles above the earth's surface.



Mechanistic schematic

Chlorofluorocarbons cause ozone depletion

- Chlorofluorocarbons historically used in cooling systems and aerosols can reach the stratosphere.
- Chemical reactions involving these compounds destroy ozone molecules.




Mechanistic schematic

The ozone hole became a global policy problem

- Marked Antarctic ozone depletion was recognized in 1985.
- The Montreal Protocol of 1987 established international phase-out commitments for ozone-depleting compounds.
- Declining Antarctic ozone-hole size following reduced CFC use.

Ground-level ozone is a secondary pollutant

- Nitrogen oxides and volatile organic compounds react in the presence of sunlight.
- The reactants are released by motor vehicles and industrial sources.



Mechanistic schematic

Precursors are not identical to the product

Emitted precursors


- Nitrogen oxides (NO_x)
- Volatile organic compounds (VOCs)
- Vehicle exhaust and industrial emissions

Secondary product

- O₃ is generated by atmospheric reactions
- Sunlight enables the relevant chemistry
- Ground-level concentration need not track a single source directly

Ozone toxicity begins with oxidant chemistry


- A major component of ozone toxicity to free radical generation.
- Oxidant stress injures the respiratory epithelium and initiates inflammatory responses.



Mechanistic schematic

Ozone injures multiple respiratory cell sites

- Both conducting airway epithelia and type I alveolar cells are described as vulnerable.
- Mediator release links direct cell injury to local pulmonary inflammation.



Mechanistic schematic

Healthy individuals may develop acute effects

- Upper respiratory tract inflammation follows exposure.
- Symptoms include chest discomfort and reduced pulmonary function.
- The response may be modest in a healthy individual yet clinically important in reduced-reserve patients.

Preexisting obstructive disease increases risk

Healthy respiratory system

- Acute inflammatory symptoms
- Mild reduction in measured lung function
- Reduced exercise tolerance is possible

Asthma or chronic obstruction


- More consequential airway narrowing / reactivity
- Asthma-related hospitalization burden rises
- Greater clinical impact from a similar ambient exposure

Exercise changes the clinical impact of exposure

- Athletes and outdoor workers are exposed during sustained outdoor activity.
- Ozone exposure in these populations with reduced exercise capacity.
- Exposure assessment should include activity and environment, not only diagnosis.

Ozone contributes to a combined irritant burden

- Ozone often accompanies sulfur dioxide and particulate matter in polluted air.
- Multiple inhaled agents can produce overlapping irritation, inflammation, and reduced lung function.



Mechanistic schematic

Nitrogen dioxide affects airway reactivity

- NO₂ is one of the six regulated ambient air pollutants.
- Robbins' health-effects table associates exposure with increased airway reactivity in healthy adults.
- Its clinical effects are particularly relevant to susceptible respiratory populations.

NO₂: Different groups, different outcomes

Adults with asthma

- Decreased lung function in the source table
- Underlying hyperreactive airways increase clinical significance

Children


- Increased respiratory infections in the source table
- Observed outcome differs from acute physiologic reactivity

Sulfur dioxide is linked to fuel combustion

- Coal- and oil-burning power plants emit sulfur dioxide.
- Copper smelting and paper mills are additional sources.
- Source recognition is useful for interpreting industrial and urban exposure contexts.

Atmospheric transformation produces irritant species

- Released sulfur dioxide can be converted into sulfur trioxide and sulfuric acid.
- These irritant products contribute to burning sensations and respiratory symptoms.



Mechanistic schematic

Upper-airway irritation is a prominent effect

- Burning sensations may occur in the nose and throat.
- Exposed individuals can experience difficulty breathing.
- These symptoms reflect direct irritation of respiratory mucosa.

Underlying lung disease alters severity

Without established disease

- Increased respiratory symptoms
- Acute irritation and breathing difficulty

With lung disease / asthma


- Asthma attacks and reduced pulmonary function
- Increased hospitalizations reported in the source table
- Higher vulnerability among chronic lung disease groups

Acid aerosols disrupt airway defense

- The source table associates acid aerosol exposure with altered mucociliary clearance.
- Mucociliary transport normally helps remove inhaled material from conducting airways.
- Children show increased respiratory infections in the source table.

Impaired clearance creates a clinical link

- Loss of efficient airway clearance provides a mechanistic interpretation for observed infection associations.
- The table also identifies lower lung function and hospitalization in people with asthma.



Mechanistic schematic

Particulate matter is heterogeneous


- Particulates are commonly called soot discussion.
- Different materials share a size-based classification despite variable chemical composition.
- The source highlights the pulmonary and cardiovascular morbidity associated with particles.

Combustion is a major source of particulates

- Coal- and oil-fired power plants emit particulates.
- Industrial combustion processes add to ambient particulate levels.
- Diesel exhaust is another important source.


Particle diameter influences where material deposits

- Larger particles are more often removed in the nose or by mucociliary transport.
- Fine particles penetrate deeper and may reach the alveolar compartment.




Coarse fraction

Upper-airway capture



PM₁₀

Thoracic penetration



PM_{2.5}

Fine; Reaches alveoli

Upper-airway capture versus alveolar access

Particles $>10\text{ }\mu\text{m}$


- More likely to be filtered by the nose
- May become trapped in mucus and cleared by airway cilia
- Reduced access to alveolar spaces

Fine particles

- Can enter small airways and alveoli
- Encounter alveolar inflammatory cells
- Provoke distal lung inflammation

Alveolar phagocytes recognize inhaled particles


- Phagocytosis of inhaled particles by macrophages and neutrophils.
- These activated cells release inflammatory mediators following uptake.



Mechanistic schematic

Inflammation is the central injury pathway

- Particle-induced inflammatory signaling can worsen preexisting respiratory disease.
- This is central to the morbidity associated with airborne particles.



Mechanistic schematic

Children are a vulnerable respiratory group

- Robbins' health-effects table associates particle exposure with increased respiratory infections in children.
- Respiratory outcomes vary across age groups and underlying disease status.
- The association concerns exposure-related population risk, not a disease-specific histologic pattern.

Pulmonary and cardiac consequences coexist

Pulmonary

- Airway / alveolar inflammation
- Decreased lung function in chronic lung disease
- Exacerbation of existing pulmonary illness

Cardiovascular

- Secondary adverse cardiac effects
- Greater concern when baseline heart disease exists
- Cardiac exacerbations contribute to mortality burden

Particle exposure is linked to daily mortality


- An association between increased ambient PM₁₀ and overall daily mortality.
- Exacerbation of pulmonary and cardiac disease accounts for much of this risk.
- The exact numeric estimate in the project Markdown contains an apparent unit-transcription problem and is intentionally not reproduced here.

Fine particles may promote lung carcinogenesis

- Experimental work suggesting inhaled particulates can promote lung cancer.
- The statement concerns experimental evidence and should not be represented as a quantified clinical effect.
- The source excerpt does not specify a single molecular pathway, histologic subtype, or attributable risk.

Pulmonary inflammation may affect distant organs

- The textbook suggests systemic inflammation as one explanation for effects outside the lung.
- This model links the primary inhalational exposure with cardiac complications without assuming direct particle translocation.



Mechanistic schematic

Interpret particle risk in context

- Size predicts deposition pattern, but composition and individual susceptibility remain relevant.
- Particulate concentration is a population-level exposure measure rather than a clinical diagnosis.
- The same ambient episode may present as respiratory or cardiac deterioration across patients.

Carbon monoxide is produced by incomplete oxidation

- CO forms when carbonaceous fuels are incompletely burned.
- Common sources include engines, furnaces, and cigarettes.
- Unlike the irritant pollutants, CO primarily threatens oxygen transport and tissue oxygenation.

CO is difficult to detect without measurement

Physical properties

- Colorless
- Odorless and tasteless
- Nonirritating gas

Clinical implication

- Exposure may not be noticed
- Symptoms can progress insidiously
- The environment and exposure history are essential clues

Ambient CO elevations tend to be localized

- CO is relatively short-lived in outdoor air because it is oxidized to carbon dioxide.
- Elevated outdoor concentrations may therefore be transient and close to sources.
- Confinement near combustion equipment can produce disproportionately hazardous exposures.

Occupational contexts can produce chronic exposure


- Tunnels, underground garages, and highway toll booths as settings with automobile fumes.
- Persistent low-level exposure is relevant even without an acute high-dose event.
- Source proximity and repeated exposure help distinguish chronic from single-event poisoning.

Closed spaces magnify acute CO danger

- An enclosed garage containing a running engine can rapidly accumulate dangerous CO.
- Improperly used gasoline-powered generators may produce poisoning during power outages.
- Mine fires are another setting described for potentially severe exposure.

CO displaces oxygen from hemoglobin


- Hemoglobin has markedly higher affinity for CO than for oxygen—about 200-fold.
- Carboxyhemoglobin cannot transport oxygen in the usual manner.



Mechanistic schematic

Loss of functional hemoglobin causes tissue hypoxia

- Tissue injury reflects impaired oxygen transport rather than primary airway inflammation.
- High oxygen demand makes the central nervous system especially vulnerable.




Mechanistic schematic

Severity correlates with carboxyhemoglobin burden

- Systemic hypoxia when CO occupies roughly 20–30% of hemoglobin.
- Unconsciousness and death become likely with approximately 60–70% saturation.
- These textbook ranges illustrate severity; Clinical presentation must still be assessed in context.

Neurologic depression can be deceptively insidious

- Progressive central nervous system depression may occur without a marked irritant warning.
- Affected individuals may become unable to recognize or escape the exposure.



Mechanistic schematic

Chronic CO exposure can accumulate biologic burden

- The stability of carboxyhemoglobin after it forms.
- Repeated or persistent exposure can produce substantial carboxyhemoglobin levels.
- Slowly developing hypoxia may cause widespread ischemic injury in the brain.

Basal ganglia are notably vulnerable

- Chronic CO poisoning can produce CNS ischemic injury.
- The textbook particularly identifies the basal ganglia and lenticular nuclei.
- Regional vulnerability should be understood as hypoxic injury, not a unique toxic deposit.

Neurologic sequelae may persist after exposure ends

- Many patients improve following removal from exposure.
- Permanent deficits may affect memory, vision, hearing, and speech.
- Persistent neurologic impairment reflects the consequences of prior tissue injury.

Acute poisoning may be fatal before lesions evolve

- Short, intense CO exposures can induce unconsciousness or death.
- When death occurs very rapidly, morphologic changes may be absent.
- The absence of gross lesions therefore does not exclude severe CO poisoning.

Morphology in survivors reflects hypoxia

Possible brain findings

- Mild edema
- Punctate hemorrhages
- Hypoxia-associated neuronal changes

Interpretive limitation


- Not specific to CO
- Depend on survival time and degree of hypoxia
- Require exposure history and blood testing

Cherry-red discoloration is a limited clue

- Cherry-red skin and mucous membranes in light-skinned individuals with acute poisoning.
- Coloration may obscure the severity of underlying oxygen deprivation.
- It should not be used alone to exclude or establish a diagnosis.

Confirm exposure by blood measurement

- Robbins specifies measurement of blood carboxyhemoglobin for diagnosis.
- Clinical interpretation also depends on the exposure history and neurologic findings.



Mechanistic schematic

Irritant injury differs from systemic asphyxiation

O₃ / SO₂ / particles

- Airway epithelium or alveolar interaction
- Oxidative and/or inflammatory responses
- Respiratory symptoms; Downstream cardiac burden for PM

CO

- High-affinity hemoglobin binding
- Reduced blood oxygen transport
- CNS depression and hypoxic organ injury

Characteristic associations across pollutants

Agent	Mechanism emphasized	Characteristic clinical effects
Ozone	ROS and epithelial injury	Airway inflammation; ↓ lung function
NO ₂	Airway reactivity	Asthmatic ↓ function; infections in children
SO ₂ / acid aerosols	Respiratory mucosal irritation	Asthma attacks; altered clearance
Particulates	Phagocyte activation, inflammation	Pulmonary + secondary cardiac disease
CO	Carboxyhemoglobin / systemic hypoxia	Neurologic depression; ischemic injury

Susceptibility patterns Table 9.1

Population	Pollutant concern	Associated effect
Healthy adults	O ₃ / NO ₂	Airway reactivity or lower lung function
Asthma	O ₃ / SO ₂ / acid aerosols	Reduced function; hospitalization
Children	NO ₂ / acid aerosols / PM	Respiratory infections
Chronic pulmonary or cardiac disease	PM; SO ₂ in chronic lung disease	Exacerbation and mortality burden
Athletes and outdoor workers	O ₃	Reduced exercise capacity

Case: Episodic wheeze during urban smog

CLINICAL PRESENTATION

A patient with established asthma develops chest tightness and worsening breathlessness after prolonged outdoor activity on a heavily polluted afternoon.

QUESTION

Which pollutant-related mechanisms most directly explain the deterioration?

Interpretation: Oxidant and irritant exposure

- Ground-level ozone can produce reactive oxygen species, epithelial injury, and inflammatory mediator release.
- Sulfur dioxide and acidic derivatives may add airway irritation and trigger asthma attacks.
- The mixed nature of smog prevents attributing the episode to one pollutant without exposure data.

Case: Coma beside combustion equipment

CLINICAL PRESENTATION

A person is found unconscious in an enclosed space containing a running fuel-powered device. There is no obvious smell or irritant warning.

QUESTION

What exposure should be suspected, what is the mechanism, and which blood test supports the diagnosis?

Interpretation: Carbon monoxide poisoning

- Suspect CO produced by incomplete oxidation of fuel in the enclosed environment.
- CO forms carboxyhemoglobin and impairs systemic oxygen transport, producing CNS hypoxia.
- Blood carboxyhemoglobin measurement supports the diagnosis; Characteristic skin color is not required.

Case: Cardiac deterioration during particulate smog

CLINICAL PRESENTATION

A patient with chronic heart disease develops worsening cardiopulmonary symptoms during a sustained episode of particulate-rich urban pollution.

QUESTION


Which particulate mechanism could link a respiratory exposure to an adverse cardiac effect?

Interpretation: A lung-to-systemic inflammatory link

- Fine particles may reach alveoli and be taken up by macrophages and neutrophils.
- Release of inflammatory mediators explains pulmonary inflammation and suggests a pathway to systemic effects.
- The textbook links particulate exposure with exacerbations of preexisting pulmonary and cardiac disease.

The major mechanism map

- For ozone and particulates, epithelial oxidant injury or inflammatory-cell activation dominates.
- For carbon monoxide, impaired hemoglobin-mediated oxygen delivery dominates.



Mechanistic schematic

Population protection and individual care address different levels

- Ambient emission control targets exposure in whole populations.
- Clinical risk assessment identifies asthma, chronic lung disease, cardiac disease, activity patterns, and combustion environments.
- A suspected CO exposure requires prompt recognition because it may occur without odor or airway irritation.

Core clinicopathologic conclusions

- Ambient air pollution is a mixture: Gas chemistry and particle size determine exposure biology.
- Oxidant gases injure epithelium; Fine particles recruit inflammatory cells and have systemic associations.
- CO is mechanistically different: Carboxyhemoglobin produces hypoxia and often neurologic injury.
- Preexisting pulmonary or cardiac disease changes the consequences of exposure.

Primary textbook and source limitations

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th edition (2026), Chapter 9, printed pp. 371–373.
- Table 9.1 (Health Effects of Outdoor Air Pollutants) cites Bascom R et al. Am J Respir Crit Care Med. 1996;153:477–498.
- Content derived from supplied project Markdown; original chapter PDF and source figures were not supplied for visual cross-checking.