

Pelvic Inflammatory Disease

Ascending infection · tubal inflammation · reproductive sequelae

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning objectives

- Trace infection from the lower genital tract into the adnexa.
- Contrast gonococcal mucosal disease with puerperal deep-tissue spread.
- Recognize acute suppurative and chronic fibrosing salpingitis.
- Explain infertility, ectopic pregnancy, and systemic complications.

Organizing question

How does the site and pattern of infection determine the morphology and long-term consequences?

PID is typically an ascending genital tract infection



Anatomical significance

The infection usually begins in the vulva or vagina and ascends.

The endocervix, tubes, and tubo-ovarian region can be involved.

The distribution of inflammation depends on the organism and route.

PID describes a multi-organ inflammatory process

- A single ascending infection can affect multiple genital tract sites.
- Salpingitis is a principal lesion responsible for late reproductive harm.
- Spread into the ovary or peritoneum increases disease extent.

Pathology perspective

PID is a disease process; Salpingitis, pyosalpinx, and tubo-ovarian abscess name particular morphologic manifestations.

Typical clinical presentation reflects pelvic inflammation

- Pelvic pain accompanies inflammation of reproductive and pelvic tissues.
- Adnexal tenderness localizes involvement near tubes and ovaries.
- Fever and vaginal discharge may accompany infection.
- Clinical features vary; The source describes this typical syndrome.

Clinical correlation

Symptoms arise from the anatomic distribution of infection and inflammatory exudate—not from a unique single histologic lesion.

Two major etiologic settings

Sexually associated PID	Postabortion / puerperal PID
Neisseria gonorrhoeae	Often polymicrobial infection
Chlamydia infection	Staphylococci and streptococci
Typically ascends from lower tract	Coliforms and Clostridium perfringens may contribute
Mucosal inflammation emphasized with gonococcus	Deeper tissue inflammation emphasized after instrumentation or delivery

Gonococcal infection remains an important cause

- *Neisseria gonorrhoeae* is a recognized cause of ascending PID.
- PID is a serious complication of gonorrhea in females.
- The endocervical mucosa is the most frequent initial focus.
- Ascending spread commonly emphasizes tubal mucosal injury.

Mechanism to remember

The tubal mucosa is an early target; Neutrophilic inflammation and exudate later compromise luminal architecture.

Chlamydia can progress beyond cervicitis

- Chlamydia trachomatis often causes cervicitis.
- In some patients, infection ascends into the uterus and fallopian tubes.
- Endometritis and salpingitis explain its contribution to PID.

Clinical distinction

Only some infections ascend beyond cervicitis; Progression is not inevitable.

Pregnancy-related events define another PID context

- Infection can follow spontaneous or induced abortion.
- Puerperal infection may develop after normal or abnormal delivery.
- Uterine procedures provide another setting for nongonococcal spread.
- Microbial diversity shapes the clinical and morphologic pattern.

Terminology

Puerperal refers to infections associated with the postpartum period; Postabortion infections can also be polymicrobial.

Polymicrobial invaders named in the source

Organism group	Examples explicitly cited
Gram-positive cocci	Staphylococci
Gram-positive cocci	Streptococci
Enteric-type bacteria	Coliforms
Anaerobic spore-forming bacteria	Clostridium perfringens


Etiologic context predicts a different tissue distribution

- Gonococcus produces conspicuous mucosal and luminal inflammation.
- Nongonococcal puerperal infections often affect the deeper wall.
- Wall and serosal involvement facilitates extension outside the organ.

Why it matters

Different routes lead to different patterns of salpingitis, pelvic spread, and bacteremia—even when symptoms overlap.

Gonococcal spread follows an ascending mucosal pathway




Pathogenic logic

The organism originates in lower tract mucosa and reaches the tube along connected mucosal surfaces.

Causal / anatomical sequence — schematic

Early inflammatory changes follow gonococcal inoculation




Interpretation

The interval describes approximate onset of inflammation, not a universal clinical incubation period.

Causal / anatomical sequence — schematic

The endocervix is the usual initial gonococcal focus



Anatomical significance

Mucosal infection usually starts within the endocervix.

The site is anatomically continuous with the upper tract.

Local inflammation can precede tubal extension.


Alternative initial gonococcal sites are glandular

- Infection may begin in the Bartholin gland.
- Other vestibular glands may be involved.
- Periurethral glands provide another possible initial site.
- The same organism can then ascend from the lower tract.

Anatomic nuance

The endocervix is the common starting site, but glandular lower-tract infection is explicitly described as an alternative.

Ascending spread makes salpingitis a central lesion




Anatomical significance

Infection progresses from lower genital structures toward tubes.

Tubal mucosal inflammation causes epithelial and plical injury.

Further extension may involve the ovary and adjacent pelvis.

Nongonococcal infection may spread via deeper channels



Different from gonococcus


Lymphatic or venous channels are proposed routes after uterine procedures, contrasting with mucosal ascent.

Causal / anatomical sequence — schematic

Mucosal and deep-wall PID are distinct patterns

Feature	Gonococcal pattern	Puerperal / other nongonococcal
Major route	Mucosal ascent	Lymphatic / venous spread proposed
Predominant inflammation	Mucosa and tubal lumen	Deep organ-wall layers
Frequent extension	Tubal lumen → adnexa	Serosa, ligaments, pelvic tissues
Bacteremia	Less frequent in comparison	More frequent with staphylococcal / streptococcal disease

The primary anatomic route shapes downstream injury



Integrated inference

Compare mucosal exudation and tubal scarring with deep-wall inflammatory extension and systemic dissemination.

Causal / anatomical sequence — schematic


Acute gonococcal disease targets the mucosal surface

- Gonococcal inflammation may be intense on the involved mucosa.
- Neutrophils predominate in the acute exudative response.
- Tubal mucosal injury provides the substrate for later fibrosis.
- The tubal lumen can accumulate abundant pus.

Morphologic anchor

The defining acute lesion is a suppurative salpingitis involving tubal mucosa and lumen.

Acute salpingitis: Edema and purulent luminal material



Purulent material in the lumen

Morphologic interpretation

The tube is congested and its mucosal folds become inflamed.

The lumen can contain abundant purulent exudate.

Inflamed plicae and luminal exudate are illustrated schematically, not as a tissue section.

Original anatomical/morphologic schematic — not a histologic specimen


Inflammatory infiltrates alter tubal mucosal architecture

- Neutrophils infiltrate actively inflamed tubal mucosa.
- Plasma cells and lymphocytes may also be present.
- Inflammation expands and injures the mucosal plicae.
- These changes impair the mucosal surface before permanent scarring develops.

Histopathologic logic

The infiltrate is mixed but acute suppuration is central; Chronic inflammatory cells can coexist with neutrophils.

Acute inflammation leads to epithelial injury




Consequences

Denudation exposes opposing mucosal folds to adhesion and reparative fusion later in the process.

Causal / anatomical sequence — schematic

Purulent exudate may fill the tubal lumen



Neutrophilic exudate fills much of the lumen

Morphologic interpretation

The involved lumen becomes distended with inflammatory exudate.

Neutrophilic pus characterizes the acute suppurative phase.

Persistent exudate can form a pyosalpinx.

Original anatomical/morphologic schematic — not a histologic specimen

An open fimbrial end permits leakage of pus



Fluid / exudate may pass beyond open fimbriae

Morphologic interpretation

Purulent material may escape at the fimbriated end.

Spread beyond the tube helps explain pelvic/peritoneal involvement.

This is an anatomic sequence, not a reproduction of a tissue section.

Original anatomical/morphologic schematic — not a histologic specimen

Gonococcal infection may relatively spare the endometrium

- Relative endometrial sparing is described in gonococcal tubal infection.
- The cause of relative endometrial sparing is explicitly stated to be unclear.
- Do not generalize this statement to all PID etiologies.
- Chlamydia may cause endometritis in ascending infection.

Interpretation caveat

Relative sparing does not exclude gonococcal disease when endometritis is identified.

Intraneutrophilic diplococci support gonococcal infection

- Smears of exudate may reveal phagocytosed Gram-negative diplococci.
- Morphologic identification alone is not definitive.
- Organism identification requires culture or gonococcal RNA/DNA detection.


Diagnostic distinction

Cellular morphology suggests etiology;
Microbiologic testing establishes identification.
No synthetic micrograph is presented.

Morphology and microbiology answer different questions

Observation / test	What it establishes	Limitation
Suppurative salpingitis	Tissue inflammatory pattern	Does not uniquely identify organism
Smear: Intracellular diplococci	Morphologic clue to gonococcus	Not sufficient alone for definitive diagnosis
Culture	Gonococcal detection	Diagnostic sample and assay matter
Gonococcal RNA or DNA detection	Evidence of organism	Does not alone map severity or extent

Ascending infection may reach the tubo-ovarian region



Anatomical significance

Spread may extend from fallopian tubes to the ovary.

Combined tubal and ovarian inflammation is salpingo-oophoritis.

Localized pus may eventually form a tubo-ovarian abscess.


Acute suppurative salpingitis is the key tubal lesion

- Tubal mucosa is congested and inflamed.
- Plicae show inflammatory infiltration and epithelial injury.
- Neutrophilic exudate expands the tubal lumen.
- Spread can create adnexal suppurative collections.

Teaching definition

Salpingitis is inflammation of the fallopian tube; The acute suppurative form features purulent exudation and mucosal destruction.

Pyosalpinx is pus collected within the tubal lumen



Pus retained within an inflamed fallopian tube

Morphologic interpretation

Location: Within the fallopian tube lumen.

Material: Purulent inflammatory exudate.

Contrast with hydrosalpinx, which describes fluid distention after obstructive scarring.

Original anatomical/morphologic schematic — not a histologic specimen

Salpingo-oophoritis indicates extension to the ovary

- Initial tubal infection spreads to adjacent ovarian tissue.
- The salpingo-oophoritis label records involvement of both sites.
- Persistent focal suppuration may create tubo-ovarian abscess.
- Anatomical extent matters because a localized collection is harder to eradicate.


Site distinction

Salpingitis = tube;

Oophoritis = ovary;

Salpingo-oophoritis = both.

Tubo-ovarian abscess is an adnexal purulent collection



Morphologic interpretation

Localized collections of pus can involve tube and ovary.

A walled-off collection may be difficult for antibiotics to eradicate.

The original schematic does not depict a clinical specimen.

Original anatomical/morphologic schematic — not a histologic specimen

Three related acute morphologic labels

Term	Anatomic emphasis	Principal pathologic feature
Acute salpingitis	Fallopian-tube mucosa and wall	Acute inflammatory infiltration
Pyosalpinx	Fallopian-tube lumen	Collection of pus
Tubo-ovarian abscess	Tubal–ovarian region	Localized suppurative collection


Abscess encapsulation changes antimicrobial accessibility

- A tubo-ovarian abscess is a localized collection of pus.
- A walled-off pus collection can be difficult to eradicate with antibiotics.
- Selected refractory cases may require surgical removal of affected organs.
- Morphologic organization therefore has direct management implications.

Caution


Morphology identifies an abscess but does not, by itself, specify antibiotics or a surgical decision threshold.

Chronic tubal damage can persist after microbes disappear




Causal / anatomical sequence — schematic

Mucosal denudation predisposes plicae to adhesion



Causal / anatomical sequence — schematic

Chronic salpingitis: Plical fusion and architectural distortion



Fused mucosal folds → blind pockets

Morphologic interpretation

Tubal mucosal folds adhere and gradually fuse.

Repair produces fibrous distortions of the lumen.

Blind pouches and glandlike spaces may develop.

Original anatomical/morphologic schematic — not a histologic specimen

Glandlike spaces are a reparative pattern

- Fusion of tubal plicae can enclose residual luminal pockets.
- These glandlike spaces reflect architectural rearrangement in chronic salpingitis.
- They are not evidence of a neoplastic gland-forming process.
- Fibrotic reorganization can compromise tubal transport.

Interpretation

A glandlike appearance can arise from nonneoplastic tubal folding and scarring; Assess the underlying inflammatory architecture.

Chronic salpingitis is a scar-forming lesion

- Organisms may no longer be identifiable in late disease.
- The plicae become fused by reparative scarring.
- Blind luminal pouches alter internal tubal geometry.
- The scarring can extend to the fimbriae and narrow the lumen.


Why chronic disease persists

The permanent clinical effect follows loss of tubal architecture, not necessarily persistence of active infection.

Acute and chronic salpingitis are morphologically different

Feature	Acute salpingitis	Chronic salpingitis
Dominant process	Active purulent inflammation	Repair with scarring
Tubal folds	Congested, infiltrated, denuded	Adherent and fused
Lumen	Purulent exudate may accumulate	Distorted; Blind pouches possible
Clinical implication	Adnexal extension / sepsis	Infertility / ectopic pregnancy

Fimbrial scarring impairs normal oocyte uptake




Reproductive consequence

The fimbriae are needed to collect the released oocyte; Scarring interferes with that function.

Causal / anatomical sequence — schematic

Tubal infertility follows disrupted passage




Pathology → function

Tubal scarring and blockage interfere with reproductive transport and may produce infertility.

Causal / anatomical sequence — schematic

An ectopic pregnancy can follow altered tubal transit




Causal / anatomical sequence — schematic

Clinical hazard

An implantation outside the uterine cavity—often in the tube—is a recognized possible consequence of scarring.

Hydrosalpinx develops when secretions cannot drain



Morphologic outcome


Hydrosalpinx indicates obstruction with retained tubal secretion, not an inherently pus-filled lumen.

Causal / anatomical sequence — schematic

Pyosalpinx and hydrosalpinx must not be conflated

Feature	Pyosalpinx	Hydrosalpinx
Luminal contents	Purulent exudate	Retained tubal secretion / fluid
Dominant setting	Suppurative infection	Fimbrial occlusion after scarring
Morphologic process	Intraluminal pus collection	Tubal dilation from impaired drainage
Key distinction	Pus	Fluid distention

Hydrosalpinx is a distended, obstructed tube



Distended, fluid-filled, obstructed tube

Morphologic interpretation

Fusion of fimbriae prevents normal outflow.

Retained secretion enlarges the tubal lumen.

Persistent distention signals an anatomic late sequela.

Original anatomical/morphologic schematic — not a histologic specimen


Chronic PID may produce persistent pelvic pain

- Chronic pelvic pain can be a long-term sequela of PID.
- Adhesions and residual inflammatory damage can alter pelvic tissue mobility.
- Pain does not establish the specific morphology or current pathogen.

Clinicopathologic limitation

The association is source-established; An individual patient's pain mechanism requires clinical assessment.

Pelvic adhesions can lead to intestinal obstruction



Extrareproductive complication

Bowel-to-pelvis adhesions can mechanically impair intestinal movement and result in obstruction.

Causal / anatomical sequence — schematic


Puerperal infection often extends into deeper tissue

- Compared with gonococcal disease, deep-wall inflammatory involvement is emphasized.
- Tubal mucosa and lumen may be relatively less affected.
- The inflammatory process can extend through organ walls.
- This pattern favors serosal and pelvic extension.

Interpretation

This is a characteristic tendency, not a binary organism-specific microscopic diagnostic criterion.

Deep-wall inflammation can breach organ boundaries



Why severe spread occurs

Full-thickness extension can involve adjacent anatomical compartments beyond the tubal lumen.

Causal / anatomical sequence — schematic

Peritonitis is an important acute complication

- Purulent material can escape from the tubal fimbrial region.
- Other invasive infections may extend through serosa to peritoneum.
- Both mechanisms provide anatomic routes to pelvic peritonitis.
- Peritonitis is an important acute complication of ascending PID.


Clinical consequence

Peritoneal involvement marks spread beyond a localized genital-tract infection.

Bacteremia differs across etiologic patterns

Comparison	Gonococcal PID	Staphylococcal / streptococcal PID
Bacteremia	Described as less frequent	Described as more frequent
Tissue involvement	Strong mucosal component	Often more extensive deeper layers
Major concern	Suppurative tubal injury	Systemic dissemination in addition to local injury

Bacteremia enables distant suppurative complications



Systemic extent

Distant suppurative complications are possible but not universal or typical presentations.

Causal / anatomical sequence — schematic

Early and late consequences require separate thinking

Acute complications	Chronic sequelae
Peritonitis	Infertility and tubal obstruction
Bacteremia	Ectopic pregnancy
Distant infection (e.g., endocarditis)	Persistent pelvic pain
Meningitis or suppurative arthritis	Adhesive intestinal obstruction

Early gonococcal infection is generally antibiotic responsive

- Early gonococcal infection is generally responsive to antimicrobial treatment.
- Response is less straightforward when infection becomes a localized abscess.
- Antimicrobial treatment does not necessarily reverse prior fibrotic injury.

Separate endpoints

Microbiologic control, acute complication control, and preservation of tubal function are different outcomes.

Gonococcal antimicrobial resistance complicates care

- Antimicrobial resistance complicates management of gonococcal infection.
- This challenges assumptions that historical responses remain reliable.
- The supplied text does not specify current susceptibility patterns or treatment regimens.
- Management choices should use separately verified contemporary guidance.

Management boundary

Treatment choices require current antimicrobial susceptibility information and guidelines; This pathology lecture gives no drug regimen.

Tubo-ovarian abscess may require surgical intervention

- A walled-off pus collection can be difficult to eradicate medically.
- The affected organs may occasionally need removal when infection cannot be controlled.
- This management issue follows the morphologic state of abscess formation.

Important boundary

The presence of an abscess alone does not determine the timing or type of surgical intervention.

Polymicrobial postabortion and puerperal PID is harder to control

- The pathogen spectrum is broad and may involve several bacterial groups.
- These infections may be harder to control than early gonococcal PID.
- Both antibiotics and management of the source of infection may matter clinically.

Pathologic explanation

Bacterial diversity and deeper tissue spread help explain the management challenge; No regimen or quantitative claim is assumed.

Case 1: Acute adnexal suppuration

CLINICAL VIGNETTE

A patient presents with pelvic pain, fever, vaginal discharge, and adnexal tenderness. A fallopian tube is dilated by purulent material, with epithelial sloughing and marked inflammation.

Which lesion best accounts for the tubal lumen finding?

Clinicopathologic explanation

Acute suppurative salpingitis with pyosalpinx. Purulence inside the tubal lumen is the defining clue; Adjacent ovarian involvement would support salpingo-oophoritis or tubo-ovarian abscess.

Case 2: Infertility after an earlier infection

CLINICAL VIGNETTE

Months after a prior pelvic infection, a patient is assessed for infertility. A tubal specimen shows fused mucosal folds and enclosed glandlike pockets, without the earlier abundant pus.

What morphologic mechanism explains the infertility?


Clinicopathologic explanation

Chronic salpingitis.
Reparative fusion and scarring distort the tube and fimbriae, impeding oocyte capture and transit. Similar damage also increases ectopic pregnancy risk.

A compartment-based interpretation of PID

Location of injury	Characteristic lesion	Potential consequence
Tubal mucosa/lumen	Acute suppurative salpingitis; Pyosalpinx	Spread to adnexa / peritoneum
Tubal plicae and fimbriae	Fusion, scarring, hydrosalpinx	Infertility / ectopic pregnancy
Tubo-ovarian region	Localized abscess	Difficult eradication
Deep wall / serosa	Invasive puerperal-type inflammation	Peritonitis / bacteremia

Integrated disease progression in four connected stages



Central synthesis

Anatomic route and tissue response explain both early complications and late reproductive harm.

Causal / anatomical sequence — schematic

High-yield clinicopathologic conclusions

- PID is an ascending infection with variable genital and pelvic extent.
- Gonococcal disease prominently causes mucosal suppurative salpingitis.
- Puerperal polymicrobial infection often spreads through deeper tissue.
- Fimbrial and tubal scarring lead to infertility, ectopic pregnancy, and hydrosalpinx.

Exam-level distinction

Pyosalpinx = pus in tube;
hydrosalpinx = obstructed
fluid-filled tube; tubo-
ovarian abscess = localized
adnexal pus.