

Perinatal Brain Injury

A neuropathology lecture on prenatal and perinatal destructive lesions

Hemorrhage • Ischemic white-matter injury • Cortical and deep nuclear patterns

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning outcomes

- Distinguish prenatal or perinatal destructive injury from a primary malformation.
- Identify the major sites, morphology, and consequences of germinal matrix hemorrhage and periventricular leukomalacia.
- Recognize ulegyria, status marmoratus, and their clinicopathologic correlations.
- Explain why a static injury may produce later-emerging motor deficits.

Clinical importance of injury around birth

- Perinatal brain injury contributes importantly to childhood-onset neurologic disability.
- Neurologic consequences depend on the timing, distribution, and severity of tissue destruction.
- Morphologic injury may predate the clinical recognition of a motor disorder.

Timing determines the morphologic response



- Early destructive events may lack the conspicuous reactive response seen after adult brain injury.
- Age at the time of injury is part of the pathologic interpretation.

Destructive injury versus malformation

Interpretive question	Prenatal or perinatal destruction	Developmental malformation
Primary event	Loss of previously developing tissue	Abnormal formation of tissue
Pathologic clue	Hemorrhage, infarction, cystic loss	Abnormal structure may occur without destructive residua
Caution	Early injury may leave little reactive change	Morphologic overlap can be substantial


Why chronology matters

- Injury early in gestation can destroy tissue before conspicuous gliosis develops.
- Gross appearance at birth or later may not reveal when the damage occurred.
- Interpret anatomical loss together with clinical developmental chronology.

Anatomic patterns of perinatal brain injury

Anatomic compartment	Named pattern	Core morphologic feature
Germinal matrix / ventricle	Germinal matrix hemorrhage	Localized or extending hemorrhage
Periventricular white matter	Periventricular leukomalacia	Necrosis, calcification, cysts
Cerebral cortex	Ulegyria	Thinned, gliotic gyral stalks
Basal ganglia and thalamus	Status marmoratus	Irregular, marble-like myelinization

The common destructive pathway



- Prenatal or perinatal ischemia is an important cause of destructive lesions.
- Clinical consequences vary with the injured region; The terms describe distinct morphologic outcomes.

Prematurity is an important context

- Premature infants have greater risk of germinal matrix hemorrhage with brain injury and hypoxic–ischemic events.
- Periventricular leukomalacia is particularly associated with premature infants.
- The injury pattern is therefore interpreted in the context of gestational maturity.

Site determines the lesion name

Site	Event	Recognizable endpoint
Germinal matrix	Bleeding	Hemorrhage; Possible ventricular extension
Periventricular white matter	Infarction	Necrotic foci; Later cysts
Sulcal depths	Cortical ischemia	Ulegyria
Caudate / putamen / thalamus	Deep gray ischemia	Neuronal loss, gliosis, status marmoratus


Cerebral palsy: A clinical descriptor

- The term denotes nonprogressive neurologic motor deficits attributed to prenatal or perinatal brain injury.
- Spasticity, dystonia, ataxia/athetosis, and paresis are characteristic deficits.
- Cerebral palsy is not the name of one specific histologic lesion.

Motor phenotypes represented by cerebral palsy

Term	Type of motor abnormality
Spasticity	Increased muscle tone in the motor phenotype
Dystonia	Abnormal sustained or intermittent muscle contractions
Ataxia / athetosis	Impaired coordination / involuntary writhing movements
Paresis	Weakness of voluntary movement

Why a static lesion can emerge clinically later



- The neurologic lesion is nonprogressive; The phenotype may only declare itself during development.
- Later recognition of symptoms should not automatically be interpreted as a new brain injury.

Cerebral palsy is pathologically heterogeneous

- Autopsy examinations demonstrate a wide range of neuropathologic findings.
- Destructive lesions can reflect remote episodes of hemorrhage or infarction.
- A clinical label alone does not uniquely identify the lesion site or mechanism.

Clinical label and pathology answer different questions

Clinical term	Pathologic assessment
Cerebral palsy describes persistent motor impairment.	Neuropathology identifies the injured region and tissue response.
Phenotype may be recognized as the child develops.	Remote hemorrhage and infarction may explain an established deficit.
Motor signs do not constitute a unique lesion.	Several structural patterns may underlie the same clinical category.

Clinical application: Delayed recognition

A child develops persistent abnormal muscle tone and involuntary movements after an apparently unremarkable early neonatal examination. Prior records suggest an event around delivery.

Interpret the morphologic clues

- What is meant by a nonprogressive motor deficit?
- Why might the abnormal movements become evident after birth?
- Can the motor phenotype alone identify a unique brain lesion?

Interpretation of the delayed-onset phenotype


- Nonprogressive describes the underlying motor-disorder category, not necessarily immediate visibility of signs.
- Functional demands increase as neural motor systems develop.
- Anatomical correlation is needed: Cerebral palsy encompasses varied destructive pathologies.

Germinal matrix hemorrhage: Clinical setting

- In premature infants, brain injury and hypoxic–ischemic events correlate with increased hemorrhage risk.
- The germinal matrix is the site of a characteristic hemorrhagic lesion in the developing brain.
- Hemorrhage severity varies from localized bleeding to extensive intracranial involvement.

The germinal matrix is near the caudothalamic junction

- Typical hemorrhage is near the developing thalamus–caudate junction.
- This periventricular location explains proximity to the ventricular cavity.
- Anatomic site is the distinguishing clue in a premature infant.



Hemorrhage at the caudothalamic junction

Schematic — regional relationships only; Not a specimen

Prematurity, ischemic injury, and hemorrhage

Prematurity

**Hypoxic–ischemic brain
injury**

**Germinal matrix
hemorrhage**

Associated clinical setting — not a universal causal sequence


- Prematurity and hypoxic–ischemic brain injury are associated with increased hemorrhage risk.
- The association does not establish a single obligatory pathway for every hemorrhage.

Localized germinal matrix bleeding

- Some hemorrhages remain small and anatomically confined.
- Bleeding can therefore be present without extension into the ventricular system.
- A localized focus and a ventricular cast represent different extents of injury.

Ventricular extension changes the anatomic compartment

- A hemorrhage may break into the ventricular system.
- Intraventricular blood is not synonymous with an isolated germinal matrix focus.
- Localization and extent should be described separately.



Hemorrhage extends into ventricular space

Schematic — regional relationships only; Not a specimen

Subarachnoid extension of hemorrhage

- Germinal matrix hemorrhage may also extend into the subarachnoid space.
- The ventricular and subarachnoid compartments can both be involved by extensive bleeding.
- This represents spread of hemorrhage rather than a distinct primary disease.

How severe hemorrhage can lead to hydrocephalus



- Hydrocephalus is a recognized complication of severe hemorrhage.
- The intervening CSF pathway is an explanatory schematic, not a statement of a single obligatory mechanism.

Outcome range of hemorrhagic injury

- Minor hemorrhage can remain localized; Severity is not uniform.
- Severe cases may lead to hydrocephalus.
- The most extensive hemorrhages can be fatal.

Anatomic extent of germinal matrix hemorrhage

Distribution	Compartment affected	Potential implication
Focal	Germinal matrix near caudothalamic region	Localized parenchymal hemorrhage
Extended	Ventricular system	Blood in CSF space
More extensive	Subarachnoid space as well	Complicated intracranial hemorrhage
Severe complicated	CSF pathways	Hydrocephalus may develop

Case: Preterm infant with ventricular hemorrhage

A preterm infant has a focal hemorrhage adjacent to the caudothalamic junction. Subsequent examination identifies blood within the ventricles and ventricular enlargement.

Interpret the morphologic clues

- What is the most likely site of hemorrhage origin?
- How has the distribution of bleeding changed?
- Which important complication is suggested by ventricular enlargement?

Germinal matrix hemorrhage: Integrated interpretation


- Context: Prematurity and associated hypoxic–ischemic injury.
- Site: The developing caudate–thalamus junction.
- Morphology: Localized blood or extension into ventricular / subarachnoid spaces.
- Complication: Hydrocephalus or death in severe presentations.

Periventricular leukomalacia: Definition

- A focal ischemic–necrotic lesion of supratentorial periventricular white matter.
- The lesion is particularly encountered in premature infants.
- Its characteristic trajectory can culminate in cystic white-matter destruction.

Anatomical target of periventricular leukomalacia

- Infarcts lie within the white matter adjacent to the ventricles.
- The source identifies supratentorial periventricular distribution.
- PVL is fundamentally a white-matter lesion, not a primary cortical lesion.



Periventricular white-matter necrosis

Schematic — regional relationships only; Not a specimen

Prematurity and periventricular ischemic injury

- Prematurity is an important clinical context for this injury.
- Ischemia produces discrete necrosis in periventricular white matter.
- One should not equate all neonatal ischemic lesions with PVL: Location matters.

Gross appearance: Chalky yellow plaques

- Early or established infarcted foci can appear as chalky yellow plaques.
- These plaques correspond to discrete areas of white-matter necrosis.
- The gross color is a useful descriptive feature, not a substitute for anatomic localization.

Necrosis is the defining tissue lesion

- PVL consists of tissue destruction, not merely pallor of intact white matter.
- Discrete necrotic foci account for later loss of tissue substance.
- Necrotic foci explain how cavitory spaces may remain after injury evolves.

Dystrophic calcification in PVL

- Dystrophic calcification often accompanies the necrotic white-matter foci.
- It reflects mineral deposition in previously damaged tissue.
- Calcification is not described as universal, and its absence would not negate the essential lesion.

Morphologic evolution of severe periventricular injury

01

Ischemic event

The periventricular white matter undergoes infarction.

02

Necrotic focus

Chalky yellow plaques;
Calcification may accompany
necrosis.


03

Chronic lesion

Loss of tissue produces large
cystic spaces.

Chronic periventricular leukomalacia: Cysts

- Severe PVL may leave large cystic spaces within white matter.
- Chronic cavitary lesions may affect periventricular white matter bilaterally.
- The cavities are a chronic consequence of prenatal or perinatal ischemic injury.



Cystic residua in periventricular white matter

Schematic — regional relationships only; Not a specimen

Periventricular injury: Acute and chronic morphology

Phase	Dominant pathologic observation	Interpretation
Infarction	Discrete necrotic white-matter focus	Tissue death
Established lesion	Chalky yellow plaque; Possible dystrophic calcification	Necrotic residuum
Chronic severe lesion	Large cystic spaces	Tissue loss and cavitation

Cyst formation represents tissue loss

- Cavities follow extensive destruction of periventricular white matter.
- The spaces represent a chronic morphologic footprint of prior ischemia.
- Their distribution near the ventricles helps infer the original injury site.

Multicystic encephalopathy is more extensive

- The term applies when damage becomes extensive and includes both gray and white matter.
- It indicates a broader destructive process than a discrete periventricular white-matter infarct.
- The distinction is anatomical extent, not simply the number of visible cysts.

PVL versus multicystic encephalopathy

Feature	Periventricular leukomalacia	Multicystic encephalopathy
Principal distribution	Periventricular white matter	Extensive gray and white matter
Basic event	White-matter infarction / necrosis	Widespread destructive brain injury
Late morphology	Possible large periventricular cysts	Multiple cystic areas with wider tissue involvement

Case: Chronic periventricular cavities

A child with a history of prematurity has chronic bilateral cavitary defects in the white matter immediately adjacent to the lateral ventricles.

Interpret the morphologic clues

- What is the neuropathologic term for this injury pattern?
- Which earlier tissue event produced these cavities?
- What change in distribution would justify the term multicystic encephalopathy?

PVL: What morphology can and cannot establish


- Periventricular distribution plus necrotic or cystic residua supports a destructive injury pattern.
- Morphology may indicate ischemic tissue loss but cannot, alone, pinpoint the precise timing of the insult.
- Always distinguish observed structural features from inferred prenatal or perinatal mechanism.

Perinatal cortical ischemic injury

- A characteristic ischemic lesion is concentrated at the depths of cortical sulci.
- The cortical configuration becomes distorted as injured tissue heals.
- The resulting gyral shape is named ulegyria.

Ulegyria: A characteristic cortical shape

- Ischemic damage disproportionately affects the deep portions of sulci.
- Gyri become mushroom-shaped with narrowed stalks.
- The narrowed tissue contains reactive gliosis.



**Broad-based
gyrus**

**Thinned gliotic
stalk**

Schematic geometry; Cortical sections are not reproduced

The gliotic stalk reflects remote cortical injury

- In ulegyria, the narrow gyral stalk is an outcome of previous destructive ischemia.
- Gliosis indicates a reactive response to tissue injury.
- The broad gyral crown over a thin stalk is a distinctive spatial clue.

Cortical and periventricular ischemia are different patterns


Compartment	Principal morphology	Named term
Deep cortical sulci	Mushroom-shaped gyri; Gliotic stalks	Ulegyria
Periventricular white matter	Necrosis with cystic evolution	Periventricular leukomalacia

Deep gray nuclei are vulnerable

- The basal ganglia and thalamus may sustain perinatal ischemic injury.
- The principal affected deep structures include caudate, putamen, and thalamus.
- Lesions at these sites correlate with movement disorders.

Deep nuclear ischemia: Regional distribution

- The relevant sites are caudate, putamen, and thalamus.
- Patchy injury in these nuclei may coexist with cortical ischemic damage.
- The location helps explain the specific motor phenotype.



Deep nuclear injury (schematic regions)

Schematic — regional relationships only; Not a specimen

Patchy neuronal loss and reactive gliosis


- Ischemic injury of the basal ganglia and thalamus causes patchy loss of neurons.
- Reactive gliosis forms part of the tissue response.
- These changes reflect tissue damage rather than a primary developmental malformation.

Status marmoratus: Abnormal myelin pattern

- In the chronic stage, aberrant and irregular myelinization gives the deep nuclei a marble-like appearance.
- The descriptive term for this pattern is status marmoratus.
- The term denotes abnormal architecture, not a literal deposit of marble-like material.

Why status marmoratus looks marbled


- The distribution of myelin becomes irregular after perinatal injury.
- This creates a visually mottled / marble-like pattern in deep nuclear tissue.
- A schematic can illustrate the principle but cannot substitute for authentic histology.



Abnormal irregular myelin pattern

Schematic only; Not histology

Deep nuclear injury and movement disorder



- Deep nuclear lesions correlate with choreoathetotic movement sequelae.
- The circuit step is a conceptual clinicopathologic inference rather than a measured pathway result.

Two classic ischemic injury signatures

Feature	Ulegyria	Status marmoratus
Primary site	Depths of cerebral cortical sulci	Caudate, putamen, thalamus
Morphology	Mushroom-shaped gyri; Gliotic stalks	Irregular myelinization; Marble-like pattern
Mechanistic theme	Cortical ischemic tissue loss	Deep nuclear ischemia, neuronal loss / gliosis
Clinical link emphasized	Destructive cortical injury	Choreoathetosis / movement disorder

Case: Choreoathetosis following early brain injury

A child with a history of prenatal or perinatal brain injury exhibits persistent choreoathetotic movements. Deep nuclear injury is identified at autopsy.

Interpret the morphologic clues

- Which neuroanatomic structures are implicated?
- What neuronal and glial changes would be expected?
- What does status marmoratus describe?

Across patterns: Clinical and pathologic correlation

Pattern	Site / primary injury	Important clinical or structural consequence
Germinal matrix hemorrhage	Caudothalamic region; Hemorrhage	Ventricular extension; Hydrocephalus
PVL	Periventricular white matter; Ischemia	Cysts after necrotic tissue loss
Ulegyria	Depths of sulci; Ischemia	Abnormal gyral architecture
Status marmoratus	Deep gray nuclei; Ischemia	Movement sequelae including choreoathetosis


Questions for a neonatal brain specimen

- Which region is predominantly involved: Germinal matrix, ventricular space, white matter, cortex, or deep gray nuclei?
- Is the lesion hemorrhagic, necrotic, cystic, gliotic, or characterized by abnormal myelinization?
- Are findings compatible with an immature destructive lesion that may mimic malformation?
- How does lesion distribution relate to the documented motor phenotype?

Interpretive pitfalls in perinatal neuropathology

Pitfall	Corrective reasoning
Assuming an early destructive lesion is a malformation	Consider limited reactive change in early gestation.
Treating cerebral palsy as one morphologic diagnosis	Recognize clinical syndrome with heterogeneous lesions.
Using PVL for any ischemic neonatal brain injury	Require periventricular white-matter involvement.
Treating a marbled pattern as a vascular lesion itself	Status marmoratus describes abnormal myelinization.

A practical lesion-classification pathway



- Site and morphologic evolution provide the most defensible lesion labels.
- Clinical correlation is relevant, but the motor phenotype alone does not establish the neuropathologic diagnosis.

Integrated case: Distinguish four lesion patterns

Four neonatal or childhood neuropathology descriptions are presented: (A) blood near caudothalamic junction; (B) cysts around ventricles; (C) thinned sulcal gyral stalks; (D) marble-like deep nuclear change.

Interpret the morphologic clues

- Identify the named lesion for each description.
- Which two lesions are particularly associated with prematurity?
- Which pattern specifically correlates with choreoathetosis?

Integrated case: Morphologic answers

- A — Germinal matrix hemorrhage; Consider ventricular and subarachnoid extension.
- B — Periventricular leukomalacia with chronic cystic tissue loss.
- C — Ulegyria after ischemic injury centered at sulcal depths.
- D — Status marmoratus in deep nuclei; Possible movement sequelae.

High-yield recognition table

Key observation	Morphologic diagnosis
Premature infant; Caudothalamic hemorrhage	Germinal matrix hemorrhage
Chalky yellow white-matter foci / periventricular cysts	Periventricular leukomalacia
Severe extensive lesions of gray and white matter	Multicystic encephalopathy
Mushroom-shaped gyri with thinned gliotic stalks	Ulegyria
Irregular myelination of deep gray nuclei	Status marmoratus

Final clinicopathologic synthesis

- Timing affects the visibility of tissue reaction, sometimes obscuring the distinction from malformation.
- Premature infants are especially associated with germinal matrix hemorrhage and periventricular leukomalacia.
- Ischemic injury can produce characteristic cortical ulegyria or deep nuclear status marmoratus.
- Cerebral palsy is a nonprogressive motor phenotype arising from heterogeneous destructive injuries.