

Pseudomonal Infections

Opportunism • virulence • vascular injury • chronic airway disease

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraisakhri.com

Learning objectives

By the end of the lecture, explain:

- Why *P. Aeruginosa* is dangerous in selected hosts and care settings.
- How acute toxin-mediated injury differs from chronic biofilm infection.
- The morphology of necrotizing pneumonia and vascular invasion.
- How organ-specific syndromes follow from host and exposure context.

A gram-negative opportunist

Pseudomonas aeruginosa

- Aerobic, gram-negative bacillus.
- Prominent opportunist rather than a uniform pathogen across hosts.
- Potentially severe infection, often challenging to treat.

Opportunistic bacterium



Susceptible host or device



Local infection ± invasive disease

Three high-risk host settings

The primary text emphasizes three clinically important contexts.

Cystic fibrosis

Persistent airway infection and progressive lung injury

Extensive burns


Deep tissue invasion with risk of bacteremia

Neutropenia

Serious invasive infection in an impaired host

Environmental origin and transmission

Infection most often begins with an environmental source.



Person-to-person spread is described as less frequent.

Exposure alone does not determine severity; Host factors matter.

The same organism, different syndromes

Context	Principal disease expression
Cystic fibrosis	Persistent respiratory infection
Intubation	Healthcare-associated respiratory infection
Urinary catheter	Healthcare-associated urinary infection
Skin burn	Deep invasion, possible sepsis
Contact lenses	Corneal keratitis

Host condition shapes clinical expression

The route to infection depends on which host protection is compromised.

Disrupted barrier or device

Intubation • urinary catheter • burn wound

Impaired clearance or defense

Cystic fibrosis • neutropenia

Cystic fibrosis as an airway niche

A defect in airway mucus clearance facilitates infection.

- Thick secretions impair mucociliary removal of bacteria.
- Small airways become obstructed by mucus.
- Persistent organisms encounter a locally altered environment.

Viscous mucus



Poor clearance



P. Aeruginosa persistence

Airway obstruction comes first

The sequence connects host disease to secondary bacterial infection.




The source describes mucus plugging as a frequent antecedent of infection.

Infection amplifies the clinical effects of airway obstruction.


Colonization can become chronic

The Robbins cystic fibrosis discussion describes a progression.



Mucus changes the local microenvironment

Static mucus helps create conditions favoring a mucoid phenotype.



Hypoxia and alginate are explicitly linked in the Chapter 10 text.

This is a local airway mechanism, not generalized systemic hypoxia.

Alginate supports biofilm formation

The bacterial extracellular matrix changes the infection.

- Alginate is a mucoid polysaccharide made by the organism.
- Biofilms contain alginate and other exopolysaccharides.
- A matrix-associated bacterial community becomes persistent.

Alginate / exopolysaccharides



Biofilm organization



Protected bacterial community

A biofilm is more than a bacterial mass

Biofilm organization has biological consequences.

Outside matrix

Bacteria exposed to antibodies,
complement and phagocytes

Within matrix

Reduced access by host defenses and
antibiotics

Biofilms obstruct host clearance

Four host-defense or treatment obstacles.

Antibodies

Reduced effectiveness against embedded bacteria

Complement

Protection within the matrix

Phagocytes

Bacterial persistence despite cells

Antibiotics


Reduced treatment susceptibility

Early infection and chronic infection differ

Early airway infection	Chronic biofilm-associated infection
Exotoxin A active	Exotoxin A output falls
Type III secretion expressed	Type III secretion expression declines
Strong direct toxin effects	Persistent inflammation and tissue enzymes
Acute cell injury	Progressive structural damage

Exotoxin A has an intracellular target

Exotoxin A to diphtheria toxin.



EF-2 is the molecular target stated by the source.

The sequence illustrates the A–B toxin action at a conceptual level.

Loss of protein synthesis is cytotoxic

Toxin-mediated interruption of translation injures the host cell.

- EF-2 is required for normal protein synthesis.
- Exotoxin A modification interrupts that process.
- The end result is loss of host-cell viability.

Modified EF-2




Suppressed translation



Cell death

The type III system delivers effectors

A specialized secretion apparatus transports bacterial effectors into host cells.



Conceptual schematic; Not a microscopic image of the injectisome.

Effector proteins weaken antibacterial ROS

An acute-phase virulence mechanism compromises host-cell defense.

- Type III-delivered effectors diminish antibacterial reactive oxygen species.
- This reduces one route by which host cells oppose the bacterium.
- The effect favors persistence of the early infection.

Effector delivery



Less antibacterial ROS



Impaired local defense

Effector proteins can induce apoptosis

The second effect noted in the source is host-cell apoptosis.

Injected effector proteins



Host-cell apoptotic response



Loss of host cells

The original paragraph is divided across columns in Markdown.

Apoptosis completion appears at robbins.md line 44059.

Acute virulence has several components

Distinct mechanisms converge on local tissue injury.

Exotoxin A

Inhibition of host protein synthesis

Type III effectors

ROS suppression and apoptosis

Proteases


Extracellular tissue injury

Elastases

Additional enzymatic tissue damage

Chronic infection alters virulence expression

Lower expression of several acute-phase features.



Reduced expression should not be interpreted as eradication or harmlessness.

Less acutely virulent does not mean benign

Persistent bacteria still injure lung tissue.

- Biofilm protects the organism against clearance.
- Ongoing inflammation damages the airway wall and surrounding tissue.
- Proteases and elastases directly contribute to injury.

Persistent organisms



Inflammation + enzymes



Chronic tissue destruction

Enzymes contribute to persistent damage

Chronic disease includes toxin-independent injury mechanisms.

Proteases + elastases



Local tissue degradation



Progressive lung injury

Schematic causal relationship described in the source.

Inflammation can be ineffective and harmful

An important host–pathogen paradox in cystic fibrosis.

Immune response continues

Antibody- and cell-mediated responses
against the organism

Bacterial persistence continues

Inflammatory injury despite
inadequate elimination

Two routes to antibiotic resistance

Two interacting explanations for difficult treatment.

Biofilm-related effect

Matrix-associated protection against antibiotics

Genetic effect

Antibiotic resistance caused by bacterial genetic change

Biofilm protection differs from genetic change

Biofilm-related protection	Genetic resistance
Community-level protection by extracellular matrix	Heritable changes in the bacterial population
Limits effectiveness of antibiotics in situ	Adds resistance beyond the matrix effect
Contributes to chronic persistence	Further complicates treatment

Resistance changes clinical difficulty

Why infections may persist despite an active host response and therapy.

- A protected bacterial population is harder to eradicate.
- Genetic adaptation can compound antibiotic resistance.
- Ongoing infection perpetuates inflammation and lung damage.

Biofilm + genetic changes




Treatment difficulty



Chronic pulmonary injury

Chronic cystic fibrosis lung disease

A coherent clinicopathologic chain.



The terminal lesions are explicitly named in the main topic.

Bronchiectasis is a structural endpoint

Chronic infection is linked to abnormal bronchial architecture.

- Bronchiectasis as a consequence of persistent infection.
- Obstructed airways and repeated inflammation precede structural damage.
- The abnormal airway then remains vulnerable to retained secretions.

Chronic infection



Airway wall damage



Bronchiectasis

Pulmonary fibrosis reflects lasting injury

The infection may leave permanent parenchymal damage.

- Repeated inflammation and tissue damage accumulate.
- The main source identifies pulmonary fibrosis as a chronic consequence.
- Fibrosis adds to the persistent functional burden of airway disease.

Chronic damage



Fibrotic remodeling



Long-term lung disease

Hospital-acquired infection depends on context

The primary text gives two characteristic device-linked syndromes.

Intubated airway

Healthcare-associated respiratory infection

Urinary catheter

Healthcare-associated urinary infection

Burn unit

Infected wounds with possible sepsis

Intubation-associated respiratory infection

An invasive airway device identifies a susceptible clinical setting.



This is a context diagram; Specific device-associated causal microbiology is not detailed by the source.

Catheter-associated urinary infection

The same pathogen can infect a different organ system.

- *P. Aeruginosa* causes urinary tract infections associated with urinary catheters.
- This is a healthcare-associated clinical manifestation.
- Host/device history is important when interpreting cultures.

Urinary catheter context




Pseudomonal UTI



Clinical assessment

Severe burns permit invasive infection

Infected burn surfaces can become a source of systemic disease.



Textbook explicitly describes extension deep into veins.

Vascular invasion is a central theme

Pseudomonal disease repeatedly targets vessels.

- In burns, organisms penetrate deeply into veins.
- In pneumonia, organisms may concentrate in vessel walls.
- Vascular injury is linked to necrosis, thrombosis and hemorrhage.

Bacterial vessel involvement




Vascular wall damage



Thrombotic + hemorrhagic injury

Why hematogenous spread matters

Local infection may escalate to systemic complications.



Complemented by Chapter 15 description of fulminant *Pseudomonas* septicemia.

Ecthyma gangrenosum is a vascular clue

Characteristic skin lesion described in disseminated infection.

- Well-demarcated, oval lesions are emphasized.
- They are necrotic and hemorrhagic.
- The lesion is associated with burn infection and hematogenous spread in the source.

Invasive infection



Skin necrosis + hemorrhage



Ecthyma gangrenosum

Bacteremia can provoke DIC

The main text calls disseminated intravascular coagulation a frequent complication.

P. Aeruginosa bacteremia



**Systemic coagulation
derangement**



DIC

No numerical incidence is inferred.

Risk and lesion pattern together

Predisposing context	Associated important pathology
Neutropenia	Severe invasive Pseudomonas infection
Burn wound	Venous penetration, dissemination, ecthyma gangrenosum
Cystic fibrosis	Persistent infection, bronchiectasis, pulmonary fibrosis
Hospital intubation	Respiratory infection / pneumonia

Necrotizing pneumonia: Gross pattern

The defining morphology combines necrosis and hemorrhage.

- Distribution follows terminal airways.
- Pale zones represent necrotic centers.
- Red peripheral zones represent hemorrhage.

Terminal airway distribution




Pale necrotic center



Red hemorrhagic edge

Fleur-de-lis describes lesion distribution

The source uses a branching, terminal-airway pattern.



A relational schematic, not a literal gross image.

Necrosis and hemorrhage coexist

Two distinct injury components appear in the same lesion.

Pale center

Tissue necrosis

Red peripheral zone

Hemorrhagic tissue

Microscopic bacillary aggregates

The source highlights organism-rich microscopic fields.

- Organisms may be present in large masses.
- They are often concentrated in blood-vessel walls.
- The finding should be interpreted with the associated vascular injury.

Numerous bacteria




Vessel-wall localization



Damaged host tissue

Coagulative necrosis of host tissue


Bacteria in vascular walls are associated with necrosis.



This diagram depicts causal text; Not a histologic specimen.

Thrombosis amplifies vascular pathology

The microscopic description links inflammation with thrombi and hemorrhage.



The distal phrase is a clinicopathologic inference from vascular occlusion; The main text states thrombosis and hemorrhage.

The vascular morphology triad

A suggestive constellation in the appropriate host.

Gram-negative bacilli

Organism-rich vessels

Necrotizing vasculitis

Host-cell coagulative necrosis

Thrombosis


Occluded injured vessels

Hemorrhage

Extravascular bleeding

Schematic: Infected vessel wall

Vessel-wall localization is the central morphologic relationship.



Bacterial aggregates

Vessel-wall injury

Luminal thrombus

Hemorrhage

Schematic — not a histologic specimen

Vascular pattern is suggestive, not specific

Avoid equating a characteristic pattern with a definitive organism identification.

Supported interpretation

Strongly suggests *P. Aeruginosa* when context fits

Overstatement

Histology alone proves the bacterial species

Pathology links lung and skin disease

Lung	Skin / burns
Terminal airway-centered necrosis	Deep infection of burned tissue
Organism-rich vessel walls	Venous penetration
Thrombosis and hemorrhage	Necrotic hemorrhagic skin lesions
Risk of invasive spread	Risk of bacteremia and DIC

The eye: Contact lens–associated keratitis

A local infection in a host without classic systemic immunodeficiency.

- Contact lens wear is the characteristic exposure in the source.
- *P. Aeruginosa* can infect the cornea.
- Anatomic site and exposure narrow the clinical suspicion.

Contact lens exposure



Corneal infection



Keratitis

Otitis externa spans host groups

Clinical group	Syndrome emphasized
Otherwise healthy person	External otitis (“swimmer’s ear”)
Person with diabetes	Severe external otitis

Injection drug use: Invasive foci

Two focal infections are named in the main source.

- *P. Aeruginosa* may cause endocarditis.
- Osteomyelitis is another recognized manifestation.
- Exposure context raises suspicion for deep-seated infection.

Injection drug exposure



Endocarditis or osteomyelitis




Deep focal infection

Clinical phenotype is host-specific

Host / exposure	Characteristic presentation
Cystic fibrosis	Chronic airway colonization and damage
Severe burns	Invasive wound infection $\pm$ bacteremia
Contact lens use	Keratitis
Diabetes	Severe external otitis
Injection drug use	Endocarditis; osteomyelitis

Reasoning from host to suspected pathogen

Clinical suspicion uses context before microscopic confirmation.



This is an educational reasoning framework, not a validated diagnostic algorithm.

Avoid mistaking morphology for identification

Interpret tissue findings in a clinically coherent way.

- Vasculitis with abundant gram-negative organisms, thrombosis and hemorrhage is strongly suggestive.
- This pattern is not pathognomonic.
- Species identification therefore requires organism-directed microbiologic evidence.

Histologic evidence and clinical context

Neither category should be interpreted in isolation.

Tissue pattern

Vascular organism masses •
coagulative necrosis • hemorrhage

Clinical setting

Neutropenia • burns • CF • intensive-
care exposure

A practical syndrome–mechanism matrix

Syndrome	Mechanistic emphasis
Early lung infection	Exotoxin A / type III effectors
Chronic CF lung infection	Alginate biofilm / inflammation / enzymes
Necrotizing pneumonia	Angioinvasion with hemorrhagic necrosis
Sepsis after burns	Vascular invasion and dissemination

Case: Progressive lung disease in cystic fibrosis

A person with cystic fibrosis has persistent *P. Aeruginosa* infection despite treatment, with progressive airway damage.

Which transition best explains persistence?

A. Formation of alginate-associated biofilm

B. Complete loss of bacterial tissue-damaging capacity

Case: Necrotic lesions after severe burns

After extensive burns, a patient develops well-circumscribed oval lesions containing necrosis and hemorrhage.

What systemic complication is especially important to consider?

A. Pseudomonal bacteremia with possible DIC

B. An isolated noninvasive superficial eruption

Case: Hemorrhagic necrotizing pneumonia

A neutropenic patient develops rapidly progressive pneumonia. Tissue shows abundant gram-negative bacilli concentrated around vessel walls, with thrombosis and hemorrhage.

How specific is the morphologic pattern?

A. Highly suggestive, not pathognomonic

B. Definitive organism identification from morphology alone

Two modes of pseudomonal injury

A concise unifying model.

Acute / invasive

Exotoxin A • type III effectors •
vascular necrosis

Persistent / chronic

Alginate biofilm • inflammatory injury
• enzymes

Integrated clinicopathologic summary

Four associations capture the teaching unit.

Host

CF, burns and neutropenia drive major susceptibility

Pathogenesis

Early toxins → chronic biofilm adaptation

Morphology

Necrotizing hemorrhagic pneumonia and vasculitis

Consequences

Bronchiectasis, fibrosis, sepsis and DIC

Source basis

Primary textbook

Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th edition (2026).

Pseudomonal Infections, in Infectious Diseases.

Related reading within Robbins: cystic fibrosis and bacterial pneumonia.