

Reactions of Neurons to Injury

Morphologic patterns • mechanisms • clinicopathologic interpretation

Neuronal cell death, degeneration, axonal responses, and inclusions

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Classify neuronal responses by time course and affected compartment.
- Identify the defining changes of acute “red neurons.”
- Explain synaptic loss, neuronal degeneration, gliosis, and axonal reaction.
- Interpret major types of neuronal inclusions and distal axonal degeneration.

Pattern recognition

Acute injury


Degeneration

Axonal reaction

Inclusions

The neuron is the principal CNS functional unit

- Neurons vary in electrical activity, transmitter use, and connectivity.
- Their functional and metabolic requirements differ across regions.
- Injury to a small strategically placed population may have major consequences.



Some key features of neurons:

Function depends on connections

Cell type and location matter

Loss cannot be quickly replaced

Selective vulnerability reflects neuronal diversity

- Different neuron classes do not respond identically to a given insult.
- Shared metabolic needs or connections can define a vulnerable population.
- The resulting lesion pattern may follow function rather than a single anatomic cluster.

Morphologic sequence

Shared properties



Shared vulnerability



Selective cell loss



Characteristic deficit

Anatomic location shapes clinical consequences

- Different CNS regions mediate different functions.
- Identical cellular injury can yield distinct syndromes at different sites.
- Localization and the pathologic process must be interpreted together.

Pattern recognition

Anatomic site


Neuronal role

Injury pattern

Clinical effect

Mature neurons have limited replacement capacity

- Mature neurons are postmitotic and cannot proliferate.
- Their integrity must be preserved over the lifespan.
- Destruction of neurons essential to a particular function can cause lasting dysfunction.



Some characteristics of mature neurons

Long-lived cell

No proliferative reserve

Loss may create a deficit

Neuronal injury has multiple timescales

- Acute insults can produce rapidly evolving injury.
- Subacute degeneration progresses over days or weeks.
- Chronic degenerative disease evolves over months or years.

Pattern and temporal context

Acute

Hours after irreversible insult

Subacute

Days to weeks

Chronic

Months to years

Injurious contexts span metabolism and trauma

- Acute examples include oxygen/glucose depletion and brain trauma.
- Severe hypoglycemia is an established cause of acute neuronal injury.
- Chronic disorders include neurodegenerative conditions of the brain.

Distinguishing patterns

Acute insults

Hypoxia / ischemia

Severe hypoglycemia, trauma

Long-duration disease

Neurodegenerative processes

Months to years

Continuous metabolic supply is critical

- Neurons are highly metabolically active.
- They depend on continuous oxygen and glucose delivery.
- Interruption of this supply may precipitate acute neuronal injury.

Morphologic sequence

Oxygen + glucose supply



Neuronal metabolism



Acute insufficiency



Cellular injury

Long lifespan creates proteostasis challenges

- Neurons maintain cellular proteins over unusually long periods.
- Misfolded proteins may accumulate in vulnerable neurons.
- This accumulation triggers the unfolded protein response and is implicated in degenerative CNS disease.

Morphologic sequence

Long-lived neuron



Protein misfolding



Unfolded protein response



Degenerative injury

Cellular injury produces distinguishable patterns

- Assess the cell body, nucleus, cytoplasm, synapses, and axon.
- Identify whether the process suggests death, degeneration, or axonal regeneration.
- Use surrounding glial reaction as supporting context, not a stand-alone etiology.

Pattern recognition

Soma

Nucleus


Processes

Background glia

Acute “red neurons” signal irreversible injury

Comparative schematic

- This is an acute morphologic reaction to severe CNS insult.
- It is among the earliest visible manifestations of neuronal necrosis.
- The diagnosis rests on a constellation of nuclear and cytoplasmic features.



Viable-appearing

Acute “red neuron”

Depicts key features conceptually; Not H&E imagery

Acute red-neuron change has multiple causes

- Acute CNS hypoxia/ischemia can produce red neurons.
- Severe hypoglycemia can produce a similar phenotype.
- Other acute insults may also cause this pattern; It is not etiologically specific.

Distinguishing patterns

Common setting

Hypoxia / ischemia
Red-neuron morphology

Other acute injury

Severe hypoglycemia
Same general pattern

Red neurons have a characteristic time context

- After irreversible hypoxic/ischemic injury, red neurons become evident at 6–12 hours.
- This timing pertains to that insult, not every possible neuronal injury.
- Interpret timing alongside clinical history and tissue context.

Pattern and temporal context

Acute

Hours after irreversible insult

Subacute


Days to weeks

Chronic

Months to years

Normal neuronal morphology is the reference

- Recognize a discernible neuronal soma and nucleus.
- Basophilic Nissl substance is present in the cell body.
- Acute injury is identified by simultaneous departure from these baseline features.



Step 1: Identify the normal morphology

Cell body + nucleus


Nissl substance

Compare with adjacent neurons

Soma shrinkage and eosinophilia mark red neurons

Comparative schematic

- The injured neuronal cell body contracts.
- The cytoplasm becomes intensely eosinophilic on H&E.
- Together, these changes produce the classic small, deeply pink appearance.



Viable-appearing


Acute "red neuron"

Depicts key features conceptually; Not H&E imagery

Nuclear pyknosis is a key acute feature

Comparative schematic

- The nucleus becomes pyknotic: Condensed and intensely staining.
- The normal nucleolus is no longer apparent.
- This nuclear change accompanies other indicators of neuronal necrosis.



Viable-appearing


Acute "red neuron"

Depicts key features conceptually; Not H&E imagery

Loss of Nissl substance accompanies acute injury

- Nissl substance disappears in the acutely injured neuron.
- The lost basophilic cytoplasmic detail accentuates eosinophilia.
- Assess it with nuclear pyknosis and soma shrinkage, not in isolation.

Comparative schematic



Viable-appearing

Acute "red neuron"

Depicts key features conceptually; Not H&E imagery

A red neuron is a composite morphologic diagnosis

- Look for shrinkage, pyknotic nucleus, absent nucleolus, and Nissl loss.
- Intense cytoplasmic eosinophilia completes the phenotype.
- A single nonspecific staining change should not substitute for the full pattern.

Pattern recognition

Soma shrinks

Nucleus pyknotic

Nissl lost

**Cytoplasm
eosinophilic**

Acute neuronal necrosis has a morphologic sequence

- A severe insult precedes visible cell-body alterations.
- Red-neuron morphology is an early manifestation of irreversible injury.
- The visible change is delayed relative to insult onset in hypoxic/ischemic injury.

Morphologic sequence

Acute severe insult



Irreversible injury / necrosis



Red neuron visible at 6–12 h*



Morphologic evidence

Red neurons should not be read as a unique etiology

- Similar morphology occurs in ischemia and severe hypoglycemia.
- Etiologic assignment therefore needs clinical and anatomic context.
- Neuronal necrosis is a cellular response, not a disease-specific diagnosis.

Distinguishing patterns

Morphology

Red-neuron features

Identifies acute neuronal injury

Etiologic diagnosis

Injury context

Requires external evidence

Clinical correlation: Acute ischemic neuronal insult

- Scenario: Oxygen delivery to CNS tissue is interrupted.
- Morphology later reveals small, eosinophilic neurons with pyknotic nuclei.
- Interpretation: Acute irreversible neuronal damage is supported; Mechanism is consistent with ischemia.

Interpretive task

Which cell-level findings support necrosis? Why does the context still matter?

Red-neuron phenotype

Clinical correlation: Severe hypoglycemia

- Scenario: An acute episode of severe glucose depletion precedes neurologic injury.
- Red neurons can appear despite no primary arterial occlusion.
- The shared morphology reflects neuronal metabolic vulnerability rather than one exclusive cause.

Interpretive task

Can a red-neuron pattern distinguish ischemia from severe hypoglycemia?

Etiology is not unique

Timing pitfalls in acute neuronal injury

- The 6–12-hour estimate is specific to irreversible hypoxic/ischemic injury.
- An early specimen may not yet display classic red-neuron morphology.
- Lack of a particular marker at one sampled time cannot exclude all injury.

Pattern and temporal context

Acute

Hours after irreversible insult

Subacute

Days to weeks

Chronic

Months to years

Subacute and chronic injury constitute degeneration

- Subacute degeneration involves neuronal death over days to weeks.
- Chronic degeneration accompanies progressive disorders over months to years.
- The sequence includes synaptic loss, eventual cell loss, and reactive gliosis.

Pattern and temporal context

Acute

Hours after irreversible insult

Subacute

Days to weeks

Chronic

Months to years

Synapses can disappear before neuronal somata

- Synapses are the junctions for interneuronal communication.
- Loss of synapses is often an early event in degenerative disease.
- Functional impairment may precede conspicuous neuronal depletion in tissue sections.

Sequence of neuronal degeneration

Synapses lost

Communication declines



Neurons die

Often apoptosis



Gliosis

Secondary tissue response

Synaptic pruning has normal developmental roles

- Pruning removes selected synaptic connections in normal development.
- It also contributes to nervous-system plasticity.
- The pathology discussion implicates aberrations in this process in some synaptic loss.

Morphologic sequence

Normal synaptic pruning



Development / plasticity



Aberrant regulation



Potential pathologic loss

Aberrant pruning may contribute to disease

- Aberrant synaptic pruning as a possible mechanism of synaptic loss.
- This does not imply every neurodegenerative disorder begins with pruning.
- Link a proposed mechanism to demonstrable morphologic consequences.

Distinguishing patterns

Physiologic process

Synaptic pruning
Development and plasticity

Possible pathologic change

Aberrant pruning
Associated synapse loss

Early degeneration can be difficult to see

- Initial synapse loss may not create striking neuronal soma depletion.
- Routine examination may underestimate early functional injury.
- Reactive glial changes can make underlying neuronal injury more apparent.

Sequence of neuronal degeneration

Synapses lost

Communication declines



Neurons die

Often apoptosis



Gliosis

Secondary tissue response

Chronic disorders can selectively affect neuron groups

- Neuronal apoptosis may preferentially involve functionally related populations.
- Cell loss need not be uniform throughout the CNS.
- Patterns of dysfunction reflect both cellular vulnerability and neuroanatomy.

Pattern recognition

Selective neurons

Related networks

Progressive cell loss

Functional deficits

Apoptotic death is part of degeneration

- The source describes apoptosis following synaptic loss in chronic disease.
- Cell death is accompanied by reactive changes in the tissue.
- Contrast this sequence with acute neuronal necrosis and red-neuron change.

Distinguishing patterns

Acute severe injury

Red-neuron necrosis

Hours after ischemia

Slow degeneration

Synapse loss → apoptosis

Days to years

Reactive gliosis can reveal neuron loss

- Astrocytic hypertrophy and hyperplasia define gliosis.
- It is a major histopathologic indicator of CNS injury.
- Gliosis may stand out while early neuronal depletion remains hard to appreciate.

Morphologic sequence

Neuronal/synaptic injury



Reactive astrocytes



Gliosis



Histologic clue

Gliosis is an indicator, not a specific diagnosis

- Reactive astrocytes occur in many types of CNS injury.
- The presence of gliosis alone does not identify the injury mechanism.
- Interpret glial changes together with neuron, axon, and disease distribution.

Distinguishing patterns

Positive evidence

CNS tissue injury

Reactive gliosis

Cannot infer alone


Specific etiology

Requires context

Alzheimer disease illustrates chronic degeneration

Intracellular change — schematic

- Alzheimer disease is given as an example of slow neuronal degeneration.
- Neurofibrillary tangles are a characteristic neuronal cytoplasmic inclusion.
- Injury may involve synaptic loss, neuron death, and gliosis over time.




Conceptual neuronal protein inclusion

Conceptual drawing; Cannot replace diagnostic microscopy

ALS illustrates selective progressive neuron loss

- Amyotrophic lateral sclerosis is another example of chronic neuronal degeneration.
- The injury evolves over a prolonged interval rather than as a single acute insult.
- Progressive loss of a vulnerable functional neuronal population can produce deficits.



The degeneration sequence connects morphology to function

- Loss of synaptic contact disrupts neuronal communication.
- Subsequent apoptotic cell loss reduces the functioning cell population.
- Reactive gliosis gives histologic evidence of the underlying injury.

Sequence of neuronal degeneration

Synapses lost

Communication declines



Neurons die

Often apoptosis



Gliosis

Secondary tissue response

Acute and chronic neuronal injury differ

Distinguishing patterns

- Acute injury: Red-neuron changes and necrosis associated with severe insult.
- Chronic degeneration: Synaptic reduction followed by neuron loss and gliosis.
- The temporal distinction helps structure a differential, but causes may overlap.

Acute

Red neurons

Necrosis after severe insult

Chronic


Synapse loss + gliosis

Progressive degeneration

Axonal reaction is a somatic response to axon injury

Somatic response and distal axon

- Injury of a neuron's axon can alter its cell body.
- Axonal reaction is particularly seen when regeneration is occurring.
- This is a distinct process from the small, eosinophilic dying red neuron.




Soma: Chromatolysis

Distal segment:
Degeneration

Anterior horn cells exemplify axonal reaction

Somatic response and distal axon

- The classic example is a spinal anterior horn motor neuron.
- Its motor axon has been severed or seriously damaged.
- The neuronal soma mounts a response visible by light microscopy.



Injured axon

Soma: Chromatolysis

Distal segment:
Degeneration

Axonal reaction reflects increased protein synthesis

- The injured neuron increases its synthetic activity.
- Cell-body enlargement and nuclear/nucleolar changes accompany this response.
- The morphology represents an attempt to support axonal regeneration.

Morphologic sequence

Axon cut/damaged



Protein synthesis rises



Somatic remodeling



Regenerative response

Cell-body enlargement characterizes axonal reaction

- The soma becomes larger than its resting appearance.
- It assumes a more rounded outline.
- This contrasts with the shrunken soma of the acutely necrotic red neuron.

Distinguishing patterns

Axonal reaction

Enlarged + rounded soma

Regenerative response

Red neuron


Shrunken soma

Acute necrotic injury

Nuclear displacement supports axonal reaction

Somatic response and distal axon

- The nucleus shifts toward the periphery of the enlarged soma.
- The nucleolus becomes enlarged and conspicuous.
- These features should be evaluated with the altered Nissl distribution.




**Distal segment:
Degeneration**

Soma: Chromatolysis

Central chromatolysis redistributes Nissl substance

- Nissl substance disperses away from the central cell body.
- Remaining Nissl material is displaced toward the neuronal periphery.
- The central pale region contrasts with the Nissl loss of red-neuron injury.

Somatic response and distal axon



Soma: Chromatolysis

**Distal segment:
Degeneration**

Axonal reaction is a recognizable feature cluster

- Enlarged, rounded neuronal soma.
- Peripheral displacement of the nucleus with enlarged nucleolus.
- Peripheral redistribution of Nissl material—central chromatolysis.

Pattern recognition

Enlarged soma

Off-center nucleus

Large nucleolus

**Central
chromatolysis**

Chromatolysis and red neurons are not equivalent

- Red neuron: Shrunken eosinophilic soma and pyknotic nucleus.
- Axonal reaction: Enlarged soma with peripheral nucleus and nucleolus.
- Nissl loss in acute necrosis differs from redistribution in chromatolysis.

Distinguishing patterns

Acute red neuron

Soma shrinks

Pyknosis + Nissl loss

Axonal reaction


Soma enlarges

Eccentric nucleus + Nissl redistribution

Wallerian degeneration occurs distal to axonal injury

Somatic response and distal axon

- Wallerian degeneration is degeneration of the distal axon after acute axonal injury.
- It is a process involving the axon beyond the lesion.
- Distinguish it conceptually from axonal reaction in the injured neuronal soma.



**Distal segment:
Degeneration**

Soma: Chromatolysis

Axon damage affects more than one compartment

- The proximal neuron may show axonal reaction and central chromatolysis.
- The distal axonal segment may undergo Wallerian degeneration.
- The two terms describe responses in different parts of the same neuron.

Morphologic sequence

Axonal lesion



Somatic axonal reaction



Distal Wallerian change



Compartment-based interpretation

Case: Identify the consequence of axon transection

- Scenario: A motor axon is seriously injured.
- The cell body enlarges, its nucleus moves peripherally, and Nissl substance disperses.
- Interpretation: A regenerative axonal reaction, not red-neuron necrosis.

Interpretive task

What term describes the somatic change? What happens in the distal axon?

Chromatolysis + Wallerian

Subcellular neuronal alterations are diverse

- Damage may affect neuronal organelles and the cytoskeleton.
- Accumulations can involve lipids, proteins, or carbohydrates.
- Some inclusions accompany aging; Others point toward infectious, metabolic, or degenerative disease.

Pattern recognition

Organelles

Cytoskeleton


Accumulations

Vacuolation

Lipofuscin is an age-associated neuronal inclusion

Intracellular change — schematic

- Complex lipid accumulation can occur with advancing neuronal age.
- Lipofuscin is an intracytoplasmic storage pigment.
- An inclusion in a long-lived neuron is not automatically evidence of a specific neurodegenerative disorder.




Schematic coarse pigment granules

Conceptual drawing; Cannot replace diagnostic microscopy

Inherited metabolic defects cause storage changes

Intracellular change — schematic

- Some inborn errors of metabolism result from loss of specific enzyme activity.
- Neurons may accumulate abnormally stored lipids or other substrates.
- Storage material is a category of morphologic change, not a single disease diagnosis.




Conceptual abnormal cytoplasmic accumulation

Conceptual drawing; Cannot replace diagnostic microscopy

Viral infection may produce neuronal inclusions

Intracellular change — schematic

- Certain viral infections produce characteristic cellular inclusions.
- Robbins gives herpes, rabies, and cytomegalovirus as examples.
- Type of inclusion and associated pathology support etiologic interpretation.



Stylized viral inclusion—NOT microscopy

Conceptual drawing; Cannot replace diagnostic microscopy

Viral inclusion names must be used precisely

- Herpetic infection: Cowdry A or B bodies.
- Rabies: Negri bodies.
- Cytomegalovirus can also cause viral inclusions in relevant infected cells.

Pattern recognition

Herpes → Cowdry

Rabies → Negri


CMV → inclusion

Context matters

Degenerative diseases may form protein inclusions

Intracellular change — schematic

- Alzheimer disease can show neuronal neurofibrillary tangles.
- Parkinson disease can show neuronal Lewy bodies.
- These are distinct named pathologic structures in different disease contexts.




Schematic intraneuronal aggregate

Conceptual drawing; Cannot replace diagnostic microscopy

Prion disease illustrates neuronal vacuolization

Intracellular change — schematic

- Creutzfeldt–Jakob disease is associated with abnormal vacuolation.
- Vacuoles involve the neuronal perikaryon and processes within neuropil.
- Vacuolization is distinct from an intracytoplasmic protein inclusion.



Conceptual intracellular vacuolation

Conceptual drawing; Cannot replace diagnostic microscopy

Intracellular changes require pattern-based classification

- Aging can yield lipofuscin pigment accumulation.
- Inherited metabolic disease can produce abnormal substrate storage.
- Infections, neurodegeneration, and prion disease have different characteristic changes.

Distinguishing patterns

Accumulated material

Pigment / lipid / protein

Aging, storage, degeneration

Altered structure

Viral inclusions / vacuoles

Infection or prion disease

Interpret neuronal injury using a structured approach

- First distinguish acute cell death, chronic degeneration, and axonal response.
- Then locate the abnormality: Soma, synapse, axon, or inclusion.
- Finally integrate the lesion with timing, neuroanatomy, and reactive gliosis.

Morphologic sequence

Classify time course



Identify cell compartment



Recognize morphology



Integrate clinical context

Application: Early progressive dysfunction

- Scenario: Long-standing progressive neurologic decline with limited obvious neuron loss.
- A histologic section shows prominent reactive astrocytes.
- Synaptic loss and early degeneration remain plausible despite subtle neuronal depletion.

Interpretive task

Why can gliosis be more conspicuous than neuronal loss? Is it specific?

Early synaptic injury

Application: Somatic morphology points to axon injury

- Scenario: An anterior horn neuron is enlarged with an eccentric nucleus.
- The nucleolus is prominent and Nissl granules are peripheral.
- This is central chromatolysis in an axonal reaction, not a red neuron.

Interpretive task

Identify the pattern and contrast its nucleus with that of a red neuron.

Axonal reaction

Application: Interpret neuronal inclusions

- Scenario: A biopsy shows neuronal inclusions during investigation of a CNS disorder.
- A named inclusion suggests a diagnostic category but does not replace context.
- Differentiate aging, inherited storage, viral infection, and neurodegeneration.

Interpretive task

Which categories of inclusions are linked to herpes, rabies, Alzheimer, and Parkinson?

Disease–inclusion mapping

Common diagnostic pitfalls

- Red-neuron change is not exclusive to ischemia.
- Gliosis is a sensitive but nonspecific reaction to diverse CNS injuries.
- Central chromatolysis is not the same as the loss of Nissl substance in acute necrosis.
- Schematic drawings in this deck must not be treated as patient-derived histology.

Pattern recognition

Pattern ≠ cause

Timing matters

**Compartment
matters**

Context required

Core morphologic distinctions

- Acute necrotic injury: Red neuron with shrinkage, pyknosis, and eosinophilia.
- Subacute/chronic degeneration: Synaptic loss, apoptotic death, and gliosis.
- Axonal injury: Somatic chromatolysis and distal Wallerian degeneration.
- Inclusions and vacuoles: Additional morphologic clues to specific disease classes.

Pattern recognition

Acute

Chronic

Axonal

Inclusion

Textbook foundation and evidence limits

- Primary textbook: Robbins, Cotran & Kumar, Pathologic Basis of Disease, 11th edition (2026).
- Chapter 28, “Cellular Pathology of the Central Nervous System,” pp. 1130–1131.
- Slides paraphrase the supplied text; original figure 28.15A and chapter PDF were not supplied.
- Neuronal drawings are original editable teaching schematics, not histopathology images.

Pattern recognition

Source-based

**No invented
datasets**

Editable schematics

**Morphology
focused**