


Release From the Body and Transmission of Microbes

Mechanisms of microbial exit, exposure, and host entry



Dr Abu Suraih Sakhri

MBBS, MD (Pathology)


www.abusuraihsakhri.com

Learning objectives

Explain how pathogen exit routes generate new host exposures.

- Distinguish release from a host, mode of transmission, and portal of entry.
- Compare respiratory, fecal–oral, sexual, vertical, and additional routes.
- Explain how asymptomatic shedding, barrier defenses, and the microbiome alter transmission.
- Apply route-based reasoning to selected clinical scenarios.

Transmission connects two hosts



- A route must connect an exit source with exposure of another susceptible host.
- Successful exposure does not invariably produce infection; Host defenses remain important.


Exit strategy is part of microbial success

Microbes leave the body through routes that make onward transmission possible.

- Exit can involve respiratory secretions, stool, genital secretions, saliva, milk, or vector feeding.
- The dominant exit route depends on where the organism is present and can be released.
- Release and transmission are distinct events: One does not guarantee the other.

Three patterns of microbial shedding

Brief



Release is concentrated during a limited interval.

Intermittent



Some microbes are shed periodically during disease flares.

Prolonged



Long-term release can occur even without clinical symptoms.

Asymptomatic hosts can be a source

A person may shed an organism despite having no evident illness.

- Robbins explicitly describes prolonged shedding by asymptomatic hosts.
- Symptom-based identification alone therefore may miss a source of transmission.
- Infection status, symptoms, and the period of microbial release are not equivalent.

Release, transmission, and entry are different

Release

Microbe leaves a host.
Example: Organism in
respiratory droplets.

Transmission

Microbe reaches another
host or exposure setting.
Example: Inhalation of
infectious particles.

Entry

Microbe crosses or
adheres to a host surface.
Example: Respiratory
mucosal infection.

These are linked stages—not interchangeable labels.

The principal transmission routes

Person-to-person transmission

```
graph TD; A[Person-to-person transmission] --> B[Respiratory]; A --> C[Fecal-oral]; A --> D[Sexual]; A --> E[Vertical]; B --> B1[Microbe released into the air]; C --> C1[Stool contaminates food or water]; D --> D1[Intimate or mucosal contact]; E --> E1[Mother to fetus or newborn];
```

Respiratory

Microbe released into the air

Fecal-oral

Stool contaminates food or water

Sexual

Intimate or mucosal contact

Vertical

Mother to fetus or newborn

Additional routes extend the framework

Saliva

Release of organisms replicating in salivary glands or oropharynx.


Arthropod vector

A blood meal can move some protozoan pathogens between hosts.

Zoonotic transfer


Animal contact, animal products, bites, or vectors transmit microbes.

The route must be interpreted with host defenses



- Virulence factors can help pathogens overcome intact defenses.
- When host defenses are compromised, normally low-virulence microbes may establish infection.

Respiratory release creates airborne particles



- Viruses and bacteria may be released in droplets or smaller airborne particles.
- Particle behavior helps determine where an infectious exposure may occur.

Respiratory transmission is not one distance category

Droplet-predominant example


Influenza is in association with relatively short-range droplets.

Longer-range particle example

Mycobacterium tuberculosis can spread in particles that travel farther from the source.


Examples illustrate differences in dispersal; A single distance rule is not universal.

Influenza: A respiratory transmission example




- Influenza among short-range droplet examples.

Tuberculosis: Airborne dispersal matters




- The long-range example is M. Tuberculosis.

Particles encounter distinct airway defenses



- Larger inhaled particles may be trapped in upper-airway mucus.
- Ciliary transport moves trapped material toward the pharynx.

Smaller inhaled particles can reach alveoli



- The source describes particles below 5 μm reaching alveoli.
- Alveolar leukocytes normally phagocytose inhaled microorganisms.

Respiratory infection requires a foothold

Epithelial attachment

Influenza viruses attach to and replicate in airway epithelial cells.

Ciliary injury

Bordetella pertussis toxins can impair ciliary activity.

Intracellular survival

M. Tuberculosis can survive within host macrophages.

When mucociliary clearance fails


Impaired airway defenses change susceptibility after exposure.

- Influenza, smoking, mechanical ventilation, and cystic fibrosis may compromise clearance.
- The resulting defect predisposes to bacterial respiratory infections.
- Route of acquisition and the host's capacity to clear the organism must be assessed together.

Respiratory exposure: Integrate exit and entry

Component	Source-based example	Biologic implication
Exit	Infectious respiratory particles	Microbe becomes available for inhalational spread
Transport	Shorter-range / farther-traveling particles	Distance and particle behavior influence exposure
Entry	Airway epithelium or alveoli	Local defenses determine establishment
Host factor	Defective mucociliary clearance	Greater opportunity for airway infection

The fecal–oral route is a connected sequence



- Enteric pathogens commonly spread when stool contaminates ingested material.
- Safe appearance of food or water does not define whether it is contaminated.

Waterborne enteric spread

Contaminated drinking water is an efficient vehicle for enteric pathogens.

- Waterborne spread as central to epidemic enteric infections.
- Examples include hepatitis A/E viruses, poliovirus, rotavirus, and *Vibrio cholerae*.
- The shared route does not imply identical clinical disease.

Foodborne fecal contamination



- The source groups contaminated food and water under fecal–oral transmission.

Viral examples of fecal–oral spread

Hepatitis A / E

Enteric exposure can transmit hepatotropic viral pathogens.

Poliovirus

Fecal–oral transmission may result in systemic infection.

Rotavirus

Representative enteric virus transmitted through the fecal–oral route.

Bacterial examples of fecal–oral spread

Vibrio cholerae

A waterborne pathogen in the Robbins example set.


Shigella spp.

Enteric bacteria named among epidemic fecal–oral pathogens.

Salmonella enterica

Included among fecal–oral epidemic pathogen examples.

Ingestion is followed by gastric defenses



- Gastric acidity helps eliminate many ingested organisms.
- Susceptibility depends in part on an organism's capacity to survive this barrier.

Gastric pH changes susceptibility to cholera

A strong experimental effect of neutralizing gastric acid.

- In healthy volunteers, neutralizing gastric acid increased infectivity of *Vibrio cholerae* by 10,000-fold.
- This result illustrates that infectivity is affected by host barriers, not merely by exposure.
- The figure describes a specific experiment—not a universal numerical rule for other pathogens.

Some enteric pathogens survive luminal defenses

Norovirus

Resistance to acid, bile, and pancreatic enzymes; Lack of lipid envelope contributes.

Protozoa

Acid-resistant cyst forms promote survival during passage.

Helminths


Some infective eggs have protective outer coats.

Intestinal secretions resist colonization

Gastrointestinal protection is multilayered.

- Mucus and secretory immunoglobulin A contribute to mucosal defense.
- Defensins provide antimicrobial activity in the lumen and at epithelial surfaces.
- Bile and pancreatic enzymes create additional barriers to susceptible microbes.

Peristalsis restricts microbial overgrowth



- Normal peristalsis contributes to the clearance of ingested organisms.
- Obstruction or ileus can disrupt this protection and favor microbial overgrowth.

Enteric infection depends on barrier failure

Pathogen strategy


Resists gastric acid or enzymes; Attaches to intestinal epithelium; Penetrates mucosa in some infections.

Host susceptibility

Reduced gastric acidity, impaired motility, or disrupted microbiota lowers local resistance.

A shared fecal–oral exposure can lead to different outcomes in different hosts.

A fecal route may include a different entry portal



- The source notes that certain helminths shed eggs which develop into skin-penetrating larvae.
- Distinguish the route by which material leaves the host from the route of entry into the next host.

Sexual transmission requires intimate contact

Mucosal or intimate contact permits exchange of infective organisms.

- This route applies to multiple pathogen classes, not only bacteria or viruses.
- The genital tract is an important—but not exclusive—site involved in sexually acquired infections.
- Transmission classification describes exposure rather than a single tissue response.

Sexually transmitted viral pathogens

HSV

Herpes simplex virus

HIV

Human immunodeficiency
virus

HPV

Human papillomavirus

All three as examples transmitted during sexual contact.

Sexually transmitted bacterial pathogens

Treponema pallidum

Bacterial agent of syphilis; Sexually transmitted by intimate contact.

Neisseria gonorrhoeae

A gonococcal bacterium with genital or other mucosal exposure.

Other sexually transmitted organisms

Protozoa

Trichomonas vaginalis


Ectoparasites

Phthirus pubis (pubic lice)

General implication

Sexually transmitted infections include organisms beyond viruses and bacteria.

Genitourinary barriers influence infection



- Urine flow tends to remove microorganisms from the urinary tract.
- The source describes lactobacillus-associated vaginal acidity as a protective factor.

Sexual acquisition: Route is not pathology

Pathogen class	Named Robbins example	Interpretive point
Virus	HSV, HIV, HPV	Do not infer a single disease course from shared transmission
Bacterium	T. pallidum, N. gonorrhoeae	Clinical targets and injury differ
Protozoan	T. vaginalis	Sexual contact can transmit eukaryotic pathogens
Arthropod	Phthirus pubis	Intimate contact may spread ectoparasites

Vertical transmission has three time windows

Mother → fetus or newborn

```
graph TD; A[Mother → fetus or newborn] --> B[Before birth]; A --> C[During birth]; A --> D[After birth];
```

Before birth

Across placenta into fetal tissues


During birth

Exposure in the birth canal

After birth

Some pathogens transmitted in
milk

Placental transmission reaches the fetus




- Maternal infection during pregnancy for transplacental spread.
- Timing of fetal exposure contributes to the nature and extent of injury.

Gestational age changes effects of rubella

The effect of fetal infection depends partly on developmental timing.

- Rubella is the Robbins example of an infection interfering with fetal development.
- The degree and type of injury vary with fetal age at infection.
- A teratogenic outcome cannot be inferred from the route alone without considering timing.

Transmission during delivery



- This intrapartum process differs from transplacental fetal infection.

Neonatal conjunctivitis illustrates intrapartum spread

Gonococcal infection

Neisseria gonorrhoeae is a named cause of birth-associated conjunctivitis.

Chlamydial infection

Chlamydial conjunctivitis is the second textbook example.

Inference

Clinical presentation shortly after birth may reflect delivery exposure, not placental transfer.

Postnatal transmission in maternal milk


The source describes milk as a potential vehicle after delivery.

- Cytomegalovirus, HIV, and hepatitis B virus as examples.
- This statement addresses a possible route; It is not a patient-specific estimate of risk.
- Decisions about feeding or preventive interventions require guidance beyond this excerpt.

Vertical routes are mechanistically different

Time	Transmission event	Named illustration
Prenatal	Maternal infection → placenta → fetus	Rubella
Intrapartum	Exposure while passing through birth canal	Gonococcal / chlamydial conjunctivitis
Postnatal	Infectious exposure from maternal milk	CMV, HIV, HBV named in source

A neonatal case requires a timing history




- Prenatal, intrapartum, and postnatal mechanisms should not be collapsed into one label.
- The same infant may encounter more than one potential transmission window.

Saliva can transfer oropharyngeal viruses



- Epstein–Barr virus is the explicit textbook example.

Arthropods transmit pathogens through blood meals



- Some protozoa are spread by arthropod blood meals.
- The vector is part of the transmission mechanism; It is not itself the portal of entry.

Zoonotic transmission begins with animals

Animal-associated source

```
graph TD; A[Animal-associated source] --> B[Direct contact]; A --> C[Animal products]; A --> D[Vector]; B --> B1[Including an animal bite]; C --> C1[Exposure through animal-derived material]; D --> D1[Arthropod-mediated transfer];
```

Direct contact

Including an animal bite


Animal products

Exposure through animal-derived material

Vector

Arthropod-mediated transfer

Animal bites can bypass the skin barrier



- The intact epidermis usually acts as a physical barrier to microbes.
- Bite-associated penetration circumvents that barrier.
- Rabies among bite-related examples in the entry table.

Direct and indirect contact are route categories

Direct contact

Exposure occurs through contact with an infected person or an animal source.

Indirect contact

Exposure occurs through an intermediary material or environment before host entry.

The chapter key concepts explicitly include both contact categories.

Acquisition source and entry site may differ


Example	Acquisition pathway	Entry portal
Inhaled respiratory pathogen	Infectious particles in air	Respiratory epithelium / alveoli
Enteric pathogen	Ingested contaminated material	Gastrointestinal mucosa
Bite-associated infection	Infected animal or arthropod	Injured skin
Skin-penetrating larvae	Larvae develop after fecal shedding	Skin

The microbiome is a resident microbial community


Humans coexist with bacteria, fungi, viruses, and protozoa in and on the body.

- Distinct microbial ecosystems occur on skin and at mucosal surfaces.
- The microbiome has effects on host metabolism and immune function.
- Resident microbes are not automatically pathogens; Context determines their effects.

Microbiome niches are anatomically distinct




Resident microbes also help defend portals



- The gut microbiota competitively limits overgrowth of potential pathogens.
- Vaginal lactobacilli generate a low-pH environment that suppresses some pathogens.

Antibiotics can disrupt gut colonization resistance



- Antibiotic exposure and loss of normal intestinal microbiota as a susceptibility mechanism.
- This mechanism illustrates how a host defense alters the outcome of pathogen exposure.

Dysbiosis: Association does not prove causation

Association

Microbial community changes are reported with multiple disease states.

Unresolved causality

It is difficult to establish whether dysbiosis causes disease or follows it.

The source emphasizes unresolved direction of causation in many microbiome–disease associations.

Microbiome effects may extend beyond one organ

Nutrition

Communities synthesize vitamins and amino acids needed by the host.

Immune modulation

Microbes influence immune development and responses.

Distant interactions


Gut–lung, gut–skin, and gut–brain axes are proposed in the source.

Microbiome therapies require evidence by indication

A biologically plausible microbial intervention is not automatically clinically effective.

- The source supports fecal microbiota transplantation for recurrent *C. Difficile* infection.
- Other chronic disease applications have yielded mixed results.
- Microbiome-targeted therapies should not be generalized across diagnoses.

One exposure framework unifies all routes



- Identify the source, exit material, transfer mechanism, and portal separately.
- Consider microbial virulence and impaired host defenses before predicting infection.

Clinical vignette: Inhalational exposure

Clinical / epidemiologic vignette

A patient with pulmonary tuberculosis shares air in an enclosed setting with other people. The discussion concerns how organisms may reach another host.

Interpretation

- Source: Infected respiratory tract
- Exit: Infectious airborne particles
- Entry: Inhalation into respiratory tract
- Do not substitute an arbitrary fixed distance for assessment of dispersal.

Clinical vignette: Contaminated water

Clinical / epidemiologic vignette

Several people develop enteric illness after consuming water contaminated with human stool. Which transfer pathway best describes the exposure?

Interpretation

- Route: Fecal–oral
- Exit: Stool
- Vehicle: Contaminated drinking water
- Entry: Ingestion followed by GI barriers.

Clinical vignette: Neonatal conjunctivitis

Clinical / epidemiologic vignette

A newborn develops conjunctivitis after delivery to a mother with a genital infection. The task is to classify the probable window of acquisition.

Interpretation

- Consider transmission during birth.
- Relevant examples: Gonococcal or chlamydial infection.
- Do not automatically infer placental infection.

Clinical vignette: Antimicrobial disturbance

Clinical / epidemiologic vignette

A patient develops intestinal overgrowth by an opportunistic organism after broad-spectrum antimicrobial therapy. What protective mechanism may have been lost?

Interpretation

- Normal intestinal colonization resistance
- Loss of microbial competition
- C. Difficile is the source example
- Distinguish host susceptibility from the initial exposure route.

Compare transmission routes by release material

Mode	Exit material / source	Exposure mechanism
Respiratory	Respiratory particles	Inhalation
Fecal–oral	Stool-contaminated food or water	Ingestion
Sexual	Organisms present at sites of intimate contact	Mucosal / intimate contact
Vertical	Maternal organism before, during, or after birth	Placenta, delivery, or milk
Other	Saliva, animal source, vector	Contact or vector-mediated transfer

Synthesis: Predict spread from the route

Transmission is a sequence of biologically distinct requirements.

- Recognize when and how the infected source sheds a pathogen.
- Match the vehicle or contact mode to an entry portal in the new host.
- Assess epithelial barriers, mucosal defenses, and microbial survival strategies.
- Use maternal timing, animal exposure, vectors, and asymptomatic shedding when appropriate.

Route-based reasoning: final checks

Identify exposure mechanism before inferring disease or intervention.

- Which host or animal source releases the agent—and in what material?
- Does the next host encounter it by inhalation, ingestion, intimate contact, vertical transfer, or a vector?
- Which portal and local defenses must the pathogen negotiate?
- Does the clinical history distinguish asymptomatic shedding, gestational timing, and environmental transfer?