

Retinoblastoma

Molecular pathogenesis • Histomorphology • Clinicopathologic risk

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com


Learning objectives

- Explain heritable and nonheritable RB1 inactivation using the two-hit model.
- Recognize defining gross and microscopic patterns of retinoblastoma.
- Relate optic nerve, choroidal and extraocular spread to prognosis.
- Connect pathology with early detection, treatment and genetic surveillance.

Retinoblastoma: The essential definition

- Malignant retinal neoplasm arising in early childhood.
- Most common primary intraocular malignancy of childhood.
- Linked chiefly to biallelic loss of RB1 tumor-suppressor function.
- A paradigm of hereditary cancer susceptibility and somatic tumor evolution.

The retina is the anatomic site of origin



Original schematic — not a histologic specimen

Retinoblastoma develops within the retina and can enlarge into the vitreous space or extend into adjacent ocular tissues.

Why this neoplasm is distinctive

- A developing retina creates a time-limited population of susceptible precursor cells.
- One germline pathogenic allele can predispose every retinal precursor to a second hit.
- Multiple independent retinal tumors may form in predisposed children.
- Early recognition can preserve life, ocular structures and useful vision.

Opening clinical case: An abnormal red reflex

A young child is brought for assessment after a parent notices a white pupil in several flash photographs. No family history of eye cancer is known.

DISCUSS

- What serious intraocular diagnosis must be excluded urgently?
- Why does a negative family history not settle hereditary risk?


Leukocoria is a clinical warning sign

- A white pupillary reflex can reflect a retinal mass or other ocular pathology.
- It warrants prompt dilated examination by an ophthalmologist.
- Retinoblastoma is a critical diagnosis, not the only cause of leukocoria.
- Delayed diagnosis can allow ocular and extraocular tumor progression.

Other presenting features

- Strabismus can signal impaired fixation from a retinal lesion.
- Reduced vision may be difficult to detect in a preverbal child.
- Advanced intraocular tumors may cause pain, redness or secondary glaucoma.
- Symptoms alone cannot establish the diagnosis or determine heritable status.

A clinically oriented diagnostic sequence



Retinoblastoma is generally diagnosed clinically; Diagnostic intraocular biopsy is avoided because of seeding risk.

What ocular examination and imaging contribute

Ophthalmic assessment

- Indirect ophthalmoscopy identifies retinal tumor sites and foci.
- Examination under anesthesia may be needed in young children.
- Ultrasonography can identify intraocular mass and calcification.


Imaging and risk

- MRI assesses optic nerve, orbital and intracranial extension.
- Imaging complements the ocular examination rather than replacing it.
- Avoid unnecessary ionizing radiation in heritable retinoblastoma.

Retinoblastoma is primarily a genetic-pathogenesis model


- RB1 is a tumor-suppressor gene at chromosome 13q14.
- Loss of both functional RB1 alleles removes an important proliferative brake.
- The initiating event may be constitutional or entirely somatic.
- A minority of tumors follow an RB1-intact, MYCN-amplified pathway.

Knudson's two-hit principle




Both hits occur in the same tumor precursor; Germline predisposition changes where the first hit resides.

Heritable retinoblastoma: Constitutional first hit



Nonheritable retinoblastoma: Two somatic hits



Heritable versus nonheritable disease

Feature	Heritable RB1-associated	Nonheritable RB1-associated
First hit	Germline / constitutional	Somatic in a retinal cell
Second hit	Somatic loss of remaining allele	Second somatic loss
Typical laterality	Often bilateral or multifocal	Usually unilateral and unifocal
Family implication	Relatives may be at risk	Usually no inherited predisposition

Germline mutation is not synonymous with family history

- Heritable retinoblastoma can arise from a de novo constitutional RB1 variant.
- Affected children may have no relatives with retinoblastoma.
- Bilateral or multifocal tumors strongly support constitutional susceptibility.
- Unilateral disease does not exclude constitutional or mosaic RB1 alteration.

Dominant predisposition, recessive cellular phenotype

Organism-level inheritance

- A pathogenic constitutional RB1 variant confers inherited cancer susceptibility.
- The predisposition can be transmitted as an autosomal dominant trait.
- Each retinal precursor begins one step closer to transformation.


Cell-level oncogenesis

- A single RB1 loss-of-function allele does not itself produce a tumor.
- The neoplastic phenotype generally requires inactivation of both alleles.
- This is tumor suppression behaving recessively at the cellular level.

Why a single RB1 hit is insufficient


- An intact RB1 allele can still produce functional retinoblastoma protein.
- The remaining checkpoint function restrains inappropriate G1-to-S entry.
- A subsequent event must disable the retained functional allele.
- The two-hit concept illustrates selection of biallelic tumor-suppressor loss.

Normal RB protein restrains S-phase entry



RB is a central checkpoint integrator; Its status reflects mitogenic and antiproliferative signals.

Cyclin–CDK complexes release the RB brake



Functional RB inhibition versus RB1 gene loss

Genetic inactivation

- Loss of both RB1 alleles removes the functional suppressor protein.
- In a retinal precursor this can initiate retinoblastoma.
- RB1 is the canonical tumor-suppressor prototype.

Pathway-level inactivation

- Other cancers may impair RB activity through cyclin or CDK dysregulation.
- RB1 may be structurally intact but functionally inactivated.
- The distinction matters when interpreting molecular mechanisms.

What is the susceptible cell?

- Retinoblastoma emerges from a developing retinal lineage.
- Modern genetic studies implicate a cone photoreceptor precursor context.
- The disease illustrates that molecular lesions interact with cellular lineage.
- RB1 expression elsewhere does not mean equivalent tumor susceptibility everywhere.

An important RB1-negative molecular subset

- A small minority of retinoblastomas lack identifiable RB1 inactivation.
- High-level MYCN amplification can drive an RB1-intact tumor.
- The older name NMYC refers to MYCN.
- This route should not be inferred solely from morphology.

Why hereditary tumors are frequently bilateral

Heritable predisposition

- All developing retinal precursors start with an altered RB1 allele.
- Separate second hits can create independent tumor foci.
- Bilateral or multifocal presentation can therefore occur.

Sporadic disease

- Two independent somatic events must meet in one retinal lineage.
- This is a rarer combination than one additional hit.
- The usual presentation is a single unilateral tumor.

Pathologic morphology does not reveal germline status

- Tumors from heritable and nonheritable cases have the same basic morphology.
- Rosettes, necrosis and calcification do not identify the inherited first hit.
- Laterality, family history and molecular testing establish genetic risk.
- Histology answers local tumor questions, not constitutional genotype alone.

Case: Two independent retinal masses

A toddler has separate tumors in both eyes. Ophthalmoscopy shows multiple retinal foci; The parents report no affected relatives.


DISCUSS

- Which RB1 susceptibility pattern is most likely?
- Should germline testing still be considered?

Macroscopic pathology: Intraretinal tumor expansion

- Retinoblastoma begins as a retinal neoplasm.
- The enlarging tumor may replace retinal tissue and form an intraocular mass.
- Tumor extension toward the vitreous and other structures influences local stage.
- The true macroscopic extent must be determined by direct pathologic examination.

Tumor growth changes the normal retinal compartment




Original schematic — not a histologic specimen

The lesion displaces normal retinal architecture; The schematic depicts anatomy, not clinical fundus imaging.

Undifferentiated tumor: Primitive neuroblastic cells

- Dense collections of small, round tumor cells are characteristic.
- Nuclei are deeply basophilic and hyperchromatic.
- Tumor cells may appear relatively uniform in poorly differentiated areas.
- Microscopic heterogeneity includes differentiated foci in some tumors.

Flexner–Wintersteiner rosettes: True lumina




- Tumor cells encircle a genuine lumen.
- Shows photoreceptor differentiation.
- Not required for a diagnosis.

Interpreting a true Flexner–Wintersteiner rosette

- Tumor cells are radially arranged around a central lumen.
- The architecture implies retinal / photoreceptor differentiation.
- Rosettes can be present among less differentiated cells.
- Their absence does not by itself exclude retinoblastoma.

Homer Wright pseudorosette: Fibrillary center



- Cell ring surrounds neuropil-like fibers.
- A pseudorosette, not a true lumen.
- Less specific than Flexner–Wintersteiner rosettes.

Editable schematic — not microscopy


Histologic rosettes and differentiated elements

Structure	Central feature	Interpretation
Flexner–Wintersteiner	True lumen	Retinal differentiation
Homer Wright	Fibrillary neuropil-like material	Neuroblastic differentiation
Fleurette	Flower-like cells; No true lumen	Greater photoreceptor differentiation

Fleurettes indicate photoreceptor differentiation

- Fleurettes form flower-like arrangements of differentiated tumor cells.
- They represent a maturation pattern within retinal tumor tissue.
- Fleurettes and Flexner–Wintersteiner rosettes reflect related differentiation.
- Differentiation alone is not a reliable prognostic classifier.

Viable tumor cuffs and ischemic necrosis



Mechanistic schematic — distance from blood supply is explanatory, not a measured scale

Why necrosis develops away from vessels

- Rapidly growing retinal tumor tissue depends on its vascular supply.
- Viable tumor remains closest to patent vessels.
- Relatively avascular regions develop ischemic cell death.
- The viable-to-necrotic gradient explains a frequent histologic pattern.

Dystrophic calcification in retinoblastoma

- Calcific foci commonly accompany necrotic tumor tissue.
- This is dystrophic calcification, not evidence of systemic hypercalcemia.
- Calcification is diagnostically helpful on histology and ocular imaging.
- Necrosis and mineralization should not be mistaken for a tumor grade.

Histologic recognition: Positive and negative evidence

Supportive findings

- Malignant small round cells arising in the retina.
- True rosettes or fleurettes may show differentiation.
- Perivascular viability and dystrophic calcification support the pattern.

Interpretive limits

- Rosettes are not invariably present.
- Degree of histologic differentiation does not by itself predict prognosis.
- The hereditary and sporadic forms are morphologically indistinguishable.

Retinocytoma / retinoma: A related lesion

- Retinoma (retinocytoma) is a relatively differentiated retinal proliferation.
- It is considered a possible precursor of retinoblastoma.
- An affected eye may contain retinoblastoma while the fellow eye has retinoma.
- This combination is particularly pertinent to heritable disease.

Retinoma versus retinoblastoma

Retinoma / retinocytoma

- A relatively differentiated retinal lesion with precursor potential.
- May be detected during hereditary retinoblastoma assessment.
- Still warrants directed ophthalmic surveillance.


Retinoblastoma

- Malignant proliferating retinal tumor.
- Includes primitive cells, necrosis and sometimes differentiated structures.
- Needs assessment of local invasion and overall clinical extent.

Histologic differentiation is not the main prognostic axis

- The textbook explicitly notes poor association between differentiation and prognosis.
- A well-formed rosette should not reassure against invasive spread.
- Extent of local invasion is more clinically consequential.
- Reporting must prioritize optic nerve, uveal and extraocular involvement.

Choroidal invasion connects microscopic spread to risk



Pathology: Depth and extent of invasion matter more than microscopic differentiation

Massive choroidal invasion: A reporting priority

- Choroidal invasion is an adverse feature when extensive.
- A consensus framework uses a maximum invasive focus dimension of at least 3 mm for “massive” invasion.
- The local pathology protocol must define orientation and measurement.
- Prognostic thresholds and treatment consequences remain protocol-dependent.

Optic nerve invasion must be anatomically localized



Determine anatomic level and margin status; Postlaminar spread is a high-risk feature

Postlaminar optic nerve invasion is high risk

- Invasion beyond the lamina cribrosa implies an advanced neural route.
- Microscopic level is more informative than simply recording invasion present.
- Tumor at an optic nerve transection margin is particularly consequential.
- Pathology reports must document the deepest invasion level.


Adverse features of retinoblastoma

Feature	Why clinically important
Extraocular extension	Tumor has escaped the confines of the globe
Postlaminar optic nerve invasion	Access to an important route for extraocular spread
Massive choroidal invasion	Substantial uveal invasion associated with metastasis risk
Optic nerve cut end involved	Concern for residual tumor along nerve pathway

There is no single universal high-risk feature set

- Definitions of high-risk pathology differ across retinoblastoma centers.
- A global survey found consensus for postlaminar nerve invasion, a positive nerve transection and extrascleral extension.
- Other criteria, including choroidal and anterior segment findings, varied by practice.
- Interpret a histologic feature in the context of the center's documented protocol.

Anatomic routes of spread



Brain and bone marrow dissemination; Lung involvement is relatively uncommon.

Metastatic behavior is anatomically patterned

- Spread to the brain is a major concern with advanced retinoblastoma.
- Bone marrow is another characteristic systemic site.
- Lung dissemination is less characteristic than in many other solid tumors.
- Risk rises with extraocular extension and invasive histologic findings.

Trilateral retinoblastoma

- Heritable retinoblastoma can be associated with an intracranial primitive neuroblastic malignancy.
- The classic association is pineal-region disease.
- This constellation is termed trilateral retinoblastoma.
- A particularly poor outcome for this association.

A second cancer risk beyond the eye

- Constitutional RB1 loss predisposes to later primary cancers.
- Osteosarcoma and soft-tissue sarcomas are characteristic examples.
- The risk is not limited to recurrent ocular disease.
- This motivates individualized survivorship counseling and exposure awareness.

Three treatment goals must be ordered correctly


First: Protect survival

- Eradicate a potentially fatal malignancy.
- Control intraocular or extraocular disease.
- Assess histologic and clinical metastatic risk.

Next: Preserve eye and vision

- Preserve the affected globe where oncologically safe.
- Maintain or recover useful vision where feasible.
- Protect the child from avoidable long-term morbidity.

Chemoreduction plus focal tumor consolidation



Selective intra-arterial ophthalmic chemotherapy

- Medication can be delivered selectively through the ophthalmic artery.
- This approach concentrates treatment at the eye in eligible cases.
- Its suitability depends on laterality, tumor distribution and specialist evaluation.
- Local control still requires longitudinal ocular assessment.

When enucleation is clinically important

- A severely involved eye may be unsafe or impossible to salvage.
- Enucleation can remove the tumor while allowing complete pathologic staging.
- Optic nerve length, cut margin and uveal invasion deserve meticulous evaluation.
- The decision balances systemic safety against ocular preservation.

Enucleation pathology: Necessary observations

- Document the tumor distribution, size and associated retinal changes.
- Sample tumor in relation to choroid, sclera and optic nerve head.
- State the deepest optic nerve infiltration and optic nerve margin status.
- Describe any extraocular tissue extension and recognized adverse findings.

Treatment choice is not determined by rosette abundance

What histology tells us

- Well-differentiated histology does not guarantee a low-risk tumor.
- High-risk invasion findings may change postoperative management.
- Need to distinguish histologic structure from disease extent.


What clinical staging adds

- Tumor size, laterality, foci and eye condition influence local treatment.
- Optic nerve or orbital spread changes systemic evaluation.
- Vision potential affects globe-sparing decisions when oncologically safe.

Genetic evaluation guides family-level prevention

- Identify whether a constitutional RB1 pathogenic variant is present.
- Offer variant-specific assessment of at-risk relatives when feasible.
- Germline negative blood testing does not always exclude low-level mosaicism.
- Family counseling must distinguish a heritable predisposition from tumor-only mutations.

Genetic testing: A practical interpretation pathway



Gene testing is complementary to urgent clinical treatment, not a reason to postpone it.

Surveillance in germline RB1 carriers

- The developing retina is examined repeatedly during early childhood.
- Some examinations require anesthesia until reliable cooperation is possible.
- Frequency and duration are individualized by genetic and clinical risk.
- Specialist schedules are more reliable than a one-size-fits-all interval.

An eye saved is not necessarily a child discharged

- Local control does not abolish the constitutional cancer predisposition.
- The fellow eye can develop additional lesions when RB1 is heritable.
- Long-term follow-up includes functional vision and subsequent cancer awareness.
- Surveillance is matched to genotype, disease status and treatment history.

Clinicopathologic case: Optic nerve extension

An enucleated globe contains retinoblastoma with tumor extending beyond the lamina cribrosa into the optic nerve; The cut margin is negative.

DISCUSS

- Which histologic feature is adverse?
- What additional anatomic sites and report elements should be reviewed?

Morphology case: Differentiated islands

Microscopy of a retinal tumor shows sheets of hyperchromatic small round cells, numerous necrotic foci, and several true rosettes surrounding lumina.

DISCUSS

- Which differentiated structure is illustrated by the rosettes?
- Can rosette frequency establish a favorable prognosis?

Genetic counseling case: A unilateral tumor

A child has a single unilateral retinal tumor and no affected family members. An RB1 pathogenic variant is detected in peripheral blood.

DISCUSS

- Does unilateral disease exclude a constitutional predisposition?
- What does this finding imply for follow-up and relatives?

High-yield reasoning: Phenotype to inference

Observation	What it supports	What it does not prove
Bilateral multifocal tumors	Constitutional predisposition	A known positive family history
True retinal rosettes	Photoreceptor differentiation	Favorable clinical outcome
Necrosis with calcification	Typical tumor architecture	A specific genetic subclass
Postlaminar nerve invasion	High-risk invasive spread	Established distant metastasis

Retinoblastoma: Five integrative conclusions

- Retinal neoplasia generally follows biallelic RB1 tumor-suppressor loss.
- A germline first hit predisposes to bilateral, multifocal and second primary tumors.
- Small blue cells, true rosettes, perivascular necrosis and calcification define the morphologic framework.
- Differentiation is not a stand-alone prognostic marker.
- Optic nerve, choroid and extraocular spread drive adverse pathologic risk.

Further reading

- Robbins, Cotran & Kumar: Pathologic Basis of Disease. 11th edition. Chapters 7 and 29.
- GeneReviews: Retinoblastoma (NCBI Bookshelf NBK1452).
- International Retinoblastoma Staging Working Group: pathology consensus (2009).
- Retinoblastoma high-risk pathology: multicenter practice survey (2021).