

Rheumatoid Arthritis

Autoimmune mechanisms · synovial morphology · clinicopathologic correlation

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraisakhri.com

The central clinicopathologic problem

- RA is a chronic autoimmune disorder in which proliferative synovitis drives joint damage.
- An inflamed synovial membrane expands into a destructive pannus.
- Cartilage loss, periarticular bone erosion and sometimes ankylosis follow.
- Extraarticular involvement confirms that RA is a systemic immune disease.

Learning outcomes

- Connect loss of self-tolerance to T-cell, B-cell and macrophage responses.
- Explain how cytokines, proteases and osteoclasts produce specific joint lesions.
- Recognize the defining synovial and rheumatoid-nodule morphology.
- Interpret joint distribution, serology, radiography and principal differentials.

RA and primary osteoarthritis: Distinct initiating processes

Rheumatoid arthritis

- Primary autoimmune synovitis
- Inflammatory pannus is destructive
- Systemic manifestations may occur

Osteoarthritis

- Primary cartilage injury and degeneration
- Inflammation may exacerbate damage
- Usually confined to joints

Why the synovium matters



Schematic — not a specimen

Intact joint

Uninflamed synovial lining

A normal joint has a smooth articular surface and a thin synovial lining.

Nonsuppurative proliferative synovitis

- “Synovitis” identifies synovial membrane inflammation as the principal lesion.
- “Proliferative” denotes synoviocyte expansion and thickened synovial projections.
- “Nonsuppurative” distinguishes chronic immune synovitis from purulent bacterial arthritis.
- The lesion ultimately extends beyond synovium into cartilage and adjacent bone.


Population pattern

- US prevalence is approximately 0.25%–1% in the supplied edition.
- Incidence peaks during the third and fourth decades of life.
- Females are affected approximately three times as often as males.
- These are textbook estimates, not current population surveillance statistics.

Why RA is classified as an autoimmune disease

- T lymphocytes react to self-antigens associated with joint tissues.
- Many patients produce autoantibodies to modified self-proteins and IgG.
- Susceptibility alleles and environmental candidates are implicated.
- Immune-directed treatment reduces inflammation and joint injury.

The disease spans joints and extraarticular tissues




Schematic

The joint lesion is the dominant expression of systemic autoimmunity.

The initiating joint antigen is not established

- CD4+ helper T-cell reactivity is central to the pathogenic model.
- Candidate targets include collagen and chemically modified peptides.
- Multiple joint antigens have been proposed; No single initiating antigen is proven.
- This uncertainty matters when distinguishing a plausible mechanism from a demonstrated cause.

Predisposition to chronic autoimmune arthritis



Schematic

The proposed sequence is multifactorial; Triggers remain incompletely defined.

Genetic contribution: Risk, not destiny

- That inherited susceptibility accounts for about half of RA risk.
- This estimate describes population-level genetic contribution, not individual prediction.
- Genetic predisposition influences development, persistence and progression.
- Environmental exposures and immune regulation remain necessary considerations.


HLA-DR4 and ACPA-positive disease

- The HLA-DR4 allele is associated with the ACPA-positive subset of RA.
- Class II HLA molecules present peptide antigens to CD4+ T cells.
- Proposed molecular mimicry involving some microbial epitopes.
- An HLA association supports susceptibility but is not a stand-alone diagnostic test.

Environmental factors: Proposed triggers

- Smoking may promote modification of self-proteins through citrullination.
- Infection, including periodontitis, is discussed as a possible contributor.
- The source explicitly states that environmental determinants are not clearly defined.
- Do not assume that a proposed association identifies a proven causal exposure.

Citrullination can alter immune recognition




Schematic

Citrullination is established; Its precise initiating role in any given case is uncertain.

Joint proteins implicated as ACPA targets

- Citrullinated fibrinogen and type II collagen are among described targets.
- Modified α -enolase and vimentin are additional candidate antigens.
- A broad antigen repertoire is consistent with multiple self-reactivities.
- A target antigen is not automatically the initiating cause of the disease.

CD4+ T-cell activation initiates amplification



Schematic

T-cell-driven inflammation links failed tolerance to destructive synovitis.


Th1 cells: The macrophage-activation arm

- Th1 helper cells release interferon- γ (IFN- γ).
- IFN- γ activates macrophages and resident synovial cells.
- Activated cells participate in chronic inflammation and tissue injury.
- The Th1 effect is complementary to other helper-T-cell pathways.

Th17 cells recruit additional inflammatory cells

- Th17 cells release interleukin-17 (IL-17).
- IL-17 recruits neutrophils and monocytes into inflammatory sites.
- Incoming cells help sustain the inflamed synovial environment.
- Cell recruitment converts antigen-specific activation into widespread joint inflammation.

Macrophages amplify destructive inflammation



Schematic

Macrophage-derived cytokines link immune activation to matrix destruction.


TNF is a major inflammatory mediator

- TNF contributes to leukocyte recruitment and persistent synovial inflammation.
- It helps induce proteolytic pathways that damage articular cartilage.
- The therapeutic benefit of TNF antagonists supports its pathogenic importance.
- TNF is an important mediator, not the sole explanation for RA.

IL-1 and IL-6 reinforce synovitis

- Macrophage-derived IL-1 and IL-6 act alongside TNF.
- These cytokines support inflammatory-cell recruitment.
- They stimulate synovial cells toward cartilage-degrading protease production.
- Parallel cytokine activities help sustain the destructive inflammatory milieu.


Cartilage destruction is cytokine-linked



Schematic

An immune lesion causes mechanical failure by degrading cartilage.

RANKL provides a mechanism for bone erosion




Schematic

Cartilage and bone destruction involve distinct cellular effectors.

B-cell responses are part of RA synovitis

- The inflamed synovium often contains B cells and abundant plasma cells.
- Lymphoid follicles with germinal centers may appear locally.
- Some plasma cells secrete antibodies recognizing self-antigens.
- Autoantibodies can also be produced in lymphoid organs outside the joint.

Where autoantibodies fit in the lesion



Schematic

Antibody-mediated mechanisms are plausible contributors, not the only known cause.

ACPA: Anti–citrullinated peptide antibodies

- ACPAs recognize epitopes created by citrullination of proteins.
- The supplied Robbins text describes detection in up to 70% of RA patients.
- They are diagnostically useful and associated with an important immune phenotype.
- The source suggests—but does not establish completely—a contribution to severity and persistence.

Rheumatoid factor targets immunoglobulin G

- Rheumatoid factors bind the Fc region of IgG.
- The source describes IgM and IgA rheumatoid-factor autoantibodies.
- Rheumatoid factors are found in approximately 80% of persons with RA.
- They may occur in people without RA and are absent in some affected patients.


ACPA and rheumatoid factor answer different questions

Feature	ACPA	Rheumatoid factor
Antigen	Citrullinated self-peptides	Fc region of IgG
Main value	Supports RA diagnosis	Supports RA diagnosis
In supplied source	Up to 70% of patients	About 80% of patients
Limitation	Not universal	May also occur without RA

Immune complexes: Contributory rather than exhaustive

- Rheumatoid factor can participate in immune-complex formation.
- Immune complexes may be deposited within joints.
- T-cell cytokines, macrophages and osteoclasts remain central in the mechanism.
- RA cannot be reduced to a purely immune-complex arthritis.

The destructive RA network




Schematic

Several linked effector pathways converge on joint destruction.

Separate demonstrated events from hypotheses

- Established in the source: Synovial inflammation, autoantibodies, pannus and erosions.
- Well-supported mediators: TNF, IL-1, IL-6, IL-17, IFN- γ and RANKL.
- Less certain: The initial antigen and exact environmental trigger.
- The balance of T-cell and antibody contributions is not identical in every patient.

Synovial transformation in early RA



Schematic — not a specimen

Early RA

Thickened, villous synovium

Synovial hypertrophy, inflammation and angiogenesis precede tissue destruction.

Synovium changes from smooth to villous

- Inflamed synovium becomes thickened, edematous and hyperplastic.
- Its previously smooth contour develops delicate and bulbous villi.
- Villi expand the surface of inflammatory synovial tissue.
- Villous hyperplasia is one characteristic feature of chronic RA synovitis.

Microscopic signature: Synoviocyte proliferation

- Synovial lining cells proliferate in response to sustained inflammation.
- The expanded lining is part of the synovial hypertrophy seen grossly.
- The pathologic lining sits adjacent to inflammatory exudate and cartilage.
- It participates in the formation of a growing pannus.

Microscopic signature: Inflammatory cell mixture

- CD4+ T lymphocytes are prominent in the synovial inflammatory infiltrate.
- B lymphocytes, plasma cells, dendritic cells and macrophages are also present.
- The infiltrate explains ongoing local cytokine and antibody production.
- The composition supports immune-mediated rather than purely degenerative disease.

Lymphoid follicles and local immune organization

- B-cell aggregates can form recognizable follicles within inflamed synovium.
- Secondary follicles may contain germinal centers.
- Plasma-cell accumulation accompanies chronic antigen-driven responses.
- These findings help explain local autoantibody production.

Synovial angiogenesis sustains inflammation

- Vascularity increases because new vessels form within diseased synovium.
- Angiogenesis supports an expanded mass of inflammatory tissue.
- Blood vessels provide routes for continued leukocyte recruitment.
- The result is a metabolically active, enlarging synovial lesion.

Fibrin and neutrophils at the articular surface

- A fibrin-rich exudate coats synovial and articular surfaces.
- Neutrophils are found within the exudate.
- This surface exudate coexists with a predominantly chronic cellular infiltrate.
- Neutrophils here do not make RA equivalent to septic suppurative arthritis.


Osteoclastic activity marks the bone interface

- Osteoclast activity increases in bone neighboring inflamed synovium.
- Activated synovial tissue can invade periarticular bone.
- This produces marginal erosions and subchondral cysts.
- Bone erosion is a structural consequence of inflammatory osteoclast stimulation.

Five pathologic components of RA synovitis

Component	Morphologic change	Clinicopathologic significance
Lining cells	Hyperplasia and villi	Expanding synovial mass
Leukocytes	T, B, plasma, myeloid cells	Immune activity
Microvessels	Angiogenesis	Inflammatory supply
Articular surface	Fibrin-rich neutrophilic exudate	Active surface inflammation
Bone interface	Osteoclasts and erosions	Structural joint injury

Pannus: Inflammatory tissue becomes invasive



Schematic — not a specimen

Progressive RA


Pannus invading articular tissues

Inflammatory pannus expands over cartilage and promotes erosion at joint margins.

What constitutes a pannus?

- Edematous hyperplastic synovium forms its scaffold.
- Inflammatory cells, fibroblasts and granulation tissue expand the lesion.
- The pannus extends over articular cartilage.
- It promotes erosion of cartilage and ultimately altered joint architecture.

Pannus converts inflammation to joint destruction




Schematic

Pannus is not simply a synonym for synovial swelling.

Cartilage erosion and bone erosion are distinct

- Synovial proteases degrade articular cartilage matrix.
- Activated osteoclasts resorb periarticular and subchondral bone.
- The pathways intersect but are not anatomically identical.
- Both contribute to reduced joint space, deformity and limited movement.

How inflammatory fusion can develop



Schematic

Ankylosis is a late potential outcome, not a necessary feature of every case.

Fibrous and bony ankylosis are sequential possibilities

Fibrous ankylosis

- Fibrous tissue bridges a joint
- Mechanical movement becomes restricted
- May precede ossification


Bony ankylosis

- Ossified bridge fuses opposing bones
- Represents structural bone fusion
- Late destructive joint consequence

Rheumatoid nodules: Extraarticular morphology

- They are described as an infrequent manifestation of RA.
- Typical sites include subcutaneous forearm and elbow regions.
- Occipital and lumbosacral sites are also described.
- The nodules are usually small, firm, oval and nontender.

Rheumatoid nodule: A layered necrobiotic lesion



Palisaded activated macrophages

Lymphocytes and plasma cells
around the rim

Schematic — not a histologic specimen

Systemic vasculitis in severe RA

- Severe disease may be accompanied by leukocytoclastic vasculitis.
- An acute necrotizing process affecting small and large arteries.
- Pleural, pericardial and pulmonary tissues may be involved.
- Some vascular and serosal lesions later evolve toward fibrosis.

Extraarticular pulmonary and cardiac disease

- RA-associated pathology may extend into lungs and pleura.
- Cardiac and pericardial involvement are also recognized.
- Vascular lesions can overlap with other systemic autoimmune phenotypes.
- A joint-predominant presentation does not exclude important visceral injury.

Ocular manifestations fit systemic autoimmunity

- Uveitis can occur in the setting of RA.
- Keratoconjunctivitis and sicca-like changes may accompany disease.
- The textbook explicitly cross-references Sjögren syndrome.
- Extraarticular symptoms merit evaluation beyond the joint examination.

Systemic symptoms can precede arthritis

- The source describes malaise, fatigue and generalized musculoskeletal pain.
- In about half of patients, these symptoms precede joint involvement.
- Recognizable arthritis may appear over subsequent weeks or months.
- The prodrome reflects the systemic nature of the inflammatory process.

RA typically presents as symmetric polyarthritis

- Small joints of the hands and feet are characteristically involved early.
- Joint involvement is usually approximately symmetric.
- Larger joints may be affected as disease progresses.
- The pattern is a strong clinical clue, not a substitute for full evaluation.

Joint distribution: Hands and feet first

- Hand and foot symptoms are generally prominent early.
- Wrists and ankles are subsequently common sites.
- Elbows and knees may be involved with wider disease.
- Small-joint predominance helps differentiate RA from primary OA.

Hand-joint distribution guides differential diagnosis

Typical RA distribution

- Metacarpophalangeal joints
- Proximal interphalangeal joints
- Symmetric small-joint inflammation

Primary osteoarthritis emphasis

- Degenerative articular cartilage pathology
- Different hand-joint pattern
- Often weight-bearing joints elsewhere

Inflammatory pain follows inactivity

- RA joints are characteristically swollen, warm and painful.
- Pain commonly worsens in the morning or after inactivity.
- Inflammatory stiffness differs from the typical mechanical OA pattern.
- Link this timing to active synovitis rather than cartilage wear alone.

Clinical swelling reflects synovial disease

- Synovial hypertrophy increases the volume of periarticular tissue.
- Inflammatory exudation contributes to effusions.
- Warmth and tenderness track active inflammatory changes.
- Progressive structural damage restricts range of motion.

Natural history is heterogeneous

- Most cases follow a chronic course with episodes of increased and decreased activity.
- Continued activity can cause joint enlargement and declining motion.
- Some patients, particularly seronegative ones, stabilize or regress.
- Robbins also describes a small group with abrupt, severe polyarticular onset.

Periarticular inflammation produces deformity

- Tendons, ligaments and adjacent soft tissues can also become inflamed.
- Progressive soft-tissue changes affect joint alignment and function.
- Ulnar deviation and flexion–hyperextension deformities may develop.
- The characteristic appearance reflects both synovial destruction and periarticular injury.


Interpreting hand deformities carefully

- Ulnar deviation describes deviation of fingers toward the ulnar side.
- Flexion–hyperextension deformity describes abnormal postures across finger joints.
- These are distinct descriptors, not interchangeable synonyms.
- Clinical configuration must be described explicitly rather than inferred from a label.

Radiographic signs of structural RA

- Joint effusions may accompany active synovial inflammation.
- Juxtaarticular osteopenia is a characteristic associated finding.
- Periarticular erosions reflect osteoclast-mediated bone destruction.
- Joint-space narrowing reflects cartilage loss.

Clinical diagnosis integrates three evidence types



Schematic

No single test replaces the overall clinicopathologic assessment.

ACPA is useful but not universal

- A positive ACPA test helps support RA in an appropriate clinical setting.
- The source states that detection is possible in up to 70% of patients.
- A negative ACPA assay cannot, by itself, exclude RA.
- Interpret serology together with joint pattern and radiographic changes.

Rheumatoid factor requires contextual interpretation

- Rheumatoid factor is common in RA but not specific to it.
- Some people without RA may have detectable rheumatoid factor.
- Some patients with definite RA have no rheumatoid factor.
- The diagnostic question is whether laboratory results fit the inflammatory joint phenotype.

Seronegative RA is a recognized diagnosis

- That about 20% of patients are seronegative.
- In seronegative patients, diagnosis relies on clinical and radiographic features.
- Negative autoantibody tests do not erase synovitis and erosive morphology.
- Tissue biopsy is rarely required in the routine diagnostic approach.

Why routine synovial biopsy is seldom necessary

- A characteristic clinical distribution often directs the diagnostic workup.
- Serology and radiographic evidence are supportive.
- Chronic synovial inflammation itself is not unique to RA.
- Biopsy is therefore rarely needed for typical cases, according to the source.

Rheumatoid arthritis versus osteoarthritis

Feature	Rheumatoid arthritis	Osteoarthritis
Pathogenic driver	Autoimmune synovitis	Mechanical cartilage injury
Inflammation	Primary, persistent	May be secondary
Joint predilection	Small joints, polyarticular	Weight-bearing joints
Morphology	Pannus, erosions; possible fusion	Cartilage damage, osteophytes, cysts
Serology / organs	ACPA, RF; extraarticular lesions	No characteristic autoantibodies or systemic disease

RA versus spondyloarthritis: Tissue compartment

Rheumatoid arthritis

- Synovial inflammation is dominant
- Symmetric small-joint disease is typical
- ACPA or RF may be present

Spondyloarthropathies

- Inflammation frequently targets entheses
- Sacroiliac and vertebral involvement
- HLA-B27 association; Often RF absent

RA versus septic arthritis: Urgency and mechanism

Rheumatoid arthritis

- Chronic autoimmune synovitis
- Often symmetric polyarticular pattern
- Nonsuppurative inflammatory lesion


Acute bacterial arthritis

- Direct microbial infection of a joint
- Typically abrupt painful single joint
- Aspiration and organism identification are important

Therapy is directed at immunologic injury

- Treatment seeks to reduce pain and inflammation.
- An equally important goal is to slow or arrest irreversible joint destruction.
- Corticosteroids and methotrexate among treatment options.
- Mechanism-directed biologic approaches target TNF or lymphocyte responses.

Why blocking TNF can improve RA



Schematic

Therapeutic success is evidence of TNF involvement, not exclusive causation.

Anti-TNF therapy introduces infection risk

- TNF antagonists suppress an important arm of host inflammatory defense.
- The source specifically highlights increased risk of opportunistic infection.
- *Mycobacterium tuberculosis* is a named example.
- This is a therapeutic risk-benefit implication, not a reason to mislabel RA as infection.

Application case: Inflammatory small-joint disease

Hypothetical patient: Symmetric pain and swelling in metacarpophalangeal and proximal interphalangeal joints, with prominent stiffness on waking. ACPA is detected.

Which immune and morphologic mechanisms best explain this pattern?

Case interpretation: Link phenotype and tissue injury

- Symmetric small-joint inflammation favors an RA-type clinicopathologic pattern.
- ACPA supports autoimmunity involving citrullinated antigens.
- CD4+ responses, macrophage cytokines and hyperplastic synovium explain active synovitis.
- Pannus, proteases and osteoclasts explain later cartilage and bone destruction.

Application case: Antibodies are absent


Hypothetical patient: Chronic symmetric inflammatory arthritis of the hands and wrists; ACPA and rheumatoid factor are both negative. Radiographs show periarticular erosions and joint-space narrowing.

Does negative serology exclude rheumatoid arthritis?

Case interpretation: Seronegative does not mean noninflammatory

- The absence of ACPA and rheumatoid factor does not rule out RA.
- A compatible clinical pattern and structural radiographic findings retain diagnostic value.
- Other causes of inflammatory arthritis still require differential assessment.
- The source states that biopsy is seldom necessary for a typical clinical picture.

Mechanistic synthesis: One disease, linked compartments



Schematic

RA is an autoimmune synovitis whose tissue damage produces its clinical phenotype.

Eight findings to retain

- RA is a systemic autoimmune disease centered on chronic proliferative synovitis.
- CD4+ T cells, cytokine-activated macrophages and B cells participate.
- TNF, IL-1, IL-6, IL-17, IFN- γ and RANKL connect inflammation to tissue damage.
- Pannus destroys cartilage; Osteoclasts cause erosions; Ankylosis may develop.
- ACPA and RF assist diagnosis; Neither is universally positive.

Primary Robbins references

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. 2026.
- “Rheumatoid Arthritis” within the Diseases of the Immune System section.
- “Rheumatoid Arthritis” within the Bones, Joints, and Soft Tissue Tumors section.
- “Comparative Features of Osteoarthritis and Rheumatoid Arthritis.”