

Role of Infections and Other Environmental Factors

Infectious stimuli

- APC costimulation
- Molecular mimicry

Environmental exposures

- Microbiome and tissue damage
- Ultraviolet radiation; smoking

Immune balance

- Potential protection by infection
- Sex-related susceptibility

Dr Abu Suraih Sakhri

MBBS, MD (Pediatrics)


www.abusuraihsakhri.com

Autoimmunity emerges from interactions between immune tolerance, susceptibility, and environmental context.

Learning objectives

- Explain infection-induced APC costimulation and loss of T-cell anergy.
- Distinguish molecular mimicry from nonspecific microbial B-cell activation.
- Trace tissue injury, ultraviolet radiation, and smoking to proposed autoimmune mechanisms.
- Evaluate microbiome, protective infection, and sex effects without overstating causality.
- Integrate environmental triggers with genetic susceptibility and immune tolerance.

Autoimmunity requires failure of self tolerance



- Environmental stimuli are enabling factors, not sufficient causes in every individual.
- The same self antigen can be innocuous or immunogenic depending on presentation context.

Why infection is an attractive trigger

Immune activation

- Microbes activate antigen-presenting cells
- Costimulatory capacity can increase

Antigenic overlap

- Microbial structures may resemble self
- Cross-reactivity becomes possible

Inflammatory damage

- Tissue injury exposes self constituents
- Cytokines recruit additional cells

Genetic susceptibility does not determine destiny

Inherited predisposition

- HLA alleles influence peptide display.
- Non-HLA variants alter immune regulation.
- Most carriers do not necessarily develop disease.

Environmental contribution

- Infections can activate APCs.
- Tissue damage can change antigen access.
- Exposure interacts with an already susceptible host.

Genetic association and an environmental exposure are not individually equivalent to causation.

Central tolerance and peripheral tolerance


Central tolerance

- Immature autoreactive lymphocytes are removed.
- B cells may undergo receptor editing.
- Some regulatory T cells develop centrally.

Peripheral tolerance

- Anergy silences antigen recognition without activation.
- Regulatory T cells suppress responses.
- Inhibitory pathways and apoptosis restrain self-reactive cells.

The two-signal requirement for T-cell activation



- Antigen recognition without sufficient costimulation normally favors functional unresponsiveness.
- Infection can change the second-signal context of an otherwise familiar self antigen.

Anergy: Recognition without effective response

Resting APC context

- Self antigen may be presented.
- B7 expression may be inadequate.
- Autoreactive T cells remain functionally anergic.

Infection-activated context

- APCs increase costimulatory signals.
- Self-antigen presentation may continue.
- A previously unresponsive T cell can be activated.

The critical change is immune context, not necessarily the identity of the self peptide.

T-cell costimulation is a checkpoint

APC

- Displays peptide on MHC
- Provides B7 costimulation

T lymphocyte

- Recognizes peptide–MHC
- Receives B7 via CD28

Regulation

- Limits response to self
- Failure increases autoimmune potential

Two principal infection-linked mechanisms


Costimulation mechanism

- Microbe activates an APC.
- APC displays self antigen and increases B7.
- Anergy may break; Autoreactive T cell responds.

Molecular mimicry mechanism

- Microbial antigen resembles self antigen.
- Pathogen-directed lymphocytes may cross-react.
- Self tissue becomes a target.

Microbes can increase APC costimulation



- The source proposes costimulator induction as an explanation for infection-associated autoimmunity.
- Activation is favored when self antigens are simultaneously presented.

What changes on the antigen-presenting cell?


Before infection

- May display tissue-derived self peptides.
- Inadequate costimulation limits response.
- T-cell anergy can be preserved.

During infection

- APC activation increases costimulatory molecule expression.
- Self peptide can still be displayed.
- The same T-cell specificity may now respond.

B7–CD28 links innate signals to adaptive activation



- B7 on APCs and CD28 on T cells provide an illustrative second-signal interaction.
- This is a proposed pathway to loss of anergy, especially with impaired regulation.

Self-antigen presentation is the crucial coincidence

Self peptide available

- APC presents tissue antigen
- Autoreactive T cell can recognize it


APC activated

- Microbial stimulus increases costimulation
- Antigen encounter becomes activating

Regulation inadequate

- Protective brakes may be defective
- Autoimmune reaction becomes more likely

Regulatory defects amplify infection-triggered activation



- The textbook explicitly emphasizes that deficient regulation makes this mechanism more plausible.
- An infectious trigger need not directly damage the target tissue to alter APC function.

An antigen-specific response can be context dependent

Nonactivating encounter

- Self peptide recognized.
- Insufficient second signal.
- Anergy or other tolerance persists.

Activating encounter

- Same self peptide may be recognized.
- Infection changes APC costimulation.
- Autoreactive T-cell activation is possible.

Distinguish altered antigen presentation context from a newly created self antigen.

Costimulation model: Clinical interpretation


- A temporal association between infection and autoimmune symptoms is biologically plausible.
- The sequence is not proof that the infection caused disease in an individual patient.
- Host susceptibility and regulatory pathways modify the response.
- The pathogen need not share an antigen sequence with the affected organ.

How to identify the costimulation model

A microbial infection activates antigen-presenting cells; A self-reactive T cell recognizes a tissue peptide during this response.

- Which change provides the second T-cell signal? B7 expression on the APC.
- Which checkpoint may fail? Peripheral T-cell anergy.
- What remains uncertain? Whether this sequence causes a particular human autoimmune disorder.

Molecular mimicry: Antigen resemblance



- Shared amino acid sequences are one proposed basis of antigenic resemblance.
- The source describes molecular mimicry as an important proposed mechanism.

Mimicry differs from nonspecific activation


Molecular mimicry

- Antigenic similarity is central.
- Antimicrobial specificity overlaps a self target.
- Cross-reaction can involve cells or antibodies.

Bystander costimulation


- APC activation is central.
- Self antigen is displayed during infection.
- No shared microbial/self sequence is required.

The microbial immune response is the starting point




- The original stimulus is foreign, even when part of the resulting response recognizes self.
- Recognition of self is an unintended consequence of overlapping specificity.

Humoral mimicry: Cross-reactive antibodies



- The source uses rheumatic heart disease to demonstrate antibody cross-reactivity.
- An antibody to a pathogen does not automatically imply clinically important autoimmunity.

Cellular mimicry: Cross-reactive T-cell recognition



- Robbins Fig. 6.22 depicts a T cell that recognizes a microbial peptide resembling self.
- The illustration is mechanistic; Individual disease causality still requires evidence.

Rheumatic heart disease: Classic example

Trigger

- Streptococcal proteins initiate antibody responses
- Antibodies target microbial epitopes


Cross-reaction

- Some antibodies recognize myocardial proteins
- Shared antigenic features underlie recognition

Consequence

- Cardiac inflammatory injury can follow
- The textbook links this process with myocarditis

Rheumatic cross-reactivity: Causal chain



- This example supports the plausibility of pathogen–self antigenic cross-reactivity.
- The full pathology of rheumatic heart disease is outside this source unit.

Do not generalize the rheumatic example uncritically

- Rheumatic heart disease supplies a concrete textbook example of molecular mimicry.
- Robbins suggests subtler forms may operate in classic autoimmune disorders.
- The text does not establish specific microbial mimics for every autoimmune disease.
- Shared sequences alone do not prove a pathogenic autoimmune mechanism.

What would support a mimicry hypothesis?


Proposed link	Mechanistic observation	Interpretive limitation
Resemblance	Microbial and self antigens share relevant features	Similarity alone is insufficient
Immune response	Host develops pathogen-reactive lymphocytes or antibody	Response may remain pathogen-specific
Cross-reactivity	Same immune specificity recognizes self	Pathogenicity still needs demonstration
Tissue injury	Self-tissue reaction follows immune recognition	Other inflammatory mechanisms may coexist

Molecular mimicry: Applied classification

**A patient develops cardiac inflammation after streptococcal infection;
Antistreptococcal antibodies bind myocardial proteins.**

- Identify the mechanism: Antigenic cross-reactivity (molecular mimicry).
- Identify the effector highlighted in the source: Antibody.
- Exclude the unsupported leap: Not all postinfectious myocarditis is caused by mimicry.

Viruses may activate B cells polyclonally



- Epstein–Barr virus and HIV as examples of viral B-cell activation.
- This pathway does not require a single microbial antigen to resemble a self antigen.

Epstein–Barr virus and HIV: Source examples

Epstein–Barr virus

- Listed as capable of polyclonal B-cell activation.
- The broad response can include autoantibody production.
- No specific autoimmune diagnosis is inferred here.

Human immunodeficiency virus

- Also listed as causing polyclonal B-cell activation.
- Autoantibodies may result from altered B-cell activation.
- Autoantibody detection does not establish a disease mechanism.


Polyclonal activation versus molecular mimicry

Dimension	Polyclonal B-cell activation	Molecular mimicry
Initiating process	Broad B-cell activation	Response to resembling microbial antigen
Antigenic similarity	Not required by model	Central to model
Potential product	Autoantibodies among other antibodies	Cross-reactive antibody or lymphocyte
Interpretation	Autoantibodies may be nonpathogenic	Cross-reactivity may damage tissue

Autoantibodies do not equal autoimmune disease


- Microbial activation may produce antibodies that recognize self antigens.
- A detectable autoantibody is not, by itself, proof of organ injury.
- Assess whether an immune response can plausibly account for tissue pathology.
- Avoid attributing every postinfection autoantibody to molecular mimicry.

Infection-associated tissue injury releases self antigens



- Infections may expose self antigens independently of microbial–self similarity.
- Antigen availability changes when cells or tissues are injured.

Damage can change the antigen itself



- The textbook links tissue injury with both antigen release and structural modification.
- Neoantigen formation is mechanistically distinct from unmodified antigen exposure.

Antigen release and neoantigen formation

Release of self antigens

- Damage liberates previously sequestered components.
- Antigen becomes available for immune recognition.
- Sequence does not require chemical modification.

Formation of neoantigens

- Tissue injury alters the structure of self components.
- Modified structures can create novel immune determinants.
- T cells may recognize these determinants.

Cytokines create a permissive inflammatory site



- Inflammation can bring potentially autoreactive lymphocytes into antigen-rich tissue.
- Recruitment is a proposed enabling event, not definitive evidence of autoimmunity.

Recruitment links systemic immunity to local injury

Systemic compartment	Inflamed tissue	Outcome
• Self-reactive lymphocytes may circulate • Tolerance normally restrains activity	• Cytokines recruit lymphocytes • Self antigens may be released	• Cells and antigens encounter each other • Tissue-directed reactions may result

Multiple microbial mechanisms can coexist

APC activation

- Costimulation rises
- Previously anergic T cells may respond

Antigenic overlap


- Pathogen-specific recognition overlaps self
- Cross-reactivity is possible

Damage and inflammation

- Self antigens exposed or modified
- Cytokines recruit lymphocytes


These are complementary proposed pathways—not mutually exclusive diagnostic categories.

Environmental insults can change self-antigen display



- Robbins explicitly extends this mechanism beyond infection.
- The tissue microenvironment and antigen display are central to the hypothesis.

Ultraviolet radiation and systemic lupus erythematosus



- This as a proposed explanation for photosensitive lupus flares.
- It does not claim every flare follows UV exposure or that UV is sufficient by itself.

Why nuclear antigens matter after UV injury

Cell injury

- Ultraviolet exposure may cause cell death
- Nuclear material becomes accessible

Antigen availability

- Exposed nuclear constituents can be immunogenic
- Existing autoimmune recognition becomes relevant

Clinical association

- Lupus flares may follow sunlight
- Proposed mechanistic link: Antigen exposure

Sunlight-associated lupus flare: Interpretation


An individual with systemic lupus erythematosus develops a flare after substantial sunlight exposure.

- Proposed initiating event: Ultraviolet-induced cellular injury.
- Potential antigenic consequence: Exposure of nuclear antigens.
- Causal caution: The vignette illustrates the textbook hypothesis, not a unique diagnostic test.

Smoking is a risk factor for rheumatoid arthritis

- Smoking as an environmental risk factor for rheumatoid arthritis.
- A proposed mechanism is chemical modification of self antigens.
- Modified self determinants could become targets of an immune response.
- Risk-factor status is not equivalent to inevitability or to proof of an individual mechanism.

Smoking: Proposed antigen-modification pathway




- The precise chemical modification is not specified in the supplied topic excerpt.
- Avoid presenting molecular intermediates absent from the assigned source.

UV radiation and smoking: Distinct routes

Feature	Ultraviolet radiation	Smoking
Disease association	Lupus flares following sunlight	Rheumatoid arthritis risk
Emphasized perturbation	Cell death and nuclear-antigen exposure	Possible chemical modification of self antigens
Proposed mechanism	More antigen available to immune system	Altered self-antigen structure
Inference limit	Not a sufficient cause of lupus	Not a sufficient cause of RA

Local tissue injury is a generalizable trigger



- Tissue damage from various causes can release self antigens.
- It is the loss of normal antigenic context that unifies multiple exposures.

Environmental association is not mechanistic proof

Observation	What it supports	What it does not establish
Exposure precedes flare	Temporal compatibility	Unique causation
Exposure associated with disease	Population-level relationship	Sufficiency in an individual
Plausible immune pathway	Biological mechanism	Clinical pathogenicity in all cases
Multiple host factors	Multifactorial susceptibility	Single-trigger explanation

The microbiome is an environmental interface

Gut microbiome

- Commensals coexist with mucosal immune cells
- May influence immune differentiation


Skin microbiome

- Commensals interact with barrier immunity
- May modulate immune responses

Autoimmune relevance

- Balance of effector and regulatory T cells may shift
- Human disease-specific effects remain uncertain

Microbes can potentially shift effector–regulatory balance



- Robbins frames this as a possibility rather than a proven causal map for human diseases.
- The direction of effect depends on host–microbe interactions.

Effector and regulatory T cells have opposing roles

Effector T-cell response

- Can drive tissue-directed inflammation.
- May become pathogenic when directed at self.
- Microbial influences could alter its prominence.

Regulatory T-cell response

- Suppresses inappropriate immune activation.
- Contributes to peripheral self tolerance.
- Changes in its maintenance may alter disease risk.

Which microbial communities cause human autoimmunity?

- The source does not identify definitive causal microbes for specific human autoimmune diseases.
- Experimental or population associations need cautious interpretation.
- Host predisposition can confound apparent microbiome–disease links.
- Robbins leaves disease-specific causal attribution unresolved.

Microbiome manipulation: Therapeutic uncertainty

Biological hypothesis

- Commensals help shape immune response.
- Changing communities could alter effector/Treg balance.
- This makes prevention or treatment attractive conceptually.

What Robbins does not establish

- Specific causal organisms in people.
- A validated microbiome manipulation to prevent disease.
- A proven intervention for treating these autoimmune disorders.

Microbiome and pathogen effects are not synonymous

Feature	Acute microbial infection	Commensal microbiome
Typical exposure model	Infection with immune activation	Resident microbial communities
Mechanisms emphasized	APC activation; Mimicry; Injury	Longer-term shaping of immune balance
Potential direction	Can promote autoimmune activation	May shift toward or away
Causality	Several mechanisms postulated	Specific human links unresolved

Infections may also protect against autoimmunity

Potential pathogenic effect

- APC activation increases costimulation.
- Molecular mimicry may create cross-reactivity.
- Tissue injury promotes antigen exposure.

Potential protective effect

- Some infections reduce disease in animal models.
- Protective immune regulation may increase.
- Mechanism is incompletely established.

A population paradox: Infection control and autoimmunity

- Autoimmune disease incidence has increased in some high-resource countries.
- This trend has occurred alongside improved infection control.
- This as epidemiologic context for a protective infection hypothesis.
- Parallel trends alone do not show that reduced infection exposure causes autoimmunity.

Animal models show infection-associated protection

Experimental observation

- In some type 1 diabetes animal models, infections reduce disease incidence.


Possible explanation

- Infections may support regulatory pathways instead of triggering autoimmunity.

Translation limit

- Results in animal models do not prove protection in humans.

IL-2 may help maintain regulatory T cells



- This pathway is explicitly described as one possibility; The mechanism remains unclear.
- IL-2 is important for regulatory T-cell maintenance in the tolerance context.

Triggering and protection can both be plausible

Pathway	Immune consequence	Autoimmune implication
Costimulator induction	Activates self-reactive T cells	Potential triggering
Mimicry	Cross-reactive immune recognition	Potential triggering
Inflammatory injury	Self-antigen exposure and recruitment	Potential triggering
Regulatory maintenance	Supports Treg function	Potential protection

Sex bias is a major feature of autoimmunity

- Many autoimmune diseases occur more frequently in females than in males.
- This with a sex-distribution figure across several disorders.
- The underlying biological explanations remain incompletely understood.
- A population sex bias does not determine risk for an individual patient.

Sex hormones and X-linked genes: Hypotheses


Hormonal contribution

- Sex hormones may modify immune responses.
- Potential relevance to tolerance and disease bias.
- The source does not establish a singular hormonal mechanism.

X-chromosome contribution


- Genes carried on the X chromosome may contribute.
- Potential relevance to sex-associated susceptibility.
- Specific causal gene pathways are not resolved in the excerpt.

Multiple exposures converge on self tolerance



- Genetic background may alter how an infection or insult is interpreted by immunity.
- Autoimmune disease as a multifactorial result, not one exposure with one outcome.

Chronic autoimmune injury can outlast a trigger



- The general-features subsection describes chronicity and relapsing/remitting courses.
- Persistence of self antigens helps explain why disease may continue after an initiating event.

One trigger can intersect several checkpoints

Recognition

- Self-antigen availability or mimicry
- Determines what can be recognized

Activation

- APC costimulation and cytokines
- Determines whether response proceeds

Regulation

- Anergy, Tregs, inhibitory pathways
- Determines whether it is restrained

Interpreting environmental links: Three questions

Question	Example	Why it matters
What changed?	APC costimulation / antigen exposure	Identifies the proposed initiating step
What responds?	Cross-reactive antibody / self-reactive T cell	Identifies the proposed immune effector
What is demonstrated?	Association / model / tissue damage	Prevents overstating human causation

Case: Postinfectious cardiac inflammation

Antibodies raised against a streptococcal protein also recognize myocardial proteins, followed by myocarditis.

- Most direct concept: Molecular mimicry.
- Effector pathway highlighted: Cross-reactive antibody recognition.
- A distinct mechanism from APC costimulation without shared epitopes.

Case: Ultraviolet-associated lupus flare

A person with lupus experiences a flare temporally associated with intense sunlight exposure.

- The implicated insult is ultraviolet radiation, not microbial mimicry.
- The proposed sequence is cell death followed by nuclear-antigen exposure.
- The association is plausible but not sufficient to prove causation in that episode.

Case: Infection without antigen resemblance

An activated APC displays a self peptide and expresses strong B7 costimulators during a microbial infection.

- Which molecular interaction changes? B7–CD28 costimulation.
- Which tolerance process is challenged? T-cell anergy.
- Must microbial peptide resemble self? No.

Case: Unexpected autoantibody during viral infection

Autoantibodies are detected after an infection known to cause broad B-cell stimulation, without evidence of organ injury.

- Consider polyclonal B-cell activation among possible explanations.
- Do not equate autoantibody detection with established autoimmune disease.
- Do not infer molecular mimicry without demonstrated cross-reactivity.

Integrated comparison of environmental mechanisms

Mechanism	Immediate immune change	Textbook example or context
APC costimulation	Self-reactive T-cell activation	Infection-activated APCs
Molecular mimicry	Cross-reactive recognition	Streptococcal proteins / myocardium
Polyclonal B activation	Broad autoantibody production	EBV; HIV
Tissue injury	Self-antigen release / change	Infection; UV; Smoking (proposed)
Regulatory shifts	Effector/Treg balance	Microbiome; Possible IL-2 effects

What is established, and what remains proposed?

Direct textbook statements

- Infections may precipitate autoimmune reactions.
- Smoking is a rheumatoid arthritis risk factor.
- Female predominance characterizes many autoimmune diseases.

Explicit mechanistic uncertainty

- Microbial contributions to specific human diseases are incompletely defined.
- Microbiome interventions are not established here.
- Protective IL-2 and sex-linked explanations remain hypotheses.

Principal teaching conclusions

- Microbial costimulation can convert self-antigen recognition into T-cell activation.
- Molecular mimicry is cross-recognition of microbial and self antigens.
- Viral B-cell activation, tissue injury, and cytokine recruitment offer additional routes.
- Ultraviolet radiation, smoking, and commensal microbiota may change immune context.
- Potential protective effects and sex bias emphasize multifactorial, incompletely resolved causation.