

Schistosomiasis

Pathogenesis • Morphology • Clinicopathologic correlations

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Explain how freshwater transmission and vascular migration establish infection.
- Connect egg-induced immune responses to granulomas and tissue fibrosis.
- Recognize the dominant hepatic, intestinal, pulmonary, renal and urinary patterns.
- Relate characteristic morphology to the major clinical complications.

Schistosomiasis: A tissue-based helminth disease

- Infection by blood flukes of the genus *Schistosoma*.
- Disease expression depends on the schistosome species and organ distribution.
- Persistent host reactions to parasite eggs are central to tissue injury.
- The supplied textbook estimates more than 200 million people infected; This is a textbook figure, not a verified current prevalence estimate.

The two principal organ patterns

Hepatointestinal disease

- *S. Mansoni* and *S. Japonicum* predominate.
- Egg deposition affects bowel and liver.
- Fibrosis and portal hypertension drive severe disease.

Urinary disease

- *S. Haematobium* predominates.
- Egg-induced bladder and ureteral inflammation.
- Hematuria, obstruction and bladder neoplasia association.


Species, setting and predominant injury

Parasite	Geographic setting	Main lesion
S. Mansoni	Latin America; Africa; Middle East	Gut and liver
S. Japonicum	East Asia	Gut and liver
S. Mekongi	East Asia	Severe hepatointestinal involvement
S. Haematobium	Africa	Bladder and urinary tract

Freshwater exposure is the transmission setting

- Freshwater snails are required intermediate hosts.
- Slow-moving tropical rivers, lakes and irrigation ditches support the cycle.
- Human contact with cercaria-containing freshwater enables skin penetration.
- Water and irrigation infrastructure can therefore influence exposure.

Life cycle: Freshwater to human




Schematic

- Eggs entering fresh water can infect snails and renew the cycle.
- Only certain developmental stages are infective to humans.

Cercariae initiate human infection

- Free-swimming cercariae leave the freshwater snail.
- They contact and penetrate human skin.
- Proteolytic enzymes assist passage through the keratinized surface.
- Minimal skin reaction at this stage.

Migration occurs through vascular compartments



Schematic

- Schistosomes disseminate through the circulation after entering the host.
- The hepatic vasculature is the principal site of maturation described for intestinal species.

Liver maturation precedes adult localization

- *S. Mansoni* and *S. Japonicum* reach hepatic vessels during maturation.
- Mature male and female flukes form pairs.
- Pairs subsequently relocate into venous systems where egg production occurs.
- Adult localization differs from the final tissue distribution of trapped eggs.

Adult worm site is not the sole disease site

Adult residence

- Mature worm pairs inhabit venous beds.
- Commonly portal or pelvic veins.
- The vascular location influences where eggs are deposited.

Egg deposition

- Eggs migrate across tissue walls or lodge in organs.
- Local inflammation follows the eggs.
- Injured tissue can be distant from mature worms.

Egg production maintains the inflammatory stimulus

- Adult females release hundreds to thousands of eggs per day (textbook description).
- Eggs secrete proteases that contribute to local inflammatory reactions.
- Continued deposition maintains repeated episodes of tissue injury.
- Both transit and trapping of eggs are pathologically important.

Egg passage produces two excretion pathways

Stool pathway




Urine pathway



Inflammation has a paradoxical role

- Local host inflammation assists movement of eggs through bowel or bladder walls.
- This permits parasite eggs to reach the external environment.
- The same reaction produces mucosal injury and chronic granulomas.
- Transmission biology is therefore linked directly to tissue pathology.

A trapped egg initiates the tissue lesion




Schematic

- The principal target of host injury is often an egg rather than an adult worm.
- Duration and location of egg retention help explain chronic organ damage.

Acute clinical disease follows early infection

- Acute schistosomiasis may present as a severe febrile illness.
- It peaks approximately two months after infection.
- Fever is linked in the source to early inflammatory cytokine responses.
- The chronic dominant lesions develop through retained eggs and fibrosis.

Early immune response: Th1 and fever




Schematic

- Robbins explicitly links the early Th1-skewed response with febrile disease.
- This is an immune-mediated systemic manifestation.

Inflammatory cytokines explain the acute syndrome

- IFN- γ activates macrophages in the early response.
- Activated macrophages secrete TNF, IL-1 and IL-6.
- These cytokines contribute to systemic fever.
- This systemic pathway differs from the later fibrosing tissue response.

Chronic immune response: Profibrotic axis



Schematic

- The source describes Th2 dominance in chronic infection, not total absence of Th1 activity.
- Activated macrophages and Th2 responses are central to severe hepatic fibrosis.

Th1 and Th2: Complementary pathogenic roles

Early Th1-skewed response

- IFN- γ production.
- Macrophage inflammatory cytokines.
- Strong connection to acute fever.


Chronic Th2-dominant response

- Activated profibrotic macrophages.
- Persistent egg-associated granulomas.
- Strong connection to collagenous scarring.

Granulomas involve more than one helper T-cell subset

- Both Th1 and Th2 cells contribute to granulomas around retained eggs.
- Th2-biased activity becomes prominent in chronic infection.
- Granulomas localize a persistent antigenic focus but can injure organs.
- The fibrosis is a host-response consequence of prolonged parasitism.

The schistosome egg is the granuloma focus



Macrophages

Lymphocytes

Eosinophils

Neutrophils


Schematic: Cell types are labeled, not drawn to histologic scale

- The egg may contain a miracidium.
- The cellular infiltrate is explicitly mixed, with eosinophils characteristic of helminth lesions.

The egg changes during lesion evolution

- In early lesions, an egg containing a miracidium lies centrally.
- The miracidium degenerates with time.
- Egg remnants may become calcified.
- Old calcified eggs can coexist with chronic inflammation or scarring.

The liver is injured by portal egg retention




Schematic

- S. Mansoni and S. Japonicum cause the dominant hepatointestinal pattern.
- Portal delivery of eggs explains lesions within hepatic tissue.

Early liver and bowel morphology

- White, pinhead-sized granulomas scatter through gut and liver.
- Each granuloma is centered on a schistosome egg.
- The granulomas may contain a prominent eosinophilic component.
- Early inflammatory foci precede widespread fibrous remodeling.

Granuloma composition is diagnostically meaningful



Macrophages

Lymphocytes

Neutrophils

Eosinophils

Schematic: Cell types are labeled, not drawn to histologic scale

- Helminth-associated granulomas are characteristically eosinophil rich.
- The label is a simplified cellular map, not a genuine histology image.

Pigment deposition is a separate morphologic clue

- Schistosome intestinal pigment is heme-derived and regurgitated by worms.
- The dark pigment accumulates in Kupffer cells and splenic macrophages.
- This pigment is iron-free.
- The hepatic surface or parenchyma may appear darkened.

Colonic disease can form inflammatory patches

- Chronic bowel exposure to eggs produces focal inflammatory lesions.
- Colonic patches can accompany *S. Mansoni* or *S. Japonicum* infection.
- Egg-centered granulomas are the local pathologic substrate.
- The intestinal phenotype may coexist with hepatosplenic damage.


Colonic pseudopolyps are chronic inflammatory change

- Pseudopolyp formation in later infection.
- They reflect patchy mucosal inflammatory remodeling.
- The term describes a protruding lesion, not an epithelial neoplasm.
- Colonic changes should be interpreted within the broader egg-related process.

Advanced hepatic morphology is distinctive

- Liver surface becomes irregular and bumpy.
- Cut surfaces show widened portal zones, fibrosis and granulomas.
- Fibrous expansion may be extensive around portal tracts.
- The source specifically notes absence of intervening regenerative nodules.

Pipestem fibrosis is a portal-centered scar



Schematic

- The name comes from thick fibrous tracks resembling clay pipe stems.
- This morphologic pattern links parasite injury to portal vascular obstruction.

Pipestem fibrosis is not defined by regenerative nodules


Schistosomal pattern

- Broad fibrous portal tracts.
- Portal expansion and obliteration.
- No intervening regenerative nodules described.

Interpretive caution

- Do not equate every fibrotic liver with nodular cirrhosis.
- Specify the observed periportal pattern.
- Portal hypertension can arise without classic cirrhotic nodularity.

Portal vein injury produces portal hypertension



Schematic

- Vascular narrowing is the mechanistic link.
- Clinical sequelae arise from elevated portal venous pressure.

Congestive splenomegaly reflects portal hypertension

- Portal resistance impairs normal splanchnic venous drainage.
- Chronic congestion enlarges the spleen.
- Severe congestive splenomegaly as a consequence.
- The splenic finding reflects hemodynamic disease rather than primary splenic parasitism alone.

Esophageal varices are a major portal consequence

- Portal hypertension promotes collateral venous pathways.
- Esophageal varices arise from the abnormal collateral circulation.
- They mark clinically important portal disease.
- Varices must be connected to portal venous obstruction in mechanistic teaching.

Ascites completes the portal-hypertension phenotype

- Advanced portal fibrosis may be accompanied by ascites.
- Ascites coexists with splenomegaly and varices in severe disease.
- It signifies an important clinical consequence of portal pathology.
- The initiating lesion remains the egg-driven inflammatory response.

Hepatosplenic disease: A causal synthesis




Schematic

- The pathway accounts for the most serious chronic intestinal-species complications.
- Each step connects morphology to clinical physiology.

Pulmonary lesions arise via diverted eggs

- Portal collateral pathways can redirect schistosome eggs to the lungs.
- Eggs lodge in pulmonary vascular beds.
- Granulomatous pulmonary arteritis may follow.
- Pulmonary disease can therefore complicate hepatosplenic schistosomiasis.

Pulmonary vascular remodeling has several stages



Schematic

- Robbins also describes scars and organizing thrombi in affected arteries.
- The lesions share some morphologic features with pulmonary hypertension.

Pulmonary arterial histology: Key features

- Granulomas and scars may disrupt arterial elastic layers.
- Organizing thrombi can occupy narrowed lumina.
- Intimal hyperplasia contributes to obstruction.
- Angiomatoid lesions resembling idiopathic pulmonary hypertension are described.

Cor pulmonale follows severe pulmonary vascular injury

- Progressive arterial obstruction increases pulmonary vascular resistance.
- Right-sided cardiac pressure burden may develop.
- Cor pulmonale as the ultimate complication.
- This complication traces back to displaced eggs and vascular granulomas.

Renal disease can accompany hepatosplenic infection

- Mesangioproliferative glomerulopathy has increased frequency.
- Membranous glomerulopathy is also described.
- These are renal associations in hepatosplenic schistosomiasis.
- The source does not establish a universal renal lesion for all infected patients.

Renal immune deposits: What is actually described

Observed glomerular findings

- Immunoglobulin deposits.
- Complement deposits.
- Membranous or mesangioproliferative morphology.


Important qualification

- Schistosome antigen is rarely found in deposits.
- Do not assume every deposit directly contains parasite antigen.
- Interpret as an associated glomerular process.

S. Haematobium targets the urinary tract

- Adult parasites favor pelvic venous systems in the source narrative.
- Eggs become deposited in bladder and urinary tract tissues.
- The dominant local injury is granulomatous cystitis.
- The clinical phenotype differs substantially from hepatointestinal disease.

Early urinary disease: Egg deposition to hematuria




Schematic

- Mucosal damage directly connects a microscopic inflammatory lesion with visible bleeding.
- Urinary infection can be recognized before late calcification develops.

Urinary granulomas undergo calcification

- As urinary lesions age, eggs and granulomas calcify.
- Numerous calcified foci can create a sandy mucosal appearance.
- Extensive disease can involve large regions of the bladder wall.
- The morphologic sequence reflects chronicity, not merely recent infection.

Calcified bladder: A radiologic correlate




Schematic

- A dense concentric rim when bladder-wall calcification is severe.
- The schematic describes the cause of the appearance; It is not a patient radiograph.

Ureteral fibrosis is a frequent urinary complication

- Inflammation extends to or occurs within ureteral walls.
- Healing produces ureteral fibrosis.
- Progressive narrowing can obstruct urine drainage.
- This as the most frequent complication of S. Haematobium infection.

Obstructive urinary injury: Full causal pathway



Schematic

- Obstruction explains the linkage between local ureteral injury and renal damage.
- The upstream tissue injury originates in granulomatous schistosomiasis.

Hydronephrosis is a downstream consequence

- Fibrotic ureteral narrowing impedes urinary drainage.
- The collecting system becomes dilated (hydronephrosis).
- Persistent obstruction predisposes to chronic renal complications.
- In this infection, renal damage may therefore be obstructive rather than glomerular.

Two different renal pathways should not be conflated

Hepatosplenic infection

- Mesangioproliferative or membranous glomerulopathy.
- Immunoglobulin and complement deposits.
- Association with severe intestinal-species disease.


Urinary schistosomiasis

- Ureteral inflammation and fibrosis.
- Obstruction and hydronephrosis.
- Chronic pyelonephritis.

Urinary schistosomiasis and bladder carcinoma

- S. Haematobium infection is associated with bladder squamous cell carcinoma.
- The source describes squamous metaplasia as a typical precursor change.
- Chronic inflammation provides the tissue context for the association.
- This is an association; No incidence figure is supplied.

Chronic cystitis creates a metaplastic context



Schematic

- Do not present the diagram as a deterministic fate for every patient.
- The pathologic association matters for urinary tract evaluation.

Dominant pathology by organ

Organ	Characteristic morphology	Major consequence
Liver	Periportal granulomas; Pipestem fibrosis	Portal hypertension
Colon	Granulomas; Inflammatory patches; Pseudopolyps	Chronic intestinal injury
Pulmonary arteries	Granulomatous arteritis; Intimal thickening	Cor pulmonale
Bladder / ureter	Cystitis; Calcification; Ureteral scarring	Hematuria; Obstruction

Mechanism-to-manifestation matrix

Mechanism	Morphologic lesion	Clinical correlate
Th1-related cytokines	Systemic inflammatory activity	Acute fever
Egg-centered granulomas	Hepatic / bowel inflammatory foci	Chronic tissue injury
Portal vein obliteration	Pipestem fibrosis	Portal hypertension
Urothelial erosion	Granulomatous cystitis	Hematuria

Diagnostic sampling follows egg exit routes

- Intestinal species release eggs through bowel walls into stool.
- *S. Haematobium* releases eggs through bladder tissue into urine.
- The correct specimen depends on the suspected species and organ pattern.
- The lecture source supports egg-shedding routes but does not provide test sensitivity or formal diagnostic algorithms.


Case: Hepatosplenic disease

Clinicopathologic vignette

A patient from a freshwater-exposure setting has severe splenomegaly, ascites and esophageal varices. Liver tissue is described as portal-expanded, fibrotic and granulomatous without regenerative nodules.

Which parasite-associated mechanism best unifies the liver morphology and portal findings?

Case reasoning: Portal egg injury



Schematic

- Venous obstruction, rather than nodular regeneration, explains the clinical syndrome.
- The most relevant parasites are *S. Mansoni* or *S. Japonicum* with scope.


Case: Chronic urinary disease

Clinicopathologic vignette

A patient with freshwater exposure has hematuria and a dense calcified bladder-wall rim. Later evaluation demonstrates ureteral narrowing and hydronephrosis.

What tissue lesion links these findings, and which *Schistosoma* species is implicated?

Case reasoning: Chronic urinary egg lesions



Schematic

- Long-standing egg deposition explains both the bladder wall and ureteral pathology.
- Bladder squamous carcinoma is an important associated long-term neoplastic lesion.

Do not confuse the dominant pathologic outcomes

Hepatic / portal

- Eggs reach liver through portal blood.
- Periportal scarring obstructs portal veins.
- Varices, splenomegaly and ascites are downstream.

Urinary / ureteral

- Eggs traverse and lodge in bladder tissue.
- Cystitis, calcification and ureteral scars develop.
- Hematuria and obstructive renal injury are downstream.

Schistosomiasis: Integrating the mechanisms

- The disease begins with freshwater cercarial penetration and vascular migration.
- Adult worm location shapes where eggs accumulate.
- Egg-driven granulomatous inflammation explains organ injury.
- Chronic fibrosis governs portal and urinary obstruction.
- Pulmonary, renal and neoplastic associations should be recognized separately.

Reading schistosomiasis tissue: key questions

- Is an egg visible in the center of a granulomatous lesion?
- Are eosinophils present in the mixed inflammatory infiltrate?
- Is there evidence of egg degeneration, calcification or chronic fibrosis?
- Does the tissue distribution indicate portal, bowel or urinary disease?
- Which vascular, obstructive or epithelial complication follows the observed lesion?