
Sickle Retinopathy, Retinal Vasculitis, and Radiation Retinopathy

Retinal vascular injury, ischemia, neovascularization, and detachment

Dr Abu Suraih Sakhri


MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Identify the shared ischemia-driven mechanism across the three retinal vascular disorders.
- Relate sickle-cell vaso-occlusion to hemorrhage, characteristic fundus lesions, and sea-fan neovascularization.
- Explain how vascular damage in vasculitis or after irradiation may lead to angiogenesis.
- Trace retinal neovascularization to hemorrhage, membrane organization, traction, and detachment.


Vascular injury is the organizing concept



The initiating lesion differs among the three conditions.

Downstream consequences converge on hemorrhage, traction, and potential retinal detachment.

The retina is a layered neurosensory tissue



- Neurosensory retina lies internal to retinal pigment epithelium (RPE).
- Hemorrhage position determines its structural context.
- Vitreous lies immediately internal to the retinal surface.


Retinal vascular injury can have two outputs

- Blood-retinal barrier disturbance permits leakage, edema, hemorrhage, and exudation.
- Vascular occlusion produces focal or regional retinal ischemia.
- Ischemia induces proangiogenic signaling; Abnormal new vessels may form.
- Neovascular membranes create mechanical as well as vascular complications.

Occlusion and vessel-wall injury: Distinguish the starting event


Feature	Occlusion-dominant	Vessel injury–dominant
Initiating event	Impaired microvascular flow	Damage to existing retinal vessels
Exemplar in this lecture	Sickle retinopathy	Peripheral vasculitis; Irradiation
Shared consequence	Retinal ischemia / nonperfusion	Retinal ischemia / nonperfusion
Potential outcome	Growth-factor response and neovascularization	Growth-factor response and neovascularization

Nonperfusion connects upstream and downstream pathology



The retinal periphery is particularly important for sickle retinopathy.
Ischemia is a mechanism; New-vessel formation is a response to that mechanism.

Growth-factor signaling precedes proliferative lesions




VEGF is a shared mediator in ischemic retinal vascular disease.

VEGF and basic fibroblast growth factor are both implicated in sickle-retinal angiogenesis.

Anatomic location separates stages of vascular retinopathy


Pattern	Predominant location	Underlying phenomenon
Nonproliferative	Within retina	Angiopathic and hemorrhagic lesions
Proliferative	Retinal surface or preretinal region	New abnormal vascular growth
Mechanical complication	Vitreoretinal interface	Organized membrane pulls on retina

Internal limiting membrane marks the retinal surface




- Retina is bounded internally by the internal limiting membrane (ILM).
- Preretinal hemorrhage is superficial to the neural retina.
- New-vessel membranes can interact with posterior vitreous.

The vitreoretinal interface converts growth into traction



Hemorrhage and fibroglial tissue organization change surface architecture.
A vessel lesion can evolve into a mechanical threat to retinal attachment.

Retinal hemorrhages arise in different compartments




- Preretinal blood is adjacent to the vitreoretinal interface.
- Intraretinal blood occupies neural retinal layers.
- Subretinal blood lies external to neurosensory retinal tissue.

Location predicts the appearance of retinal hemorrhage

Location	Pathologic orientation	Clinical appearance
Nerve fiber layer	Parallel to retinal surface	Flame-shaped hemorrhage
Deeper retinal layers	Perpendicular to surface	Dot-like appearance
Preretinal / subhyaloid	Between retinal surface and posterior hyaloid	Superficial blood at vitreoretinal interface

Hemorrhage and its organization can precede detachment



This sequence is specifically described in sickle retinopathy.

It highlights why even prior hemorrhage can have later structural consequences.


Tractional and rhegmatogenous detachment differ

Characteristic	Tractional / nonrhegmatogenous	Rhegmatogenous
Primary mechanism	Pull from an organized surface membrane	Full-thickness retinal break allows vitreous fluid access
Retinal tear	Not required initially	Required
Important nuance	Severe traction can tear retina	May follow severe traction

Three disorders, one ischemic retinal response

Disorder	Primary upstream insult	Convergent consequence
Sickle retinopathy	Low-O ₂ -induced red-cell sickling and occlusion	Peripheral ischemia; Angiogenesis
Retinal vasculitis	Injury to retinal vessels	Ischemia; Neovascularization
Radiation retinopathy	Retinal vascular injury following irradiation	Ischemia; Neovascularization


Sickle hemoglobinopathy begins at the molecular level



The hemoglobin mutation is the initiating systemic mechanism.

The retinal lesion reflects its downstream microvascular consequences.

Deoxygenation makes HbS erythrocytes less deformable




Schematic cellular forms — not a blood film

- HbS polymerization changes intracellular physical properties.
- Progressive sickling distorts red cells.
- Distorted, stiff cells participate in microvascular obstruction.

Why microvascular transit matters

- Slow capillary transit permits deoxygenated HbS to polymerize.
- Sickled erythrocytes become mechanically less deformable.
- Increased red-cell adhesion and local flow slowing can further promote obstruction.
- Reduced perfusion creates conditions for additional regional hypoxia.

Vaso-occlusion can reinforce itself



Repeated obstruction may sustain a cycle of impaired perfusion and deoxygenation.
The retinal periphery provides a particularly important low-oxygen microenvironment.


Two genotypic settings for nonproliferative sickle retinopathy

Hemoglobin state	Status in this topic	Interpretation
HbSS	Nonproliferative retinal changes described	Sickle hemoglobinopathy with ocular risk
HbSC	Nonproliferative retinal changes described	Compound HbS/HbC sickling disorder
HbAS sickle trait	Ocular profile not established here	Requires separate evidence

HbSC is a genuine sickling disorder

- HbSC red cells contain substantial HbS, unlike normal HbAA erythrocytes.
- HbC-associated cellular dehydration raises intracellular HbS concentration.
- Deoxygenation can therefore trigger polymerization and microvascular manifestations.
- Both HbSC and HbSS occur among genotypes associated with nonproliferative sickle retinopathy.

Low oxygen tension emphasizes the retinal periphery




Schematic retinal field — not an ophthalmoscopic image

- The peripheral retinal microcirculation is the emphasized site of sickling.
- Regional vaso-occlusion generates nonperfused retina.
- The ischemic territory may subsequently drive angiogenic signaling.

Nonproliferative sickle retinopathy: Core concept

- The nonproliferative stage is characterized by intraretinal angiopathic changes.
- Occlusions contribute to preretinal, intraretinal, and subretinal hemorrhages.
- Resolved hemorrhages generate recognizable ophthalmoscopic signs.
- Proliferation is a separate stage characterized by neovascularization.

Sickle-cell hemorrhages span retinal compartments



- Preretinal, intraretinal, and subretinal hemorrhage are all described.
- Their different positions imply different surface and tissue effects.
- Organization of preretinal blood is particularly relevant to traction.

Preretinal hemorrhage can become a traction source

- Blood accumulates at or near the retinal surface.
- An organized preretinal hemorrhage can produce fibrous tissue.
- Contraction may pull on the underlying retina.
- The resulting traction can contribute to retinal detachment.


Intraretinal hemorrhage indicates vascular injury inside retina

- Occluded retinal microvessels can be associated with intraretinal bleeding.
- Hemorrhage occurs within the layered neurosensory tissue.
- Its subsequent resolution may leave clinically visible abnormalities.
- Intraretinal disease does not itself establish proliferative retinopathy.

Subretinal hemorrhage occupies the RPE–retina interface

- The hemorrhage lies external to the neurosensory retina.
- This compartment may be involved in nonproliferative sickle retinopathy.
- It is anatomically distinct from preretinal and intraretinal bleeding.
- Site must not be confused with the presence of a full-thickness retinal tear.


Salmon patches: A residual hemorrhagic sign



Schematic retinal field — not an ophthalmoscopic image

- An ophthalmoscopically recognized finding in sickle retinopathy.
- Appears among changes associated with resolution of retinal hemorrhage.
- This finding alone does not establish a sea-fan proliferative lesion.


Iridescent spots: A recognizable posthemorrhagic change



Schematic retinal field — not an ophthalmoscopic image

- Iridescent spots may remain following resolution of hemorrhage.
- They belong to the described nonproliferative lesion spectrum.
- An illustrated mark is not a substitute for an authentic clinical photograph.

Black sunburst lesions: Peripheral pigmented-appearing marks




Schematic retinal field — not an ophthalmoscopic image

- Named ophthalmoscopic finding in sickle retinopathy.
- These lesions are grouped with changes following hemorrhage resolution.
- Their appearance should not be mistaken for the sea-fan neovascular lesion.

Three classic fundus descriptors do not mean the same thing


Finding	Pathologic association	Teaching interpretation
Salmon patches	Hemorrhage-resolution changes	Recognize hemorrhagic history
Iridescent spots	Hemorrhage-resolution changes	Recognize nonproliferative aftermath
Black sunburst lesions	Hemorrhage-resolution changes	Do not confuse with sea-fan growth
Sea fans	Florid peripheral neovascularization	Identify proliferative response

Nonproliferative lesions reflect vascular occlusion



Salmon patches, iridescent spots, and black sunbursts are named results of hemorrhage resolution. They illustrate the morphological footprint of earlier microvascular disease.

Ischemia explains the transition to neovascular disease



An ischemic retinal microenvironment can drive proliferative disease.
The same general sequence appears in several retinal vascular conditions.


Two growth factors named in sickle retinopathy

- Vascular endothelial growth factor (VEGF) is upregulated with retinal ischemia.
- Basic fibroblast growth factor (bFGF) is also implicated in the angiogenic response.
- These signals are linked to retinal angiogenesis downstream of vascular occlusion.
- Their mention does not establish a treatment response or biomarker cut-off.

Proliferative sickle retinopathy is defined by new vessels

- The clinically important transition is the appearance of retinal neovascularization.
- The new vessels are especially prominent in the retinal periphery.
- Florid peripheral retinal neovascularization forms characteristic “sea fans.”
- The consequence of proliferation is not limited to bleeding: Organized tissue may exert traction.


Sea fans: The characteristic proliferative lesion



Schematic retinal field — not an ophthalmoscopic image

- Peripheral florid retinal neovascularization.
- Linked to occlusion, ischemia, and local angiogenic growth factors.
- Represents proliferative, not merely hemorrhagic, retinal disease.


Location of sea fans matters to disease mechanism



Schematic retinal field — not an ophthalmoscopic image


- A peripheral ischemic field is the stated upstream context.
- The neovascular lesion occupies the peripheral retinal region.
- Bleeding and traction may arise from pathological vessels or their organization.

New vessels create hemorrhagic consequences



Pathologic retinal angiogenesis may be followed by bleeding.
Retinal neovascularization can bleed into the avascular vitreous.


Organization turns hemorrhagic lesions into traction



Mechanical effects can continue after an initial bleeding event.

Traction couples vascular pathology to anatomy of vitreous and retinal attachment.

Traction can separate neurosensory retina from RPE



- Neovascular or hemorrhagic material can organize on the retinal surface.
- Contraction transmits a pulling force onto the retina.
- Separation from RPE defines retinal detachment.

Sickle nonproliferative vs proliferative disease

Feature	Nonproliferative	Proliferative
Main process	Intraretinal angiopathy / hemorrhage	New retinal vessels
Named signs	Salmon patches, iridescent spots, black sunbursts	Sea fans
Ischemic mechanism	Occlusion already present	Hypoxic growth-factor upregulation
Major structural concern	Organization of preretinal blood	Hemorrhage and fibrovascular traction

Application: Hemorrhage versus proliferation

Teaching vignette — hypothetical

A teaching fundus description in a person with HbSS includes multiple residual hemorrhagic lesions but no stated peripheral sea-fan neovascularization.


Interpretation prompts

- Which disease phase is described?
- Which findings would establish a proliferative lesion?
- What additional structural complication can follow preretinal hemorrhage?

Interpreting the sickle case

- Residual hemorrhagic lesions are compatible with nonproliferative morphology.
- Peripheral sea-fan neovascularization would indicate proliferative disease.
- The absence of an explicitly described sea fan should not be read as proof of absent angiogenesis.
- Organized preretinal hemorrhage can exert traction even in the discussed nonproliferative pathway.

Sickle retinopathy: Two related paths from occlusion




Both outcomes emerge from occlusive microvascular disease.

Retinal hemorrhages and neovascularization are linked but are not synonymous.

Retinal vasculitis begins with injury to retinal vessels

- Peripheral retinal vasculitis can be associated with retinal neovascularization.
- Vessel-wall damage, not HbS polymerization, is the initiating disease process.
- Injured retinal vessels can create nonperfused ischemic zones.
- The morphology does not, by itself, identify a specific systemic vasculitis.


Peripheral vasculitis can leave ischemic retinal sectors



Schematic retinal field — not an ophthalmoscopic image


- Vessel injury can impair perfusion in an affected retinal region.
- Ischemic zones provide an angiogenic stimulus.
- A schematic ischemic sector does not establish a particular angiographic pattern.

Vasculitis: Vascular damage to abnormal angiogenesis



The causal sequence is explicitly grouped with sickle and radiation retinopathy.
No particular inflammatory cause is assumed here.

Complications of vasculitis-associated proliferation



The same mechanical end-stage may follow distinct initiating conditions.

Recognizing vessel damage is more informative than labelling all cases as sickle disease.


Retinal ischemia does not identify the vasculitis cause

- Retinal vasculitis is one route to retinal vessel damage and ischemia.
- The downstream ischemic pattern alone does not identify an etiology.
- Specific etiologic diagnosis requires additional clinical and laboratory evidence.
- Regardless of cause, ischemia-driven neovascularization may lead to traction.

Radiation retinopathy: The initiating clinical setting

- Irradiation for an intraocular tumor provides the initiating clinical context.
- Retinal vascular damage is the highlighted pathological link.
- Resulting nonperfusion may promote ischemia-driven angiogenesis.
- Retinal morphology alone does not establish an irradiation dose or exposure timeline.


Radiation injury converges on retinal nonperfusion



Exposure and endothelial/vascular injury are conceptually upstream.

The central mechanistic link is vascular nonperfusion followed by angiogenesis.


Radiation-associated retinal injury may create ischemic zones



Schematic retinal field — not an ophthalmoscopic image

- After irradiation, vessel damage can disturb retinal perfusion.
- Regions of ischemia provide the substrate for new-vessel signaling.
- This is a mechanistic schematic, not a post-radiotherapy clinical photograph.

Radiation retinopathy: Proliferative consequences



The result parallels the downstream events in vasculitis.

No timing or incidence claim is made beyond the supplied textbook passage.


Why radiation retinopathy is grouped with vasculitis

- Neither requires sickle hemoglobin polymerization as the initiating insult.
- Both can injure retinal vessels and leave poorly perfused retinal tissue.
- Neovascular growth is a secondary reaction to this ischemia.
- Bleeding and membrane traction can threaten retinal attachment.

Vasculitis and irradiation: Different causes, same pathway

Feature	Peripheral retinal vasculitis	Radiation retinopathy
Initiating setting	Inflammatory vessel injury	Irradiation for intraocular tumor
Primary retinal problem	Vessel damage / nonperfusion	Vessel damage / nonperfusion
Ischemic response	Retinal angiogenesis	Retinal angiogenesis
Possible complications	Hemorrhage, traction, detachment	Hemorrhage, traction, detachment

All three converge on one pathobiologic sequence



Sickle disease begins in red cells; Vasculitis and radiation begin in damaged vessels.
This shared end pathway is why they are taught together.


Identify the initiating lesion before assigning a diagnosis

Disorder	Supported upstream mechanism	What is not justified from morphology alone
Sickle retinopathy	HbS-related peripheral microvascular occlusion	Do not infer hemoglobin genotype from a sea fan alone
Retinal vasculitis	Injury to vessels in peripheral retinal vasculitis	Do not infer a named systemic disorder
Radiation retinopathy	Irradiation-associated vessel damage	Do not infer dose or timing

Shared VEGF biology links neighboring retinal disorders

- Diabetic retinopathy also features ischemia-associated VEGF upregulation.
- Retinopathy of prematurity also involves proliferation following retinal ischemia.
- The three conditions in this lecture share the broad ischemia-to-angiogenesis logic.
- Identical signaling logic does not imply identical causes, classifications, or treatments.

Why tractional retinal detachment is a distinct complication



- Vitreoretinal membranes exert a mechanical pull on neural retina.
- Detached neurosensory retina separates from underlying RPE.
- Severe traction can ultimately cause a tear and superimposed rhegmatogenous detachment.

Three brief scenarios: Identify the mechanism

Teaching vignette — hypothetical

A: HbSC disease with peripheral hemorrhagic lesions.

B: Peripheral retinal vasculitis and zones of nonperfusion.

C: Intraocular irradiation followed by retinal neovascularization.

Interpretation prompts

- Which upstream lesion fits each?
- What downstream pathway do all three share?
- Which findings represent proliferative rather than hemorrhagic disease?

Answers: Recognize mechanism before complication

Scenario	Upstream abnormality	Downstream concern
A — HbSC	Peripheral red-cell sickling and occlusion	Hemorrhage ± sea-fan neovascularization
B — vasculitis	Retinal vessel damage	Ischemia, neovascularization and traction
C — irradiation	Radiation-associated vessel injury	Ischemia, neovascularization and traction

Three routes to retinal structural damage

Starting point	Intermediate process	Possible endpoint
Sickle preretinal hemorrhage	Organization of blood	Retinal traction / detachment
Sickle sea-fan neovascularization	Preretinal abnormal vascular tissue	Hemorrhage / traction
Vasculitis or irradiation	Vascular damage → ischemic neovascularization	Hemorrhage / traction / detachment

Clinicopathologic synthesis

- Primary lesion: Intraretinal vascular occlusion or damage.
- Ischemic retina upregulates proangiogenic signals; VEGF is particularly emphasized.
- Proliferative new vessels are distinct from residual nonproliferative hemorrhagic lesions.
- Hemorrhage and organized tissue produce traction, potentially separating retina from RPE.

Clinical distinctions worth retaining

- Salmon patches, iridescent spots, and black sunbursts follow retinal hemorrhage; sea fans indicate neovascularization.
- Sickle retinopathy begins with HbS-related microvascular obstruction, especially in the retinal periphery.
- Retinal vasculitis and irradiation damage vessels, but both can converge on ischemic angiogenesis.
- Hemorrhage, membrane organization, and traction explain how a vascular disorder may culminate in retinal detachment.