

Spinocerebellar Degenerations

Neurodegeneration across cerebellar, spinal, and peripheral systems

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Define the common tissue targets and clinical heterogeneity of spinocerebellar degenerations.
- Distinguish dominant SCAs from Friedreich ataxia and ataxia-telangiectasia.
- Compare coding-repeat, noncoding-repeat, and conventional SCA mutations.
- Connect FXN and ATM abnormalities to their characteristic clinical syndromes.

One umbrella: Three vulnerable systems

Cerebellum

Neuronal dysfunction and loss contribute to disordered coordination.

Spinal cord

Degeneration contributes to motor and sensory manifestations.

Peripheral nerves

Nerve involvement can add weakness, impaired sensation, and reflex changes.

The commonality is anatomic vulnerability—not one shared molecular cause.

Definition: A heterogeneous degenerative group

- Shared pathology: Neuronal loss and dysfunction in cerebellum, spinal cord, and peripheral nerves.
- The group spans distinct genotypes, ages at onset, and clinical presentations.
- Classification increasingly follows genetic analysis rather than appearance alone.

Anatomical convergence does not imply genetic uniformity.

Heterogeneity occurs along three axes

Genetic etiology

Dominant SCA families;
Recessive Friedreich ataxia
and ataxia-telangiectasia.

Age at onset

Childhood-onset
syndromes coexist with
disorders having variable
onset.

Clinical syndrome

Ataxia may be
accompanied by sensory,
motor, ocular, cardiac, or
immune findings.

Anatomical involvement explains clinical overlap

- Cerebellar dysfunction makes ataxia a common presenting theme.
- Spinal and peripheral nerve injury broadens the syndrome beyond isolated ataxia.
- Similar motor findings may therefore arise from different inherited diseases.
- Extraneural manifestations may point toward particular etiologies.

A genetically useful first division

Dominant group

Spinocerebellar ataxias: A large group of autosomal-dominant inherited degenerations.

Recessive group A

Friedreich ataxia:
Progressive ataxia with neuropathy and cardiomyopathy.

Recessive group B

Ataxia-telangiectasia:
Childhood ataxia with telangiectasias and immunodeficiency.

A shared syndrome is not one disease

- Phenotypic overlap is substantial despite distinct causal lesions.
- Molecular categories organize the SCAs more effectively than ataxia alone.
- Friedreich ataxia exemplifies mitochondrial dysfunction from reduced frataxin.
- Ataxia-telangiectasia links neurodegeneration with impaired DNA-damage signaling.

What a neuropathologist should infer—and not infer

Supported finding

Neuronal loss and dysfunction affect multiple connected neural compartments.

Reasonable interpretation

Multisystem injury can produce combinations of gait, limb, reflex, and sensory abnormalities.

Not established

A single microscopic signature or unified lesion explains all disorders.

Distinguish the textbook's shared localization from a universal histologic pattern.

Molecular diagnosis changes classification

- More than 40 SCA subtypes are distinguished through genetic, clinical, and pathologic information.
- Genetic causes have been found for more than half of these SCA entities.
- This progress in classification has not produced complete mechanistic understanding.
- No corresponding universal effective treatments.

Spinocerebellar ataxias: Defined by inheritance

- “SCA” denotes a family of autosomal-dominant inherited spinocerebellar degenerations.
- The term encompasses many distinct genotypes rather than a single mutation.
- Molecular mechanisms and accompanying clinical findings vary across types.

SCA classification is multidimensional

Genetic

Causative variant or repeat-expansion locus.

Clinical

Ataxia plus subtype-associated neurological or other findings.

Pathologic

Patterns of nervous-system degeneration and cellular involvement.

More than 40 SCA subtypes

- More than 40 molecularly and clinicopathologically defined subtypes.
- A numbered SCA designation identifies a distinct genetically characterized category.
- Subtype diversity prevents a single genotype–phenotype description from covering all SCAs.

Use the SCA number together with a genetic mechanism—not as an explanation by itself.

Three classes of SCA mutations

Coding CAG expansions

Translated polyglutamine tracts; Proteins accumulate in neuronal nuclear inclusions.


Noncoding repeat expansions

Expanded repeats outside protein-coding sequences; Disease mechanism remains unclear.

Conventional mutations

Copy-number changes, deletions, insertions, and point mutations affect diverse proteins.

Coding expansion: Repeat to protein



Protein inclusion formation as a hallmark, not proof of one injury pathway.

Why CAG expansion changes the protein

- A CAG triplet encodes the amino acid glutamine.
- When expanded within a coding sequence, it produces an elongated polyglutamine tract.
- The affected protein is mutant despite retention of its broader protein identity.
- Accumulation of this protein can be demonstrated as nuclear inclusions.

Polyglutamine inclusions: Correct interpretation

- The source reports nuclear inclusion accumulation in the relevant SCA variants.
- The inclusion pattern suggests mechanisms analogous to Huntington disease.
- It does not identify the precise toxic step or establish inclusions as the sole cause.

Polyglutamine SCA spectrum: Named examples

SCA1, SCA2

Both are classified among CAG/polyglutamine-associated SCAs.

SCA3, SCA6

Both are also included in the polyglutamine class.

SCA7, SCA8, SCA17

Listed in the Robbins polyglutamine group; SCA7 has visual impairment.

Machado–Joseph disease corresponds to SCA3

- Spinocerebellar ataxia type 3 as Machado–Joseph disease.
- It belongs in the group associated with coding CAG expansion and polyglutamine.
- The synonym is a disease-name correspondence, not a separate mechanistic category.

SCA7: Visual impairment is a useful clue

- SCA7 is included among the polyglutamine spinocerebellar ataxias.
- Visual impairment as a relatively distinguishing feature.
- A visual complaint can help refine the pattern of an otherwise ataxic syndrome.

Polyglutamine disorders extend beyond numbered SCAs

- Dentatorubral-pallidoluysian atrophy (DRPLA) in the polyglutamine disease group.
- This classification is mechanistic; DRPLA is not itself a numbered SCA subtype.
- Shared repeat chemistry can link disorders that differ in clinical designation.

Polyglutamine SCAs versus Huntington disease

Common molecular motif

A coding CAG repeat expansion leads to a polyglutamine-containing mutant protein.


Common cellular theme

Abnormal proteins accumulate; Nuclear inclusion patterns are reported.

Distinct clinical entities

Huntington disease and SCAs differ in dominant neuroanatomic and clinical manifestations.

Noncoding repeat expansions: What is known



Do not depict an unproven toxic RNA/protein mechanism as established for all subtypes.

Six noncoding-repeat SCAs

SCA10 • SCA12

Explicitly assigned to the
noncoding-repeat class.

SCA27B • SCA31

Explicitly assigned to the
noncoding-repeat class.

SCA36 • SCA37

Explicitly assigned to the
noncoding-repeat class.

Known locus does not mean known mechanism

- For most noncoding-repeat SCAs, the implicated genetic locus has been defined.
- The causal steps connecting repeat expansion to degeneration are not established in the source.
- Do not infer that every noncoding expansion acts through an identical toxic product.

Coding versus noncoding expansions

Coding CAG repeats

Translated into expanded polyglutamine tracts within mutant proteins.

Noncoding repeats

Disease-associated expansions without the same direct polyglutamine coding mechanism.

Both are repeat-expansion diseases, but their demonstrated molecular consequences differ.

Conventional SCA mutations: Variant classes

- Copy-number variation alters the number of copies of a genetic segment.
- Deletions remove DNA sequence; Insertions add sequence.
- Point mutations change individual sequence positions.
- Robbins groups the remaining SCAs under these conventional mutation mechanisms.

Affected protein pathways in conventional SCAs

Receptors

Altered signaling at cellular reception or transduction steps.

Ion channels

Altered proteins involved in membrane electrical signaling.

Kinases

Altered proteins participating in signal-transduction cascades.

These are protein categories named; No specific variant–channel pairing is implied.

Three SCA mutation classes compared

Feature	Coding CAG	Noncoding repeats	Conventional variants
Primary lesion	Coding triplet expansion	Expansion outside coding tract	CNV / indel / point mutation
Protein consequence	Expanded polyglutamine	Not a common direct polyQ mechanism	Diverse signaling proteins affected
Source mechanism	Intranuclear mutant protein	Unclear	Altered receptors / channels / kinases

What genetic classification can establish

- A specific SCA locus or mutation can define the inherited disease category.
- Mutation class gives a mechanistic starting point for biological interpretation.
- A genetic label alone does not specify an exact age at onset or clinical course.
- Across SCAs, causal cellular cascades remain incompletely resolved.

Avoid overinterpreting nuclear inclusions

- Nuclear accumulation is reported in the polyglutamine SCA group.
- The finding supports abnormal processing or persistence of mutant protein.
- It does not demonstrate that every SCA subtype has such inclusions.
- A conventional mutation or noncoding expansion should not be equated with polyQ pathology.

Friedreich ataxia: Autosomal-recessive degeneration

- Friedreich ataxia is recessively inherited, unlike the dominant SCA group.
- Its syndrome includes progressive ataxia, weakness, spasticity, and sensory neuropathy.
- Cardiomyopathy makes the disease systemically consequential.

Friedreich ataxia: Typical clinical sequence

Early childhood

Gait ataxia, often beginning in the first decade



Progression

Hand clumsiness and dysarthria follow



Broader disability

Motor, sensory, and systemic complications accumulate

Motor examination contains mixed signs

- Progressive weakness and spasticity contribute to motor disability.
- Deep tendon reflexes are reduced or absent.
- An extensor plantar response is nevertheless typically present.
- The juxtaposition reflects involvement beyond a pure peripheral neuropathy.

Why the reflex pattern is diagnostically instructive

Depressed reflexes

Compatible with damage to the afferent or peripheral components of the reflex arc.

Extensor plantar response

Suggests dysfunction of descending motor pathways.

Both findings may coexist in a single multisystem neurodegenerative condition.

Sensory examination: Proprioception is prominent

- Joint-position sense is impaired.
- Vibratory sense is impaired.
- Pain, temperature, and light touch may also be reduced in some patients.
- Sensory abnormalities contribute to the complexity of the ataxic syndrome.

Friedreich ataxia: Orthopaedic associations

Pes cavus

A high-arched foot is a commonly observed deformity.

Kyphoscoliosis

Spinal curvature can arise as the neurological illness progresses.

Cardiomyopathy is a major determinant of outcome

- Friedreich ataxia is accompanied by cardiomyopathy.
- Arrhythmias and congestive heart failure occur with high frequency.
- Cardiac disease as contributing to many deaths.
- Cardiac manifestations distinguish this syndrome from isolated cerebellar dysfunction.

Diabetes is an associated systemic feature

- Concomitant diabetes in up to one quarter of affected individuals.
- This is a reported association, not a universal diagnostic criterion.
- An ataxia syndrome accompanied by cardiac and metabolic disease warrants a systemic framework.

Friedreich ataxia: Neurological and systemic domains

Neurological

Gait and limb ataxia;
Weakness; Spasticity;
Dysarthria; Sensory loss.

Skeletal

Pes cavus and
kyphoscoliosis in many
affected people.


Systemic

Cardiomyopathy and
associated diabetes.

Course and prognosis in the supplied Robbins text

- The source describes a relentlessly progressive loss of mobility.
- It reports that many affected people require a wheelchair within roughly five years of onset.
- It describes a typical life expectancy limited to approximately 40–50 years.
- These are textbook-level generalizations, not individualized prognostic predictions.

FXN intronic expansion: From DNA to expression



The FRDA mutation is noncoding—but distinct from SCAs

- Friedreich ataxia involves GAA expansion in the first intron of FXN.
- The gene lies on chromosome 9 in the supplied source.
- The expansion lowers frataxin expression through altered chromatin structure.
- “Noncoding expansion” describes mutation location; It does not place FRDA among dominant SCAs.

Frataxin and the mitochondrion

Protein identity

Frataxin is a mitochondrial protein encoded by FXN.


Functional role

It participates in assembly of iron–sulfur cluster enzymes.

Affected systems

Mitochondrial complexes I and II.

Reduced frataxin compromises bioenergetics



Iron accumulation adds a potential injury mechanism

- Reduced frataxin is associated with increased mitochondrial free iron.
- Free iron can contribute to oxidative stress.
- The textbook links oxidative phosphorylation deficiency and oxidative stress as consequences.
- It does not quantify the individual contribution of each pathway to neurodegeneration.

Friedreich ataxia has two reported genotypes

Common configuration

GAA-repeat expansion present in both FXN alleles.

Less common configuration

Expansion in one allele with a loss-of-function point mutation in the other.

Both configurations are compatible with recessive disease.

A unified pathogenesis for Friedreich ataxia

- An inherited FXN lesion reduces expression of mitochondrial frataxin.
- Iron–sulfur enzyme assembly and respiratory-chain function become impaired.
- Oxidative metabolism deteriorates while free iron may amplify cellular stress.
- These cellular disturbances accompany progressive neurological and systemic illness.

Ataxia-telangiectasia: A multisystem recessive syndrome

- Ataxia-telangiectasia is autosomal recessive.
- Childhood ataxia and dyskinesia coexist with telangiectasias and immunodeficiency.
- DNA-damage response dysfunction unites a broad set of disease features.

Clinical triad: Ataxia-telangiectasia

Neurological

Early childhood ataxic-dyskinetic syndrome.

Vascular

Telangiectasias of conjunctivae and skin develop subsequently.

Immune

Immunodeficiency predisposes to recurrent respiratory infections.

Early presentation: Instability and infections

- Walking instability appears in early childhood.
- Recurrent sinopulmonary infections are common initial manifestations.
- Dysarthria and abnormal eye movements develop with disease progression.


Telangiectasias add an extracerebellar diagnostic clue

- Telangiectasias occur on the conjunctivae and skin.
- They follow the onset of the early neurologic syndrome description.
- Their presence supports a multisystem ataxia syndrome rather than an isolated motor disorder.

Immunodeficiency carries clinical consequences

- Defective immune function accompanies the neurological illness.
- Repeated sinopulmonary infections are characteristic clinical problems.
- Infectious symptoms may therefore appear alongside or before an obvious neurologic diagnosis.

ATM: The central DNA-damage response



The diagram simplifies ATM signaling; It does not specify unlisted molecular intermediates.

ATM is a kinase with broad cellular responsibilities

- ATM is encoded on chromosome 11q22–q23 in the source.
- Its kinase product plays a critical role in responses to double-stranded DNA breaks.
- The broader signaling network extends beyond canonical DNA repair.

Additional processes influenced by ATM

Cell fate and genome

Apoptosis and telomere maintenance.

Stress and energy

Mitochondrial homeostasis and oxidative-stress responses.

Protein quality control

Ubiquitin–proteasomal degradation system.

Why ATM loss causes neurodegeneration remains uncertain

- ATM influences multiple cellular stress and maintenance pathways.
- The pathway responsible for neuronal degeneration is unclear.
- Defects of single-strand-break repair in other disorders produce comparable degeneration.
- This resemblance supports a role for genome-maintenance biology but does not isolate one mechanism.

Lymphoid neoplasia is an important association

- Many individuals with ataxia-telangiectasia develop lymphoid neoplasms.
- T-cell leukemias as particularly frequent among these cancers.
- The neurodegenerative and hematologic manifestations fit a systemic DNA-response disorder.

Ataxia-telangiectasia follows a progressive course

- Early disease often features walking unsteadiness and recurrent infections.
- Speech abnormalities and eye-movement abnormalities become apparent later.
- The source describes relentless progression and death typically in early adulthood.
- The stated course should be interpreted as a general textbook description.

Compare the three major inherited patterns

Axis	Dominant SCAs	Friedreich ataxia	Ataxia-telangiectasia
Inheritance	Autosomal dominant	Autosomal recessive	Autosomal recessive
Molecular classes	PolyQ / noncoding / conventional	FXN GAA intronic expansion	ATM loss of function
Key clue	Highly variable; SCA7 vision	Neuropathy + cardiomyopathy	Telangiectasia + immunodeficiency
Mechanistic emphasis	Heterogeneous; often unresolved	Low frataxin → mitochondrial dysfunction	Deficient DNA-damage response

Three mechanisms, three biological emphases

Abnormal protein

CAG/polyglutamine SCAs:
Mutated proteins
accumulate in neuronal
nuclei.

Energy impairment

Friedreich ataxia: Frataxin
deficiency compromises
mitochondrial function.

Genome maintenance

Ataxia-telangiectasia: ATM
loss affects DNA-break
responses and other
pathways.

Mechanistic categories guide understanding; They are not interchangeable diseases.

Clinical reasoning: Onset and systemic clues

A school-aged child develops progressive gait ataxia, diminished tendon reflexes, an extensor plantar response, and impaired vibration sense. A cardiomyopathy is documented.

- Dominant polyglutamine SCA
- Friedreich ataxia
- Ataxia-telangiectasia

Which source-supported syndrome best fits?

Case interpretation: Friedreich ataxia

- Early gait ataxia followed by motor progression matches the described chronology.
- Loss of deep tendon reflexes with an extensor plantar response matches the characteristic examination.
- Impaired vibration sense and cardiomyopathy reinforce the diagnosis.
- The features are pattern-recognition clues; The source associates FRDA with FXN variants.

Clinical reasoning: Neurologic plus immune disease

A young child has progressive unsteadiness and repeated sinopulmonary infections. Later, conjunctival vascular lesions appear, with dysarthria and eye-movement abnormalities.

- Friedreich ataxia
- Ataxia-telangiectasia
- Spinocerebellar ataxia type 3

Which disorder unifies the neurological and systemic findings?

Case interpretation: Ataxia-telangiectasia

- Childhood unsteadiness and recurrent infections are characteristic early features.
- Conjunctival telangiectasias and immunodeficiency distinguish the syndrome.
- ATM dysfunction provides the disease's molecular context.
- Lymphoid neoplasia is an associated long-term concern.

Clinical reasoning: Visual symptoms in dominant ataxia

Several generations in a family have progressive ataxia. One affected individual has prominent visual impairment in addition to the movement disorder.

- SCA7 within the polyglutamine group
- Ataxia-telangiectasia
- Friedreich ataxia

Which Robbins-listed subtype offers a specific visual clue?

Case interpretation: SCA7 and pattern matching

- Autosomal-dominant family structure fits the SCA category.
- SCA7 is specifically noted for associated visual impairment.
- SCA7 appears's CAG/polyglutamine disease list.
- Genetic classification remains the appropriate framework for distinguishing numbered SCAs.

What the provided textbook supports—and leaves open

- Established: Shared neuronal targets, inherited classifications, named mutation categories.
- Established: FXN/frataxin deficiency and ATM-mediated DNA-damage signaling are central molecular lesions.
- Uncertain: Mechanisms underlying most noncoding-repeat SCAs and ATM-related neuronal degeneration.
- Limitation: Source material alone does not justify subtype-specific diagnostic thresholds or treatment algorithms.

Synthesis: recognize the disease family, then mechanism

- Spinocerebellar degeneration names a pathological-anatomical family, not one disorder.
- Dominant SCAs are genetically heterogeneous; coding and noncoding repeats must be distinguished.
- Recessive Friedreich ataxia highlights mitochondrial and cardiometabolic pathology.
- Recessive ataxia-telangiectasia links nervous-system degeneration with immune and DNA-response abnormalities.