

Squamous Metaplasia of Lactiferous Ducts

Recurrent subareolar abscess · Periductal mastitis · Zuska disease

Keratinizing ductal metaplasia → obstruction → rupture
→ inflammation

Breast pathology · mechanism, morphology and clinicopathologic reasoning

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning outcomes

- Explain keratinizing squamous metaplasia in the nipple duct and its anatomic significance.
- Trace obstruction, dilation, rupture, inflammation, fistula formation, and recurrence.
- Distinguish this process from common inflammatory mimics and outline source-supported treatment logic.

Index presentation: A recurrent subareolar lesion

Clinical vignette

- A patient presents with a painful, erythematous mass immediately beneath the areola.
- The same region has been drained previously; A tiny opening has appeared at the areolar edge.
- Which primary ductal epithelial abnormality can explain both the abscess-like mass and the fistula?

One disorder, several clinical names

Squamous metaplasia of lactiferous ducts


Recurrent subareolar abscess · periductal mastitis · Zuska disease

These names emphasize different aspects of a keratin-driven duct lesion; They are not separate Robbins entities.

The primary lesion arises in a nipple duct

Nipple / duct opening

Deeper lactiferous duct




Expected transition

The key abnormality is within the lactiferous duct near the nipple, not the terminal lobules.

Simplified duct schematic — not a histologic specimen

A clinically inflammatory lesion can begin without infection



Primary keratin leakage can produce abscess-like inflammation; Bacteria may be secondary.

Breast inflammation: Etiologic framework

Cause category	Illustrative source mechanism
Infectious	Direct bacterial invasion in lactational mastitis
Foreign-body reaction	Keratin spilled from a ruptured duct
Autoimmune / immune-mediated	Inflammatory disorders may have noninfectious causes

Inflammatory breast cancer can mimic inflammation and must be considered clinically.

Clinical identity: The central triad

Painful subareolar mass


Ductal keratinization → periductal inflammation → recurrence

A subareolar location, repeated episodes, and a possible skin fistula form a characteristic clinical pattern.

Normal anatomy: Duct opening at the nipple

Nipple / duct opening

Deeper lactiferous duct



Expected transition

The nipple duct normally transitions from a squamous surface to a glandular-lined duct.

Simplified duct schematic — not a histologic specimen

The epithelial transition is anatomically important


Region	Expected epithelial identity
Nipple surface	Squamous epithelial differentiation
Duct beyond normal transition	Nonkeratinizing glandular / ductal lining
Affected deep nipple duct	Abnormal extension of keratinizing squamous epithelium

Squamous differentiation is abnormal because it reaches well past the usual transition point.

Metaplasia changes differentiation, not location

Nipple / duct opening

Deeper lactiferous duct



Abnormally deep transition

The disease is defined by a shift toward keratinizing squamous differentiation deep within the duct.

Simplified duct schematic — not a histologic specimen

The defining morphologic finding

Keratinizing squamous epithelium


Extends well into nipple duct beyond its expected transition

This histologic distribution is the key lesion; Inflammation alone is not the defining feature.

Keratin is produced within the duct

Nipple / duct opening

Deeper lactiferous duct



Cells shed material into lumen


Metaplastic cells shed keratin into a narrow luminal pathway not designed to clear this material.

Simplified duct schematic — not a histologic specimen

Trapped keratin becomes an obstructing plug

Nipple / duct opening

Deeper lactiferous duct



Obstructing luminal plug

Retained keratin can form a plug and block the involved nipple duct.

Simplified duct schematic — not a histologic specimen

Obstruction sets up ductal dilation

**Deep squamous
metaplasia**



**Keratin
accumulation**



Luminal plug




**Progressive duct
dilation**

The duct becomes enlarged because material remains trapped proximal to the obstruction.

Dilation precedes duct rupture

Nipple / duct opening

Deeper lactiferous duct




Eventual rupture of the distended, keratin-filled duct.

Simplified duct schematic — not a histologic specimen

Rupture transfers keratin to the interstitium

Nipple / duct opening

Deeper lactiferous duct



Keratin escapes into tissue
The major inflammatory stimulus is keratin escaping into adjacent periductal tissue.

Simplified duct schematic — not a histologic specimen

Extravasated keratin acts as a foreign material

Duct wall disruption



Keratin outside duct



**Tissue reaction to
debris**



**Chronic
granulomatous
inflammation**

The granulomatous response is explained by exposure of the interstitium to normally intraluminal keratin.

Granulomatous inflammation is mechanistically linked

Extravasated keratin

Intense chronic granulomatous reaction in surrounding tissue

The inflammatory infiltrate is a consequence of duct rupture, not evidence that an infectious organism necessarily initiated disease.

Why a painful erythematous mass develops

Obstructed duct



**Rupture and local
tissue injury**



**Intense periductal
inflammation**



**Painful, red
subareolar swelling**


Clinical pain and erythema reflect the host inflammatory response to escaping keratin and tissue damage.

A localized abscess-like appearance can mislead

What the examiner sees	What may underlie it
Tender subareolar swelling	Periductal keratin-induced inflammation
Erythematous surface	Inflammation beneath the areola
Recurrent drainage	Persistent abnormal duct lining and fistula


A bacterial abscess phenotype does not establish bacteria as the initiating cause.

Chronic injury produces fibrotic traction



Fibrosis explains changes in nipple contour after recurrent disease.


Acquired nipple inversion is a mechanical sequel



Inflammation and scar contraction can pull the nipple inward.

Schematic relationships; Simplified and not drawn to scale


A subareolar fistula follows a characteristic route



In recurrent cases a tract can course beneath nipple smooth muscle toward the skin.

Schematic relationships; Simplified and not drawn to scale


The fistula may open at the areolar edge



The cutaneous opening at the edge of the areola rather than necessarily at the duct orifice.


Schematic relationships; Simplified and not drawn to scale

From duct injury to external opening



The skin opening is the external expression of a deeper ductal process.

Recurrence reflects a persisting anatomic driver



Drainage may empty a collection while the metaplastic duct epithelium persists.

The disease also occurs in males

Common in females

Also reported in males

Clinical suspicion should not be restricted exclusively to female patients.


Smoking is a prominent association

More than 90%

Of affected individuals are smokers in the Robbins account

This is a reported association from the textbook, not proof that smoking alone is sufficient to cause the lesion.

Proposed tobacco-related epithelial mechanism



This mechanism as a suggestion; Do not treat it as established causality.

Separate observation from mechanistic hypothesis


Source-supported observation	Interpretive status
Strong smoking association	Descriptive observation
Relative vitamin A deficiency	Proposed association / mechanism
Tobacco toxicity alters differentiation	Hypothesized causal pathway
Keratin plugs and duct rupture	Described pathologic sequence

Association, hypothesis, and direct morphology carry different levels of evidence.

Conceptual anatomy of a keratin-plugged duct

Nipple / duct opening

Deeper lactiferous duct




Obstructing luminal plug

This is an original vector schematic; No microscopic specimen is depicted.

Simplified duct schematic — not a histologic specimen

Morphology is a sequence, not one static lesion



Different tissue samples can capture different stages of the same pathophysiologic process.

Recognize the primary epithelial abnormality


Core microscopic observation	Meaning
Keratinizing squamous epithelium	Abnormal duct differentiation
Extension into deep nipple duct	Beyond the usual squamous transition
Accumulated keratin in lumen	Potential obstructing substrate

The original textbook diagram is not reproduced; These are source-described features.

Identify the material that plugs the lumen

Nipple / duct opening

Deeper lactiferous duct



Cells shed material into lumen


The retained luminal material is desquamated keratin—not the thick secretions defining duct ectasia.

Simplified duct schematic — not a histologic specimen

Rupture creates a periductal reaction

Nipple / duct opening

Deeper lactiferous duct




Retained keratin

Keratin escapes into tissue

A ruptured keratin-containing duct links luminal pathology to tissue-level inflammation.


Simplified duct schematic — not a histologic specimen

Inflammatory morphology depends on chronology



Chronic foreign-body-type inflammation as primary and anaerobic superinfection in some recurrent lesions.

Secondary anaerobic infection can supervene



Not all abscess-like lesions begin as bacterial infections; Bacteria become relevant when secondary infection develops.

Primary versus secondary events

Primary disease process	Possible secondary consequence
Deep keratinizing metaplasia	Duct blockage and dilation
Duct rupture	Chronic granulomatous response
Persistent duct abnormality	Recurrences and fistula
Recurrent diseased area	Secondary anaerobic infection

Identifying the initiating lesion changes management logic.

Clinical presentation is anatomically focused

- Painful, erythematous mass typically centered beneath the nipple–areolar complex.
- Recurrent episodes can culminate in a fistulous opening at the areolar margin.
- Inflammatory scarring may pull the nipple inward and cause acquired inversion.

Initial and recurrent disease are not identical


First recognized episode	Recurrent episodes
Tender, red subareolar mass	Repeat inflammation in same region
May resemble a simple abscess	Fistula is more likely to emerge
Underlying duct disease may be occult	Fibrotic traction / nipple inversion may develop

Recurrence raises the importance of a persisting ductal lesion.

The broader differential includes lactational mastitis

- Lactational acute mastitis typically occurs during the first month of breastfeeding.
- Nipple cracks and fissures permit bacterial entry into breast tissue.
- *Staphylococcus aureus* is the usual organism; Streptococci are less frequent.

Why lactational mastitis becomes inflamed



This infectious portal-of-entry mechanism differs from metaplasia-driven keratin retention.

Compare mechanisms: Duct metaplasia vs mastitis

Feature	Squamous metaplasia	Lactational acute mastitis
Initiating event	Keratinizing ductal differentiation	Bacterial entry via nipple injury
Typical setting	Recurrent subareolar lesion	Early breastfeeding
Inflammatory driver	Keratin extravasation	Bacterial infection
Possible extension	Fistula and scarring	Abscess or spreading cellulitis

Features reflect Robbins descriptions, not a stand-alone diagnostic rule.

Duct ectasia is another periareolar mimic

- Often presents as a palpable periareolar mass with thick white nipple secretions.
- Pain and erythema are uncommon in the Robbins description.
- More typical in multiparous females in the fifth or sixth decade.

Duct ectasia has a different luminal substrate

Squamous metaplasia	Duct ectasia
Keratinizing epithelium within nipple duct	Ectatic duct with inspissated secretions
Obstructing keratin plug	Secretions and lipid-laden macrophages
Keratin rupture → granulomatous response	Secretions / cholesterol → chronic inflammation

Both may produce fibrosis and retraction, but their mechanisms differ.

Smoking helps separate these duct disorders

Feature	Squamous metaplasia	Duct ectasia
Cigarette-smoking association	Strong (>90% smokers)	Not associated with smoking
Frequent major symptom	Painful red subareolar mass	Mass with thick white discharge
Characteristic process	Deep squamous keratinization	Dilated secretion-filled ducts

Smoking history is supportive context, not a substitute for clinicopathologic assessment.

A swollen erythematous breast is not always benign

Inflammatory breast carcinoma

An important clinical mimic of inflammatory disorders

Robbins specifically warns that inflammatory breast cancer should be considered in females with an erythematous, swollen breast.

Cancer exclusion is an independent diagnostic duty

**Inflammatory breast
presentation**



Clinical assessment



**Appropriate breast
imaging**



**Tissue sampling
when indicated**

Chapter context supports combined clinical, imaging, and biopsy evaluation of symptomatic breast disease; A benign-looking story does not eliminate cancer.

Clinical contrasts: Avoid single-feature diagnosis

Entity	Clinical discriminator in source
Squamous duct metaplasia	Recurrent painful subareolar lesion ± fistula
Acute lactational mastitis	Breastfeeding-associated infection, often fever
Duct ectasia	White secretion; Pain / erythema uncommon
Inflammatory breast cancer	Can mimic breast inflammation; Must be considered

This is a teaching comparison, not a formally validated diagnostic scoring system.

Why simple incision and drainage may fail

**Incise and drain
collection**



**Immediate cavity
decompression**



**Keratinizing lining
remains**



**New obstruction and
recurrence possible**

Robbins explicitly identifies persistent metaplastic epithelium as the reason recurrence is common after simple drainage.


Antibiotics treat a secondary component

Secondary bacterial infection

Antibiotics have a therapeutic role when it is present

Antimicrobial therapy does not itself remove the keratin-producing epithelium.

Definitive surgery targets the disease-bearing anatomy



En bloc removal of the affected duct and contiguous fistula tract is curative in most cases.

Schematic relationships; Simplified and not drawn to scale

Why duct-and-fistula excision is logical

**Persistent abnormal
duct epithelium**



**Remove involved
duct**



**Remove connected
fistula tract**



**Break the
recurrence pathway**

Eliminating both the keratin source and contiguous tract addresses the cause rather than only draining its consequence.

Management decisions follow pathogenesis

Recurrent subareolar inflammatory lesion

Identify the characteristic site and history

Assess for cancer / other mimics

Clinical assessment, imaging, biopsy as indicated

Determine if superinfection is present

Antibiotics are relevant when bacterial infection coexists

Address persistent duct lesion

En bloc duct and fistula excision is usually curative

This is a conceptual teaching pathway derived, not a formal clinical guideline.

Case exercise: A first subareolar episode

Clinical vignette

- A patient presents with a painful, red mass directly beneath one nipple.
- The initial impression is bacterial abscess, but there is no evidence here establishing primary infection.
- Name the epithelial change and trace the two steps that produce tissue inflammation.

Case reasoning: First episode

**Keratinizing
squamous
metaplasia**



**Keratin plugs the
duct**



**Duct dilates and
ruptures**



**Keratin provokes
inflammation**


The chain accounts for a painful abscess-like mass without requiring primary bacterial infection.

Case exercise: Recurrent lesion with fistula

Clinical vignette

- The same patient returns after repeated incision and drainage of a subareolar collection.
- A tract now opens at the areolar edge and the nipple is retracted.
- Explain why drainage alone is insufficient and which anatomic structures definitive surgery should address.

Case reasoning: Prevent another recurrence



Nipple inversion reflects traction by scarring; Recurrence reflects the underlying metaplastic duct.

Microscopy challenge: What establishes the lesion?

Most discriminating pathologic observation?

A. Granulomatous inflammation somewhere in the breast

B. Keratinizing squamous metaplasia extending deeply into the nipple duct

C. An abscess cavity without an identifiable ductal abnormality

Discussion / instructor answer is in speaker notes.

Mechanism challenge: Order the events

Which sequence is source-consistent?

A. Bacterial infection → squamous metaplasia → keratinization in every case

B. Deep squamous metaplasia → keratin plug → rupture → granulomatous inflammation

C. Duct ectasia → vitamin A excess → fistula

Discussion / instructor answer is in speaker notes.

Differential challenge: Identify duct ectasia

Which feature favors duct ectasia rather than Zuska disease?


A. Thick white nipple secretions with pain and erythema uncommon

B. Recurrent areolar-edge fistula after keratin duct obstruction

C. Strong smoking association with deep ductal squamous keratinization

Discussion / instructor answer is in speaker notes.

Synthesis: The four indispensable links



Clinically: Recurrent painful subareolar mass; Sometimes fistula and nipple inversion. Definitive management addresses the ductal source.

Reference and source limitations

- Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. Elsevier; 2026.
- Primary teaching section: Squamous Metaplasia of Lactiferous Ducts; related context: Inflammatory Disorders, Acute Mastitis, and Duct Ectasia.
- Original textbook image was not available for authorized insertion. All anatomy and mechanism illustrations are original editable schematics, not pathology specimens.
- Original chapter PDF was unavailable for image-level verification; extracted Markdown and topic-unit source were reviewed.