

Thyroglossal Duct Cyst

Embryologic origin · morphology · clinicopathologic diagnosis

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

What this lesion teaches

- Trace normal thyroid descent and persistence of the embryonic duct.
- Predict location and epithelial lining from developmental anatomy.
- Identify characteristic cyst-wall histology and reactive changes.
- Distinguish clinically important mimics and explain definitive excision.

A developmental remnant, not a thyroid nodule

- A thyroglossal duct cyst develops from persistent segments of the thyroid migration tract.
- It is the most common clinically significant congenital thyroid anomaly.
- The cyst is usually in the midline anterior neck, often near the hyoid bone.
- Its wall may contain thyroid follicles, but the lesion is not simply a dilated follicle.

DEFINING RELATIONSHIP

Persistent epithelial tract → cystic cavity

Where it belongs among neck lesions

Congenital midline

Thyroglossal duct cyst

Ectopic thyroid

Dermoid / epidermoid

Other congenital

Branchial cleft cyst

Lymphatic malformation


Acquired / neoplastic

Inflammatory node

Thyroid nodule


Cystic metastatic node

Thyroid primordium: The starting point




- The thyroid anlage originates at the foramen cecum of the tongue.
- The primordial gland descends through the anterior neck.
- The final thyroid position is anterior to the upper trachea.

The normal path of thyroid descent



- Foramen cecum: Superior embryonic origin.
- Hyoid region: Key intermediate landmark.
- Pretracheal region: Definitive gland position.

The hyoid is an anatomical anchor



- The duct courses in intimate relation to the hyoid bone.
- Most clinical cysts are found near this level.
- This association explains both imaging landmarks and the surgical field.

Normal involution versus retained tract

Usual developmental outcome

Epithelial descent tract



Duct normally regresses



No persistent cystic remnant

Persistent duct remnant

Epithelial descent tract



Duct segment persists



Secretions accumulate → cyst


A segment—not necessarily the entire duct—may persist

- Embryonic tract obliteration may be incomplete and discontinuous.
- Residual epithelial segments can form one or more cystic spaces.
- A persistent tract or sinus can coexist with the cyst.

ANATOMIC CONSEQUENCE

A seemingly discrete cyst may have a deeper ductal extension.

Why retained epithelium becomes a cyst



Each step explains the clinicopathologic consequence of duct persistence.


Congenital origin does not require infant presentation

- The developmental abnormality is congenital.
- Small persistent remnants may initially be clinically occult.
- Progressive secretion or inflammation can reveal the lesion later in life.

DIAGNOSTIC CAUTION

An adult presentation
does not exclude a
congenital duct
anomaly.

One embryologic tract; Several anatomical levels



- Lingual and suprahyoid portions may form high lesions.
- Perihyoid cysts are closely related to the hyoid.
- Infrahyoid cysts descend into the anterior cervical soft tissues.

Midline is typical—but not an absolute rule

- The embryologic pathway is predominantly central.
- Infrahyoid cysts may be slightly paramedian.
- Location alone should not override evaluation of the hyoid relationship and internal architecture.

IMAGING PRINCIPLE

Hyoid proximity + cystic morphology is more informative than midline position alone.

How large is a usual cyst?

1–4 cm


Head and Neck chapter

Usually $\leq$ 2–3 cm

Endocrine System chapter

These statements describe common observed dimensions, not diagnostic cutoffs.


Macroscopic configuration



- Lesions are often rounded, but fusiform enlargement may follow the tract.
- A circumscribed cavity forms in the anterior cervical soft tissues.
- Fluid, mucus, or debris may occupy the lumen.

Schematic cut-surface; Does not depict an observed gross specimen.

Cyst structure: Lumen, epithelium, fibrous wall




- Luminal cavity contains secretions and desquamated material.
- Epithelial lining separates contents from the fibrous stroma.
- The wall may contain inflammation and thyroid remnants.

Original conceptual histology schematic — not a patient specimen.

Multiple epithelial phenotypes are expected


Lining phenotype	Typical description	Teaching implication
Squamous	Stratified epithelium	More characteristic near tongue base
Respiratory-type	Pseudostratified columnar	Often in lower tract remnants
Mixed / transitional	Coexisting patterns	A mixed lining remains compatible with TGDC

Observed epithelial linings in a 685-case series



Observed descriptive frequencies in the Thompson et al. Series; Not diagnostic thresholds.


Higher lesions: Squamous lining



- Stratified squamous epithelium is described near the tongue base.
- The lining is one legitimate variant of thyroglossal duct differentiation.
- Squamous lining alone is not evidence of branchial cleft origin.

Original conceptual histology schematic — not a patient specimen.


Lower lesions: Respiratory-type lining



- Pseudostratified columnar epithelium is described in lower locations.
- Ciliated respiratory-type differentiation may be recognized.
- This lining is not specific without location and wall findings.

Original conceptual histology schematic — not a patient specimen.


Mixed lining reflects epithelial plasticity



- Squamous and respiratory-type regions may coexist.
- Transitional epithelial differentiation is also described.
- Sampling a single lining focus may underestimate heterogeneity.

Original conceptual histology schematic — not a patient specimen.


The fibrous wall is diagnostic territory



- Collagen-rich fibrous stroma supports the cavity.
- Inflammatory aggregates may accompany the lining.
- Embedded thyroid follicles can support embryologic origin.

Original conceptual histology schematic — not a patient specimen.

Thyroid follicles may occur in the wall



- Clusters of colloid-containing follicles may persist along the migratory route.
- They can be minute and discontinuous.
- Their presence is supportive, but absence in a sample does not exclude a duct remnant.

Original conceptual histology schematic — not a patient specimen.


Lymphoid aggregates: Present but not defining

- Lymphoid aggregates in the fibrous cyst wall.
- Chronic inflammatory change may be patchy.
- Prominent nodular lymphoid tissue can instead favor branchial cleft cyst when the lesion is lateral.

INTERPRETATION

Integrate wall composition with the location of the lesion.

Superimposed infection can alter the lesion



Each step explains the clinicopathologic consequence of duct persistence.


Why an inflamed cyst can lack recognizable lining

- Repeated inflammation can erode or denude epithelium.
- Fibroinflammatory repair may replace the original cyst lining.
- Search remaining sampled wall for glandular elements, follicles, and ductal epithelium.

PITFALL


No lining in one section
is not sufficient to reject
TGDC.

Rupture produces a secondary tissue reaction



Each step explains the clinicopathologic consequence of duct persistence.

Specimen sampling can decide the diagnosis




- Document relationship to hyoid bone, tract, and adjacent soft tissues.
- Sample the lining and thickened or irregular areas.
- Look for thyroid follicles and investigate any mural nodule thoroughly.

Schematic of specimen composition; Not a gross pathology image.

A practical benign histology checklist

- Epithelial type(s) and preservation of lining
- Fibrous wall and lymphoid / inflammatory changes
- Thyroid follicles or other tract remnants
- Rupture, abscess, sinus, or fistula
- Presence or absence of atypical proliferations


Classic clinical presentation



- Slow-growing, often painless anterior cervical swelling.
- Usually near the midline and hyoid region.
- May become apparent in childhood or adult life.

Site schematic for orientation, not a clinical photograph.

A mass that moves with tongue protrusion




Swallowing movement is supportive, not definitive

- Swallowing may elevate the lesion.
- Thyroid-related and other neck structures also move with swallowing.
- The combined pattern of location, tongue movement, and ultrasound provides stronger evidence.

BEDSIDE EXAM

No single dynamic sign is a pathology diagnosis.


Acute infection changes the clinical picture



- Rapid enlargement, tenderness, erythema, and fever may develop.
- Inflamed duct remnants may become abscesses.
- Treat acute infection before planned definitive removal.

Site schematic for orientation, not a clinical photograph.

Sinus and fistula formation



- A patent epithelial remnant can form a sinus tract.
- Spontaneous drainage or previous infection may produce a skin opening.
- The full tract is relevant during definitive surgery.

Case: Intermittent discharge in a child

CLINICAL / PATHOLOGY SCENARIO


A child has a recurrent midline swelling inferior to the hyoid. The overlying skin periodically drains mucoid fluid.

Which embryologic structure explains both the cyst and a draining sinus?

REASONED ANSWER

A persistent segment of the thyroglossal duct; Evaluate the tract rather than treating only a surface opening.

Initial decision: Is the mass truly cystic?




Ultrasound is usually the first imaging study

- Assesses cystic versus solid architecture without ionizing radiation.
- Identifies location relative to hyoid, strap muscles, and thyroid.
- Can document the presence of orthotopic thyroid tissue.

PREOPERATIVE REQUIREMENT

Identify normal thyroid tissue before excising a suspected duct remnant.


Typical sonographic pattern



- Well-circumscribed cystic lesion with posterior enhancement.
- Anechoic to hypoechoic content; Internal debris is possible.
- No intrinsic solid vascular component in an uncomplicated cyst.

Synthetic schematic (native shapes), not diagnostic patient ultrasound.


Infection can complicate the ultrasound appearance



- Internal echoes and thicker walls may accompany inflammation.
- Complex fluid does not by itself imply carcinoma.
- Clinical tenderness, fever, and perilesional change aid interpretation.

Synthetic schematic (native shapes), not diagnostic patient ultrasound.

Mural nodules warrant deliberate evaluation



- A true solid intracystic component is not a typical simple cyst feature.
- Enhancing mural nodules or calcifications may signal neoplasia.
- Target the suspicious region and evaluate the thyroid and cervical nodes.

Synthetic schematic (native shapes), not diagnostic patient ultrasound.

When CT or MRI changes management

CT

- Can show relationship to the hyoid and deeper neck anatomy.
- Useful for calcifications and complicated collections.

MRI

- Defines soft-tissue extent and tongue-base involvement.
- Useful when anatomy is complex or lesion is atypical.

Verify the orthotopic thyroid gland

Key safety check

- Is a normal thyroid gland seen in its expected anatomical position?
- Is the lesion genuinely cystic?



Interpretive actions

- Ectopic thyroid tissue can be the only functioning thyroid in rare patients.
- Inspect the normal thyroid bed before removing a presumed duct cyst.
- A normal biochemical result does not alone prove normal thyroid location.

Lingual thyroid is a critical embryologic mimic

Thyroglossal duct cyst

- Fluid-filled persistent duct remnant
- Variable epithelial lining
- May contain scattered thyroid follicles

Lingual thyroid

- Ectopic functioning thyroid parenchyma
- Usually predominantly solid thyroid tissue
- Normal thyroid may be absent from its usual site

Branchial cleft cyst: Lateral versus midline

Thyroglossal duct cyst

- Near midline / hyoid
- Embryonic thyroid descent tract
- Thyroid follicles may be present

Branchial cleft cyst

- Usually lateral upper neck / SCM
- Branchial developmental remnant
- Prominent organized lymphoid tissue common

Dermoid and epidermoid cysts: Different lining, different contents

Thyroglossal duct cyst

- Secretory respiratory or squamous lining
- May contain thyroid follicles
- Often hyoid-associated

Dermoid / epidermoid

- Keratinizing squamous lining with lamellated keratin
- Dermoid: Cutaneous adnexal structures
- Typically superficial to strap musculature

Lymphatic malformation: Pattern recognition

Thyroglossal duct cyst

- Focal duct-path lesion
- Often near hyoid
- Single or clustered cystic spaces

Lymphatic malformation

- Often crosses tissue planes
- May be multilocular / trans-spatial
- Fluid–fluid levels can occur

An adult cystic neck mass needs oncologic caution

- Cystic lymph-node metastasis may mimic a benign congenital cyst.
- An irregular solid component, abnormal nodes, or unusual location changes the differential.
- Clinical age and risk profile should modify the work-up.

RED FLAG

Do not label every adult cystic neck mass “branchial” or “thyroglossal” without assessment.

Differential diagnosis by anatomical signature

Entity	Dominant anatomical clue	Additional discriminator
TGDC	Midline / perihyoid	Tongue movement, duct trajectory
Branchial cleft	Lateral, SCM margin	Lymphoid wall; Lateral upper neck
Dermoid / epidermoid	Superficial central neck	Keratin contents; Adnexa in dermoid
Lymphatic malformation	Trans-spatial	Multiloculated spaces
Cystic nodal metastasis	Nodal chain	Suspicious solid/nodal components


Fine-needle aspiration: Informative but limited

- Aspiration of cyst fluid may yield few diagnostic epithelial cells.
- Inflammatory cells and macrophages can dominate cytology.
- Sampling a suspicious solid nodule is more informative than fluid alone.

DIAGNOSTIC LIMITATION

A nondiagnostic or benign-appearing aspirate cannot exclude a small mural carcinoma.

Sistrunk excision treats the tract, not only the cyst



- Remove the cyst and involved tract.
- Excise the central portion of the hyoid bone.
- Remove a core of superior tract tissue toward the tongue base.

Why simple cystectomy has more recurrences

Cystectomy alone

Cyst excised alone



Residual epithelial tract remains



New secretion / cyst regeneration

Sistrunk excision

Cyst + central hyoid + tract




More complete remnant removal



Lower recurrence risk

What arrives in a Sistrunk specimen?



- Cystic lesion with adjacent fibrous and sometimes skeletal-muscle tissue.
- A central segment of hyoid bone is usually part of the specimen.
- The duct remnant may be inconspicuous; Orient and submit suspicious foci.

Schematic of specimen composition; Not a gross pathology image.

After surgery: What recurrence means

- Recurrent disease can reflect retained epithelial extensions.
- Prior rupture or infection may complicate tract delineation.
- Review prior procedure and histology when recurrent midline swelling appears.

CLINICOPATHOLOGIC LINK

Incomplete remnant excision is a plausible cause of recurrent disease.


Carcinoma is uncommon but not ignorable

- Most thyroglossal duct cysts are benign.
- Carcinoma associated with a cyst is distinctly rare.
- Atypical mural tissue should be sampled and reported carefully.

SOURCE NUANCE

Exceedingly rare transformation of lining epithelium; Thyroid-type carcinoma in embedded follicles is a separate consideration.

Most reported associated carcinomas are papillary type



- Neoplasm may arise in ectopic thyroid follicles within a duct remnant.
- Papillary thyroid carcinoma is the major reported carcinoma subtype.
- The conventional cyst lining may remain uninvolved.

Original conceptual histology schematic — not a patient specimen.

Histologic red flags in the cyst wall

- True papillary or infiltrative epithelial proliferation
- PTC-type nuclear enlargement / irregularity / grooves
- Nuclear clearing or intranuclear pseudoinclusions
- Invasive growth beyond a benign cyst-wall contour
- Correlation with suspicious gross or imaging nodules

Primary duct carcinoma versus thyroid primary

Duct-associated primary

- Carcinoma demonstrably centered in duct remnant
- Benign duct lining and follicles can support origin
- Requires clinical-imaging correlation

Separate thyroid carcinoma

- Thyroid gland can contain synchronous PTC
- Metastatic disease or dual primaries may be considered
- Examine orthotopic thyroid and cervical nodes

The malignancy work-up must stay proportional

Key safety check

- Is a normal thyroid gland seen in its expected anatomical position?
- Is the lesion genuinely cystic?



Interpretive actions

- Verify the lesion, orthotopic thyroid, and cervical lymph nodes.
- Discuss carcinoma cases with pathology, radiology, and head-and-neck teams.
- Additional thyroid surgery is individualized; No single approach fits every case.

Case: A mural nodule in a neck cyst

CLINICAL / PATHOLOGY SCENARIO

A 46-year-old adult has a perihyoid cyst on ultrasound. A vascular solid nodule projects into the lumen.

Does the benign appearance of most of the cyst exclude neoplasia?

REASONED ANSWER

No. Characterize and sample the solid component; Inspect the thyroid and cervical nodes.

Integrated pathology case: Lining partly absent

CLINICAL / PATHOLOGY SCENARIO

An excised midline hyoid-associated cyst is predominantly fibroinflammatory; Small residual ducts and thyroid follicles are found in deeper levels.

Does partial denudation undermine a thyroglossal duct diagnosis?

REASONED ANSWER

No. Inflammation may erase lining; The lesion site and residual ductal/thyroid elements support TGDC.

Clinical reasoning: Distinguish two neck cysts

CLINICAL / PATHOLOGY SCENARIO

Patient A: Child, central perihyoid mobile lesion. Patient B: Young adult, painless upper lateral neck cyst along the sternocleidomastoid.

Which developmental diagnosis is favored for each?

REASONED ANSWER

A: Thyroglossal duct cyst. B: Branchial cleft cyst. Histology and imaging resolve atypical cases.

Reporting exercise: Avoid unsupported certainty

CLINICAL / PATHOLOGY SCENARIO


A cyst fragment shows nonkeratinizing squamous lining and lymphocytes; The specimen site is not documented and no thyroid follicles are seen.

Can the epithelial pattern alone establish a thyroglossal duct cyst?

REASONED ANSWER

No. Squamous lining overlaps with branchial cleft cyst; Anatomical context and deeper sampling matter.

The integrated diagnostic pathway



Four anchors for retention

01 Embryology: Persistent tract from the foramen cecum.

02 Anatomy: Midline or paramidline cyst related to the hyoid.

03 Histology: Squamous / respiratory lining $\pm$ thyroid follicles.

04 Management: Sistrunk excision; Scrutinize unusual mural tissue.

Sources and interpretation boundaries

1. Robbins, Cotran & Kumar. Pathologic Basis of Disease. 11th ed. 2026; ch. 16:677–678; ch. 24:993.
2. Patel S, Bhatt AA. Thyroglossal duct pathology and mimics. Insights Imaging. 2019;10:12. doi:10.1186/s13244-019-0694-x.
3. Thompson LDR, Herrera HB, Lau SK. Head Neck Pathol. 2016;10:465–474. doi:10.1007/s12105-016-0724-7.
4. Thimsen V, et al. Virchows Arch. 2025;486:1139–1151. doi:10.1007/s00428-025-04125-2.
5. Thyroglossal Duct Cyst. NCBI Bookshelf. NBK519057.