

Vascular Dementia

Cerebrovascular mechanisms, brain pathology and clinical consequences

Dr Abu Suraih Sakhri

MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Define vascular dementia in its neuropathologic and clinical context.
- Relate vascular lesions and their distribution to cognitive and neurologic impairment.
- Distinguish large-vessel infarction, lacunar injury and diffuse subcortical white-matter disease.
- Integrate coexisting vascular and degenerative changes without assuming a single cause.

Definition: Cumulative vascular brain injury

A syndrome arising from multifocal vascular disease.


- A syndrome of dementia developing with multifocal cerebrovascular damage over months to years.
- Multiple bilateral infarcts may affect both gray and white matter.
- The hallmark is an anatomic burden of vascular injury, not one specific histologic lesion.
- Clinical findings commonly extend beyond isolated memory impairment.

Why the syndrome is clinically distinctive

- Dementia coexists with gait abnormalities and pseudobulbar signs.
- Superimposed focal neurologic deficits may be present.
- The vascular injuries may arise through several mechanisms.
- Multiple anatomic systems may be impaired by bilateral lesions.

Bilateral multifocal lesions

Conceptual distribution map




Outer rim = cortex | Inner field = deep tissue

- Infarcts occur in both cerebral hemispheres.
- Gray and white matter can be involved together.
- Distinct lesion sites provide an anatomic basis for multi-domain impairment.

Schematic — not an MRI, gross specimen, or mapped anatomy

From vascular disease to cognitive impairment



- The syndrome generally develops across months and years, rather than from a single acute event.
- Consequences depend on the amount and location of injured brain tissue.

Clinical stroke versus vascular dementia

Stroke (episode)

- Acute neurologic manifestations explained by a vascular event.
- The focal deficit depends on the vascular territory affected.
- An episode may contribute to later accumulated injury.

Vascular dementia (syndrome)

- Cognitive syndrome associated with multiple or widespread vascular lesions.
- Gait and pseudobulbar abnormalities may coexist.
- May have superimposed focal deficits.

Related concepts, not interchangeable clinical labels

A syndrome with more than one vascular cause

- Cerebral atherosclerosis is a major etiologic substrate.
- Thrombosis or embolization from carotid vessels or the heart may produce infarcts.
- Chronic hypertension causes cerebral arteriolosclerosis and small-vessel injury.
- Different mechanisms can coexist in the same brain.

Three principal vascular pathways




- Three causative vascular categories; The fourth box represents their shared tissue outcome.

Schematic relationships (not quantitative)

The key determinant is lesion anatomy

- Cerebral infarction produces deficits related to its anatomic distribution.
- Infarct size and location reflect duration of ischemia and adequacy of collateral perfusion.
- Small injuries in strategically important tracts can cause prominent functional deficits.
- Do not infer syndrome severity solely from counting visible lesions.

Atherosclerosis as an upstream substrate




- Large-artery atherosclerosis can compromise blood supply directly or supply thromboemboli.
- The carotid bifurcation and origin of the middle cerebral artery are important sites.

Atherosclerotic occlusion: Common sites

- The carotid bifurcation is a common site of thrombotic occlusion.
- The origin of the middle cerebral artery is another frequent location.
- Either end of the basilar artery may also be involved.
- Atherosclerotic cerebrovascular disease often coexists with hypertension or diabetes.

Carotid-to-cerebral thromboembolism




- Thrombi can fragment and move distally from their site of formation.
- Occlusion of cerebral branches may generate distinct territories of ischemic necrosis.

Schematic causal sequence

Cardiac sources of cerebral emboli

- Cardiac mural thrombi are important embolic sources.
- Myocardial infarction, valvular disease and atrial fibrillation are relevant predispositions.
- Emboli can lodge at branch points or previously narrowed segments.
- Recurrent embolization can create anatomically separated infarcts.

Mechanism of cardioembolic cerebral injury



- The middle cerebral artery territory is frequently affected by cerebral embolism.
- Clinical outcome depends on the distribution of damaged tissue.

Embolism versus in-situ thrombosis

Embolic occlusion

- Occlusive material arrives from a distant source.
- Cardiac or carotid origin is common.
- Vessel branching points are frequent lodging sites.


Local thrombotic occlusion

- Thrombus arises at a diseased intracranial or extracranial artery.
- Acute change in a vulnerable plaque may initiate thrombosis.
- Extension and distal fragmentation may also occur.

Blood flow and cerebral vulnerability

- The brain has substantial continuous oxygen and energy requirements.
- Cerebral autoregulation helps maintain relatively constant perfusion.
- Reduced oxygen delivery may follow low flow, impaired oxygenation or both.
- Collateral supply, insult duration and flow reduction shape tissue survival.

What determines infarct extent?



- Magnitude and rapidity of the flow reduction also modify injury.
- These determinants influence lesion shape, anatomic site and clinical manifestations.

Simplified determinant chain, not a numerical risk model

Cortical versus deep arterial supply

Cortical arterial territories

- Additional leptomeningeal anastomoses may contribute collateral supply.
- Larger territory infarcts may follow occlusion of major branches.
- Cortical involvement can accompany cognitive deficits.

Deep penetrating territories


- Deep penetrating arteries have little or no collateral supply.
- They supply basal ganglia, thalamus and deep white matter.
- Small-vessel occlusion can create lacunar infarcts.

Hypertension and the cerebral microvasculature


- Chronic hypertension affects penetrating arteries and arterioles.
- Wall thickening and hyaline arteriolosclerosis alter vessel function.
- Affected vessels supply deep gray nuclei, white matter and brainstem.
- Occlusion may produce lacunar infarcts; Rupture may cause hemorrhage.

Hyaline arteriolosclerosis reduces the lumen

Cross-sectional vessel schematic



Comparator



Affected vessel

- Vessel walls become thickened by hyaline change.
- Lumen narrowing predisposes to distal ischemic injury.
- Wall rigidity is associated with vulnerability to rupture.
- This pattern is prominent in deep brain structures.

Schematic — lumen and wall not drawn to scale


Small-vessel occlusive pathway



- The causal sequence contrasts with embolic occlusion of a larger arterial branch.
- Repeated small infarcts can accumulate in subcortical structures.

Lacunar infarcts: Deep distribution

Conceptual distribution map



Outer rim = cortex | Inner field = deep tissue

- Lacunes affect deep gray nuclei, deep white matter and pons.
- Illustrated locations are conceptual, not a diagnostic map.
- Cumulative bilateral small lesions contribute to vascular impairment.

Schematic — not an MRI, gross specimen, or mapped anatomy

Lacunae: Gross and microscopic identity

- Lacunar infarcts as small occlusive infarcts below 15 mm in maximal dimension.
- Grossly, healed lesions form small lake-like spaces.
- Microscopically, there is tissue loss with surrounding gliosis.
- This definition characterizes a lesion, not by itself the clinical diagnosis of dementia.

Common lacunar locations

Anatomic site	Pathology context	Clinical teaching interpretation
Putamen / globus pallidus	Deep perforator territory	Potential motor-network disturbance
Thalamus	Deep gray-matter infarct	Disruption of distributed brain circuits
Internal capsule	Compact white-matter tracts	Focal neurologic deficits possible
Deep white matter	Subcortical ischemic injury	Disconnection of long association pathways
Caudate nucleus / pons	Additional lacunar locations	Heterogeneous focal signs

Specific clinical outcomes depend on precise lesion location; Table is explanatory, not a fixed rule.

Large territorial infarcts versus lacunes

Territorial infarction

- Follows occlusion of a major arterial supply.
- Often has wedge- or territory-related distribution.
- Can involve cortical gray matter and subjacent structures.


Lacunar infarction

- Small, deep occlusive lesion.
- Often reflects disease of penetrating arterioles.
- Remote lesion shows cavity and surrounding gliosis.

Small-vessel disease has two major outcomes

- Arteriolar occlusion can produce a lacunar infarct.
- An abnormal rigid vessel wall can rupture into brain parenchyma.
- Both are consequences of vascular pathology in deep cerebral territories.
- The occurrence of either lesion should prompt attention to the underlying vasculopathy.

Hypertensive injury: Occlusion or rupture




- The tissue phenotype reflects the mode of vessel failure.
- Lacunar infarcts and hypertensive intraparenchymal hemorrhages have related vascular substrates.

Cerebral amyloid angiopathy: Distinct vessel injury


- Amyloid- β peptide accumulates within meningeal and cortical vessel walls.
- The altered walls become rigid and vulnerable to bleeding.
- Lobar hemorrhages and small microbleeds are characteristic manifestations.
- This distribution contrasts with hypertensive deep arteriolosclerosis.

Amyloid deposition in cerebral vessel walls

Cross-sectional vessel schematic



Comparator



Affected vessel

- The deposits are amyloid- β , not the collagen of hyaline arteriolosclerosis.
- Cortical and leptomeningeal vessels are preferentially involved.
- Cerebral amyloid angiopathy is an important alternative vascular substrate.

Schematic — lumen and wall not drawn to scale


Two small-vessel disease patterns

Feature	Hypertensive arteriolosclerosis	Cerebral amyloid angiopathy
Typical vascular territory	Deep penetrating arterioles	Cortical / leptomeningeal vessels
Dominant wall alteration	Hyaline thickening	Amyloid- β deposition
Infarct/bleeding pattern	Lacunes and deep hemorrhage	Lobar bleeding and microbleeds
Major contextual association	Long-standing hypertension	Amyloidogenic peptide accumulation

CADASIL: Hereditary small-vessel injury

- Cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy.
- Caused by pathogenic NOTCH3 variants affecting its extracellular receptor domain.
- Recurrent small-vessel strokes and progressive white-matter changes may be accompanied by dementia.
- An inherited vascular disorder is a distinctive alternative to common hypertensive injury.

CADASIL: Molecular lesion to syndrome



- The Robbins description links this genetic arteriopathy to recurrent small-vessel events and dementia.
- The pathway is a teaching simplification, not a complete molecular model.

Schematic causal sequence

Other hereditary cerebral microangiopathy

- Variants affecting COL4A1, a vascular basement-membrane component, cause inherited small-vessel disease.
- Brain hemorrhage may coexist with cataract, hematuria or epilepsy.
- The multisystem phenotype helps distinguish this disorder from sporadic hypertensive vasculopathy.
- Not every hereditary cerebral vasculopathy necessarily causes vascular dementia.

Acquired and inherited small-vessel mechanisms

Common acquired pathway


- Long-standing hypertension.
- Hyaline arteriolar wall alteration.
- Deep lacunar infarcts or hemorrhage.

Hereditary pathway

- CADASIL: NOTCH3-associated vasculopathy.
- COL4A1-associated basement membrane disease.
- Clinical and family context may aid recognition.

Gray-matter and white-matter injury

Conceptual distribution map



Outer rim = cortex | Inner field = deep tissue

- Cumulative vascular damage may affect cortical and deep gray matter.
- White-matter infarcts can interrupt cerebral connections.
- The syndrome cannot be reduced to a single microscopic pattern.

Schematic — not an MRI, gross specimen, or mapped anatomy

Cortical and subcortical injury patterns

Cortical gray matter

- May sustain infarction in an arterial territory.
- Focal neuronal loss and subsequent gliosis reflect chronic injury.
- Can coexist with subcortical infarcts.

Subcortical structures

- Deep nuclei and connecting white matter may be damaged.
- Penetrating vessel disease is an important substrate.
- Extensive white-matter involvement defines Binswanger pattern.


Binswanger disease: Defining feature

The defining distinction is topographic: Extensive subcortical white-matter injury.

- Also termed subcortical arteriosclerotic encephalopathy.
- Vascular injury preferentially involves large areas of subcortical white matter.
- It represents a distinct distribution within vascular cognitive disease.
- Other causes of extensive hemispheric white-matter abnormalities must be excluded.

Binswanger pattern: Broad subcortical injury

Conceptual distribution map




Outer rim = cortex | Inner field = deep tissue

- The predominant injury involves large subcortical white-matter regions.
- This differs from a few discrete cortical infarcts.
- The schematic is conceptual and cannot substitute for clinical imaging.

Schematic — not an MRI, gross specimen, or mapped anatomy

Why white-matter lesions matter




- The final step is a clinicopathologic interpretation of disconnection in a multi-region syndrome.
- Exact clinical deficits still depend on lesion distribution.

Binswanger pattern: Differential principle

- Extensive hemispheric white-matter damage is not specific for vascular disease.
- The distribution requires correlation with the clinical course and vascular context.
- Both clinical and radiologic distinction from other white-matter diseases are essential.
- Do not diagnose the cause solely from a broad white-matter abnormality.

From ischemia to cell death




- Glutamate-mediated NMDA receptor activation allows excessive intracellular calcium entry.
- This mechanism contributes to vulnerability of ischemic CNS tissue.

Early morphology of cerebral infarction


- Acute ischemic neuronal injury produces eosinophilic “red neurons.”
- Nuclear condensation and fragmentation accompany irreversible neuronal damage.
- Glial and endothelial swelling, edema and myelin injury emerge.
- A remote infarct has a very different appearance from an acute lesion.

Infarct histology evolves with time



- The diagram summarizes Robbins descriptions of focal infarct evolution.
- Different regions of the same lesion may exhibit different stages.

Macrophage-mediated clearance of an infarct



- Phagocytes remove myelin debris and products of damaged tissue.
- Reactive astrocytes form a glial meshwork at the lesion edge.

Remote infarct: The neuropathologic endpoint

- Old ischemic lesions contain a cystic space left by lost parenchyma.
- A glial scar with capillaries rims the defect.
- Multiple remote cavities indicate an accumulated history of tissue destruction.
- The size and distribution of remote lesions provide anatomic context for the syndrome.

Chronic lacunes on histologic examination

- A lacune is a small remote cavity in a deep cerebral structure.
- Microscopy demonstrates tissue loss and reactive gliosis.
- The surrounding parenchyma may reflect chronic microvascular disease.
- Multiple lacunes should be considered together with coexisting white-matter injury.

Pale versus hemorrhagic infarction

Initially nonhemorrhagic infarct

- An occluded end artery deprives its downstream parenchyma of blood.
- Grossly a lesion becomes pale and soft after evolving ischemic injury.
- Cavitation appears as tissue is removed.

Hemorrhagic transformation

- Reperfusion may follow dissolution or fragmentation of occluding material.
- Damaged microvessels permit punctate or confluent bleeding.
- Underlying ischemia remains the initiating event.


Distribution is more informative than lesion label

- A lesion may be large, lacunar, cortical or subcortical.
- Multiple lesion types can accumulate in the same patient.
- Network disturbance depends on the combined location of tissue loss.
- The clinicopathologic assessment therefore integrates lesion types and distribution.

Core clinical triad

- Dementia developing over months or years.
- Gait abnormalities suggesting broader neural-network involvement.
- Pseudobulbar signs accompanying bilateral cerebral injury.
- Focal neurologic deficits may be superimposed.

Clinical manifestations and lesion geography



- A strictly focal stroke syndrome is not the same as the accumulated vascular dementia phenotype.
- The symptom combination reflects the distributed anatomical lesions.

Gait abnormalities: Clinicopathologic relevance

- Gait change is a characteristic associated feature in the Robbins account.
- Deep and subcortical tissue injury can disturb distributed motor pathways.
- Its coexistence with cognitive impairment increases concern for widespread brain involvement.
- A gait complaint alone does not establish vascular dementia.

Pseudobulbar signs: Bilateral network injury

- Pseudobulbar findings may accompany vascular dementia.
- They are compatible with interruption of bilateral descending supranuclear pathways.
- Their presence supports a broader lesion burden than an isolated cognitive complaint.
- Pseudobulbar signs do not uniquely identify the cause of vascular injury.

Focal deficits may be superimposed

- A focal deficit may arise with an additional vascular event.
- Different vascular territories yield different neurologic deficits.
- An acute deficit should not be interpreted as merely part of chronic dementia.
- Clinicopathologic interpretation requires both event timing and lesion location.

Clinical phenotype: Focal and distributed effects

Focal vascular effect

- Acute territorial or deep-structure insult.
- Deficit linked to a particular lesion location.
- Can coexist with chronic vascular cognitive impairment.

Distributed chronic effect

- Numerous bilateral lesions or extensive white-matter injury.
- Combined cognitive, gait and pseudobulbar findings.
- May reflect heterogeneous vascular mechanisms.

Clinical–radiologic correlation is indispensable

- Identify whether brain lesions are cortical, deep gray-matter or subcortical white-matter predominant.
- Examine whether the distribution plausibly relates to observed neurologic deficits.
- Consider large-artery, embolic and small-vessel substrates.
- The amount of visible vascular injury alone is not a complete attribution test.

Anatomical patterns and pathologic substrates

Observed pattern	Mechanism to consider	Interpretive caution
Multiple cortical infarcts	Large-artery or embolic events	Different events may coexist
Deep lacunar cavities	Penetrating small-vessel disease	Do not equate any lacune with dementia
Confluent subcortical injury	Diffuse white-matter vascular disease	Exclude other white-matter disorders
Lobar microbleeds	Cerebral amyloid angiopathy	Pattern differs from deep hypertension

Pattern-to-mechanism links reflect supplied Robbins pathology, not formal imaging criteria.


The Binswanger differential cannot be skipped

- Widespread white-matter disease has multiple possible causes.
- A vascular cause requires compatible history and lesion distribution.
- Clinical–radiologic correlation is explicitly emphasized by Robbins.
- A generic label of “white-matter change” is not a definitive etiology.

Mixed vascular and neurodegenerative pathology

- Cerebrovascular lesions often coexist with neurodegenerative disease.
- The additional vascular burden can increase neurologic impairment.
- For comparable degenerative lesion severity, vascular disease increases the risk of cognitive impairment.
- Avoid forcing patients with overlapping disease into a single exclusive category.

Overlapping pathologies and clinical impairment



- The source emphasizes the additive contribution of cerebrovascular pathology in degenerative disorders.

Schematic relationships (not quantitative)

Vascular and degenerative contributors

Vascular contribution

- Infarcts, lacunes and white-matter vascular injury.
- Spatial heterogeneity and focal deficits may be present.
- Gait and pseudobulbar features may coexist.

Degenerative contribution


- Neurodegenerative pathology may independently impair cognition.
- Often coexists with vascular abnormalities.
- Neurologic outcome reflects overlapping pathologic burdens.

Do not treat this table as a stand-alone diagnostic algorithm.

Common attribution errors

- Assuming all white-matter abnormalities signify Binswanger disease.
- Calling any isolated lacunar infarct sufficient evidence of vascular dementia.
- Attributing cognitive dysfunction exclusively to one pathologic process when lesions coexist.
- Ignoring a new focal deficit because a chronic cognitive syndrome is already present.

A structured clinicopathologic interpretation



- Correlate dementia, gait, pseudobulbar and focal findings.
- Keep morphologic observation separate from etiologic inference.

Case 1: Multifocal vascular brain injury

Clinical vignette

A patient has developed progressive cognitive impairment over several years. Examination identifies an abnormal gait and bilateral supranuclear motor signs. Brain studies show multiple remote cortical and deep infarcts.

Questions for interpretation

- Which pathologic syndrome is suggested?
- Why is the distribution of lesions important?

Case 1: Interpretation

- Multifocal bilateral infarction with dementia, gait and pseudobulbar features suggests vascular dementia.
- Both gray-matter and white-matter lesions may contribute.
- The differential vascular substrate includes atherosclerosis, embolism and hypertensive arteriolosclerosis.
- Additional clinical evidence is still required to establish the patient-level diagnosis.

Case 2: Subcortical disease

Clinical vignette

A patient with a history of chronic hypertension has slowly evolving cognitive and gait difficulties. Brain imaging demonstrates extensive injury preferentially affecting the subcortical white matter.

Questions for interpretation

- What named vascular pattern should be considered?
- What essential Diagnostic consideration: Remains?

Case 2: Interpretation

- The distribution suggests Binswanger disease (subcortical arteriosclerotic encephalopathy).
- Chronic hypertensive disease provides a plausible small-vessel substrate.
- The same extensive white-matter appearance can be seen in other disorders.
- Clinical and radiologic differentiation.

Case 3: Combined pathology

Clinical vignette

An older adult with established neurodegenerative brain disease also has several remote vascular infarcts and a greater burden of cognitive impairment than expected from degenerative lesions alone.


Questions for interpretation

- Does the presence of a degenerative disorder exclude vascular contribution?
- How should the coexisting infarcts be interpreted?

Case 3: Interpretation

- Neurodegenerative and vascular lesions may coexist.
- Cerebrovascular lesions can increase neurologic impairment for a given degenerative disease burden.
- The appropriate conclusion is combined neuropathologic contribution, not automatic exclusive attribution.
- This relationship is emphasized in the Robbins vascular dementia paragraph.

Integrative model of vascular dementia



- Three important upstream causes: Atherosclerosis, embolic/thrombotic disease and hypertensive small-vessel disease.
- Coexisting neurodegenerative pathology may amplify clinical impairment.

High-yield distinctions

Entity	Core feature	Important caveat
Vascular dementia	Dementia with multifocal vascular brain disease	Mixed degenerative pathology is common
Lacunar infarct	Small deep infarct; remote cavity / gliosis	A lesion type, not a syndrome
Binswanger disease	Extensive subcortical white-matter vascular injury	Other white-matter disorders can resemble it
CADASIL	Hereditary NOTCH3 small-vessel disease	One specific etiology, not all vascular dementia

Discussion and retrieval questions

- Which three principal vascular mechanisms are named in the Robbins vascular dementia section?
- What pathologic characteristic distinguishes a lacune from a large territorial infarct?
- What makes the Binswanger pattern distinctive, and why is its differential necessary?
- Why does coincident neurodegenerative pathology complicate disease attribution?

References and source boundaries

- Robbins, Cotran & Kumar: Pathologic Basis of Disease, 11th ed. (2026), ch. 28, “Vascular Dementia,” p. 1143.
- Related chapter sections: “Cerebrovascular Disease,” “Focal Cerebral Ischemia,” and “Intraparenchymal Hemorrhage,” pp. 1136–1141.
- All maps, vessel cross-sections and causal diagrams are original editable schematics; none depict a real specimen or scan.
- Source limitations: original chapter PDF pages and licensed diagnostic images were not included; radiologic criteria were not independently updated.