

Ventricular Septal Defect

Anatomy • shunt physiology • pulmonary vascular disease

Pathology teaching lecture

Dr Abu Suraih Sakhri


MBBS, MD (Pathology)

www.abusuraihsakhri.com

Learning objectives

- Localize membranous, muscular, and infundibular septal defects
- Explain the left-to-right shunt and the remodeling it produces
- Predict the natural history of small versus large defects
- Recognize the path to irreversible pulmonary vascular disease

Definition: Incomplete ventricular septation



Left ventricle · Septal opening · Right ventricle

Original simplified chamber schematic · not to anatomical scale

- A ventricular septal defect (VSD) is an opening in the interventricular septum
- It permits blood to communicate between right and left ventricles
- Its physiologic importance depends on defect size and accompanying malformations

VSD within congenital heart disease

Septal defects


Obstructive
lesions

Outflow
anomalies

Conceptual comparison · not a specimen or quantitative scale

- Congenital heart disease includes septal, obstructive, and outflow-tract defects
- VSD is a septal lesion and a common cause of congenital left-to-right shunting
- A VSD may be isolated or part of a complex malformation

VSD is the most common listed malformation



Selected congenital cardiac malformations · relative shares

- VSD as the most common congenital cardiac defect
- Its tabulated share is 42% of the defects in the cited series
- Frequency depends on how small, spontaneously closing defects are counted

The denominator changes the incidence


Screen-detected
openings

Persistent clinically
significant defects

Conceptual comparison · not a specimen or quantitative scale

- Routine neonatal echocardiography detects numerous tiny septal openings
- The source reports small muscular VSDs or ASDs in over 5% of live births
- Many close in infancy; Counting all of them inflates the burden of persistent CHD

Congenital origin: When malformations arise



Conceptual stages · temporal progression, not a patient timeline

- Major cardiovascular morphogenesis occurs during gestational weeks 3–8
- The four-chambered architecture develops through coordinated growth and septation
- Faulty closure can leave a persistent communication across the septum

Two developmental heart fields


First field → LV

Second field → RV /
outflow

Conceptual comparison · not a specimen or quantitative scale

- The first cardiac progenitor wave contributes to the left ventricle
- The second contributes to the right ventricle, outflow tract, and much of the atria
- Normal partitioning requires coordination between these developing structures

Septation completes four-chamber anatomy



Conceptual stages · temporal progression, not a patient timeline

- A primitive beating heart tube is established early in embryogenesis
- Cardiac chambers take shape before definitive septation is complete
- By roughly day 50, ventricular and atrial septation produces four chambers

Why septal development can fail

**Developmental
perturbation**



Incomplete
septation




VSD

Schematic causal sequence · arrows indicate the conceptual direction

- Cardiac morphogenesis requires cell migration, tissue remodeling, and coordinated gene expression
- Perturbations can affect localized septal structures without preventing live birth
- The precise cause of most congenital cardiac defects remains unknown


Etiology is genetically heterogeneous



Broad CHD associations · not specific causal assignments to VSD

- Both inherited factors and environmental exposures contribute to CHD
- No identifiable genetic defect is found in most affected patients
- Similar structural defects can result from different developmental disturbances

Interacting cardiac transcription factors



Interacting transcription factors

Interactions are contextual, not proof of a single VSD genotype

- GATA4, TBX5, and NKX2.5 in some septal-defect cases
- These factors physically interact and coregulate cardiac developmental targets
- A defect in the network may disrupt normal septum formation

GATA4: A nonsyndromic association

GATA4

Transcription factor

ASD / VSD

Associations derived from the CHD gene table

- The congenital heart disease gene table lists GATA4 with ASD or VSD
- GATA4 encodes a transcription factor
- This is an etiologic association—not a universal molecular explanation for VSD

TBX5: A syndromic association

TBX5

Holt–Oram syndrome

Septal / conduction findings

Associations derived from the CHD gene table

- TBX5 is listed with Holt–Oram syndrome
- ASD, VSD, or conduction defects among its cardiac manifestations
- VSD can therefore be one feature of a broader developmental syndrome

Other syndromes may include VSD

DiGeorge /
TBX1


CHARGE /
CHD7

Noonan / Ras-
pathway genes

Conceptual comparison · not a specimen or quantitative scale

- DiGeorge syndrome/22q11.2 deletion is associated with VSD among other defects
- CHARGE and Noonan syndromes also appear in the table of cardiac manifestations
- A VSD alone does not establish any of these syndromic diagnoses

Environmental contributors are nonspecific



Broad CHD associations · not specific causal assignments to VSD

- Congenital rubella, teratogens, and maternal diabetes among CHD causes
- Folate supplementation early in pregnancy is associated with lower CHD incidence
- These are broad CHD associations, not VSD-specific causal assignments

Isolated and associated VSDs



Isolated defect

Associated defect

Conceptual comparison · not a specimen or quantitative scale

- VSD can be an isolated opening or coexist with other cardiac malformations
- The source reports only 20%–30% of VSDs presenting in childhood are isolated
- Clinical interpretation therefore requires attention to additional structural lesions

Location and magnitude define morphology

Ventricular septal defect

Membranous


Muscular

Infundibular

Classified by site of the opening; Size is a separate descriptor

- VSDs by their position in the septum and defect magnitude
- Membranous lesions predominate; Muscular and infundibular lesions form the remainder
- Anatomic classification does not, by itself, specify shunt severity

Membranous VSD is the dominant subtype



Aortic valve region · Membranous septum · VSD opening

Original simplified chamber schematic · not to anatomical scale

- About 90% of VSDs are located in the membranous interventricular septum
- The membranous region lies near the aortic valve
- This high septal position differs from muscular and infundibular locations

The membranous defect: Spatial reasoning

Valve-level landmarks

VSD

Interventricular septum

Aortic valve • Upper septum

Regional schematic · positional relationships simplified

- The defect is high in the interventricular septum near the valve plane
- A high septal opening still connects the two ventricular chambers
- A precise gross specimen would be preferable to a schematic for operative localization

Infundibular VSD

Valve-level landmarks

VSD

Interventricular septum

Pulmonary valve • Infundibular region

Regional schematic · positional relationships simplified

- The source identifies an infundibular subtype below the pulmonary valve
- This location belongs to the minority of nonmembranous VSDs
- The pulmonary valve is a key landmark for recognizing the defect

Muscular VSD

Valve-level landmarks

Interventricular septum

VSD

Muscular septum • Single opening

Regional schematic • positional relationships simplified

- Muscular defects lie within the muscular interventricular septum
- They may be single or multiple
- Small muscular defects have a substantial potential for spontaneous closure

Multiplicity: A practical distinction



Single VSD

Multiple muscular
VSDs

Conceptual comparison · not a specimen or quantitative scale

- Most VSDs are solitary
- Multiple openings are especially encountered in the muscular septum
- Describing a defect therefore includes both location and number


Anatomic summary of VSD variants

Site	Anatomic relation	Pattern
Membranous	Near aortic valve	Most common
Infundibular	Below pulmonary valve	Less common
Muscular	Muscular septum	Can be multiple

Editable comparison based on Robbins descriptions

- Membranous: Around 90%, adjacent to the aortic valve region
- Infundibular: Located beneath the pulmonary valve
- Muscular: Located in muscular septum, sometimes multiple

The central pathophysiology is shunting




Systemic side · LV · VSD · Pulmonary side · RV

Original simplified chamber schematic · not to anatomical scale

- A shunt is an abnormal channel allowing blood to move between circulatory compartments
- With VSD, the defect connects the systemic and pulmonary ventricular pumps
- Flow direction reflects the pressure/resistance conditions of the circulation

An uncomplicated VSD begins left-to-right



Pathway abstraction · direction represents the mechanism described

- Systemic-side pressure ordinarily favors flow from left ventricle to right ventricle
- Oxygenated blood recirculates through the pulmonary circuit
- Initial left-to-right shunting does not itself produce cyanosis

Defect size alters the circulatory effect


Smaller VSD

Larger VSD

Conceptual comparison · not a specimen or quantitative scale

- The defect's magnitude influences the amount of shunted blood
- Associated right-sided malformations also modify the physiologic consequences
- A small opening may be tolerated; A large one may generate substantial load

Why left-to-right flow increases lung exposure



Schematic causal sequence · arrows indicate the conceptual direction

- Blood that crosses into the right ventricle is pumped back through pulmonary vessels
- The lungs receive extra flow that would not occur with an intact septum
- This alters the normally low-pressure, low-resistance pulmonary bed

The pulmonary circulation is low resistance

Normal pulmonary
bed

Chronic VSD shunt

Conceptual comparison · not a specimen or quantitative scale

- Normal pulmonary vessels are adapted to lower pressures than systemic vessels
- Significant shunts chronically increase pulmonary blood flow and pressure
- Progressive exposure initiates maladaptive arterial remodeling

Why large VSDs cause early difficulty


Large defect · early
impact

Small defect · delayed
detection

Conceptual comparison · not a specimen or quantitative scale

- A large defect permits a major left-to-right communication
- Clinically important problems virtually from birth in large VSDs
- Small lesions can remain unnoticed for many years


Right-ventricular pressure remodeling



Schematic causal sequence · arrows indicate the conceptual direction

- Large VSDs produce early right-ventricular hypertrophy
- The shunt-associated pulmonary pressure load contributes to this response
- Right-ventricular hypertrophy is a structural sign of hemodynamic burden

Left-ventricular volume remodeling



Pathway abstraction · direction represents the mechanism described

- The source figure describes left-ventricular volume hypertrophy in VSD
- Recirculated pulmonary blood returns to left-sided cardiac chambers
- This creates a volume component of the ventricular adaptation

Pressure and volume effects coexist


RV · pressure loading

LV · volume loading

Conceptual comparison · not a specimen or quantitative scale

- A substantial VSD affects more than one chamber
- Right ventricular pressure hypertrophy and left ventricular volume hypertrophy can coexist
- The relative dominance depends on the hemodynamic context


Pulmonary arteries respond initially



Schematic causal sequence · arrows indicate the conceptual direction

- Increased pulmonary flow and pressure stimulate medial hypertrophy
- Small muscular pulmonary arteries also vasoconstrict
- These early adaptations help limit downstream capillary pressure


Long-term vasoconstriction becomes damaging



Schematic causal sequence · arrows indicate the conceptual direction

- Persistent vasoconstriction is not a benign steady state
- Prolonged stress promotes obstructive lesions in the arterial intima
- The remodeled vascular bed loses its normal low-resistance character


Pulmonary vascular disease is progressive



Conceptual stages · temporal progression, not a patient timeline

- Large uncorrected defects may progress to irreversible obstructive pulmonary vascular disease
- Increasing vascular resistance progressively raises the pressure seen by the right ventricle
- The shunt changes as the pressure relationship between ventricles changes


Shunt reversal is a late transition



Pathway abstraction · direction represents the mechanism described

- When pulmonary resistance approaches systemic resistance, shunt direction can reverse
- Venous blood is then delivered to the systemic arterial circulation
- Cyanosis arises from systemic admixture of poorly oxygenated blood

Eisenmenger syndrome



Schematic causal sequence · arrows indicate the conceptual direction

- Eisenmenger syndrome is reversal of a longstanding left-to-right shunt
- The mechanism is acquired severe pulmonary vascular disease
- It signals that the original congenital shunt has reached an advanced pathologic stage

Why cyanosis is absent initially

Early · left-to-right




Late · right-to-left

Conceptual comparison · not a specimen or quantitative scale

- Early left-to-right shunting does not bypass oxygenation of venous blood
- Therefore initial isolated VSD is not a cyanotic defect
- Cyanosis after reversal reflects admixture of venous blood into systemic blood


Consequences of chronic hypoxemia



Schematic causal sequence · arrows indicate the conceptual direction

- Persistent cyanotic shunts with increased red-cell counts
- Digital clubbing and hypertrophic osteoarthropathy may occur with prolonged hypoxia
- These consequences belong to the advanced right-to-left phase


Paradoxical embolism becomes possible



Pathway abstraction · direction represents the mechanism described

- Right-to-left channels allow venous emboli to bypass pulmonary filtration
- Such emboli can enter the systemic arterial supply
- This is a consequence of shunt direction rather than a defining feature of every VSD


Pathology connects defect and vasculature



Conceptual stages · temporal progression, not a patient timeline

- Septal anatomy establishes an abnormal route for blood
- The resulting load changes chamber morphology and pulmonary arterioles
- Late vascular injury can become more decisive than the septal opening itself

Associated defects change shunt physiology



Schematic causal sequence · arrows indicate the conceptual direction

- Robbins stresses the importance of right-sided accompanying malformations
- Right-ventricular outflow obstruction may alter transseptal pressure conditions
- Anatomic context must be defined before predicting the shunt

Tetralogy of Fallot includes a VSD

Isolated VSD

Tetralogy with
obstruction

Conceptual comparison · not a specimen or quantitative scale

- Tetralogy combines VSD with subpulmonary obstruction, overriding aorta, and right-ventricular hypertrophy
- The direction of shunting depends on the degree of right-ventricular outflow obstruction
- This contrasts with the usual isolated left-to-right VSD

VSD versus atrial septal defect

Property	VSD	ASD
Opening	Ventricular	Atrial
Pulmonary load	Flow + pressure	Mainly flow
Clinical pattern	Size-dependent	Often later presentation

Editable comparison based on Robbins descriptions

- VSD communicates between ventricles; ASD communicates between atria
- VSD as adding pulmonary blood flow and pressure
- ASD predominantly adds right ventricular and pulmonary outflow volume

VSD versus patent ductus arteriosus

Property	VSD	PDA
Connection	LV ↔ RV	Aorta ↔ pulmonary artery
Early flow	Left-to-right	Left-to-right
Pulmonary burden	Flow + pressure	Flow + pressure

Editable comparison based on Robbins descriptions

- VSD is an interventricular communication
- PDA is persistence of the channel between pulmonary artery and aorta
- Both can cause increased pulmonary blood flow and pressure

The small VSD phenotype

Small opening


Limited burden

Late
recognition

Conceptual comparison · not a specimen or quantitative scale

- Smaller defects are often well tolerated for many years
- They may not be recognized until later in life
- This course differs from clinically important large defects present early


Small muscular VSDs can close



Schematic causal sequence · arrows indicate the conceptual direction

- Spontaneous closure in approximately 50% of small muscular VSDs
- Closure potential is a major reason not to treat all asymptomatic openings immediately
- This estimate is for the small muscular category—not all VSDs


Adult-detected VSD has a different profile



Conceptual stages · temporal progression, not a patient timeline

- A VSD first identified in adulthood is usually isolated,
- The survival of an undetected defect suggests a relatively tolerated circulatory impact
- Timing of discovery should not be confused with timing of embryologic origin


Large untreated VSD has a different course



Schematic causal sequence · arrows indicate the conceptual direction

- Substantial shunting increases pulmonary flow and pressure
- Early RV hypertrophy and pulmonary hypertension are common consequences
- Irreversible pulmonary arteriopathy becomes a major long-term hazard


Observation is conditional—not universal



Teaching schematic, not a clinical management algorithm

- Deferring closure for asymptomatic VSD beyond infancy
- The rationale is the potential for spontaneous closure
- This approach must not be generalized to large hemodynamically important defects


Why significant defects require early repair



Schematic causal sequence · arrows indicate the conceptual direction

- Early correction prevents continued pulmonary vascular stress
- The stated goal is to prevent irreversible obstructive pulmonary vascular disease
- Timing depends on lesion significance rather than congenital diagnosis alone

Late vascular disease changes repairability



Conceptual stages · temporal progression, not a patient timeline

- Once pulmonary vascular disease is irreversible, simple closure is no longer an adequate solution
- Such congenital shunt lesions as irreparable at that stage
- Therefore natural history determines the opportunity for corrective intervention

Morphology versus clinical severity

Anatomic description

— Physiologic assessment

Conceptual comparison · not a specimen or quantitative scale

- Anatomic location specifies the septal defect
- Size and associated right-sided malformations dominate the functional consequences
- Pulmonary vascular resistance and shunt direction describe advanced physiologic progression

Practical clinicopathologic questions

- Is the defect single or multiple, and where is it located?
- How large is the hemodynamic communication?
- Are there accompanying right-sided malformations?
- Is pulmonary vascular disease developing or established?

Case 1: An incidental muscular opening

Clinical reasoning vignette

Small muscular VSD


No large shunt

Possible closure

Illustrative hypothetical case; No patient data

- A newborn screening study identifies a tiny muscular VSD
- No large-shunt manifestations are described
- Question: Which outcome discussed by Robbins is plausible during infancy?

Case 1: Interpret the prognosis



Schematic causal sequence · arrows indicate the conceptual direction

- Spontaneous closure is plausible, especially for small muscular defects
- The cited approximate 50% closure frequency belongs to this subgroup
- The scenario does not justify assuming every small lesion will close

Case 2: A large infant VSD

Clinical reasoning vignette

Cardiac chambers


Pulmonary arteries

Time-dependent changes

Illustrative hypothetical case; No patient data

- An infant has a large interventricular communication
- Evidence of significant pulmonary flow burden is present
- Question: Which two pathologic compartments are at greatest risk over time?

Case 2: Follow the remodeling



Schematic causal sequence · arrows indicate the conceptual direction

- The right ventricle develops hypertrophy under pressure load
- Pulmonary arteries undergo medial and eventually intimal remodeling
- Uncorrected shunting risks fixed pulmonary vascular obstruction

Case 3: Late-onset cyanosis

Clinical reasoning vignette

Prior left-to-right


Pulmonary resistance ↑

Flow reversal

Illustrative hypothetical case; No patient data

- An unrepaired large VSD persists into later life
- Pulmonary vascular resistance is now markedly elevated
- Question: Why can a previously acyanotic defect become cyanotic?

Case 3: Identify Eisenmenger physiology



Schematic causal sequence · arrows indicate the conceptual direction

- Fixed pulmonary vascular remodeling reverses the original shunt
- Right-to-left admixture lowers systemic arterial oxygenation
- The case illustrates the importance of repair before irreversible vascular disease


Three natural histories to distinguish

Defect context	Flow / course	Key concern
Small muscular	May close	Follow natural history
Large unclosed	Large L→R shunt	Pulmonary arteriopathy
VSD with obstruction	Flow may alter	Cyanotic physiology

Editable comparison based on Robbins descriptions

- Small muscular VSD: Possible spontaneous closure
- Large VSD: Substantial early loading and pulmonary vascular risk
- VSD with right-sided obstruction: Altered flow direction, as in tetralogy

Integrated sequence: From anatomy to outcome



Conceptual stages · temporal progression, not a patient timeline

- A structural defect creates communication between ventricular chambers
- Shunting increases pulmonary load and stimulates ventricular/vascular adaptation
- The outcome depends on defect size, associated anomalies, and progression of vascular disease

References and source boundaries

- Robbins, Cotran & Kumar, Pathologic Basis of Disease, 11th edition, Chapter 12, pp. 485–490
- Primary topic: “Ventricular Septal Defect,” pp. 488–489
- Supporting context: congenital heart disease, left-to-right shunts, and tetralogy of Fallot
- Original simplified PowerPoint schematics; no published figures reproduced